

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/03917 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US01/21701
- (22) International Filing Date: 9 July 2001 (09.07.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/216,995 7 July 2000 (07.07.2000) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/216,995 (CIP)
Filed on 7 July 2000 (07.07.2000)
- (71) Applicants (for all designated States except US): **AL-PHAX, INC.** [US/US]; P.O. Box 110307, 2 Triangle Drive, Research Triangle Park, NC 27709-0307 (US). **UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL** [US/US]; 308 Bynum Hall, Campus Box 4105, Chapel Hill, NC 27599-4105 (US).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): **OLMSTED, Robert** [US/US]; 420 Bayberry Drive, Chapel Hill, NC 27514 (US). **KEITH, Paula** [US/US]; 5004 Sadelia Place, Holly Springs, NC 27540 (US). **DRYGA, Sergey** [RU/US]; 211 Bonsill Place, Chapel Hill, NC 27514 (US). **CALEY, Ian** [GB/US]; 2 Triangle Drive, P.O. Box 110307, Research Triangle Park, NC 27709-0307 (US). **MAUGHAN, Maureen** [US/US]; 2532 Wrightwood Avenue, Durham, NC 27705 (US). **JOHNSTON, Robert** [US/US]; 101 Marin Place, Chapel Hill, NC 27516 (US). **DAVIS, Nancy** [US/US]; 132 New Castle Drive, Chapel Hill, NC 27514 (US). **SWANSTROM, Ronald** [US/US]; 7021 Knotty Pine Drive, Chapel Hill, NC 27514-8659 (US).
- (74) Agents: **MILLER, Mary, L.** et al.; Needle & Rosenberg, P.C., 127 Peachtree Street, N.E., Suite 1200, Atlanta, GA 30303-1811 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IL, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/03917 A2

(54) Title: ALPHAVIRUS VECTORS AND VIROSOMES WITH MODIFIED HIV GENES FOR USE AS VACCINES

(57) Abstract: The present invention provides methods and compositions comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit integrase, RNase H and/or reverse transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle.

**ALPHAVIRUS VECTORS AND VIROSOMES
WITH MODIFIED HIV GENES FOR USE AS VACCINES**

This application claims priority to U.S. Provisional Application No. 60/216,995,
5 filed July 7, 2000 which is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

10

The present invention relates to vaccines using viral antigens, and in particular,
to vaccines for the treatment and prevention of human immunodeficiency virus (HIV)
infection. The vaccines of this invention comprise alphavirus RNA replicon systems
which contain nucleic acid sequence encoding antigens for eliciting an immune
15 response to HIV.

Background

The successful control of the AIDS epidemic will require an effective vaccine
20 for human immunodeficiency virus type 1 (HIV) that significantly reduces or prevents
the spread of infection. Currently, several viral vector systems as well as naked DNA
are at various stages of pre-clinical and clinical evaluation as candidate HIV vaccines.
Recombinant poxviruses are the most widely studied virus vectors and are furthest
along in clinical development (e.g., ALVAC).

25

The alphavirus-based replicon particle systems, such as the ones described in
U.S. Patent No. 5,792,462 and herein referred to as "VRPs," have multiple distinct
properties that make them attractive as an HIV vaccine delivery technology. These
properties include: natural targeting to and expression in lymphoid tissues (an optimal
30 site for induction of an immune response); high antigen expression levels, e.g., up to
20% of total cell protein; induction of balanced humoral, cellular, and mucosal immune
responses; sustained efficacy over multiple simultaneous or sequential inoculations of

the vector; and a high margin of safety.

Venezuelan equine encephalitis virus (VEE) is a member of the Alphaviruses group, which also includes the prototype Sindbis virus (SIN) and Semliki Forest virus (SFV), and is comprised of enveloped viruses containing plus-stranded RNA genomes within icosahedral capsids (Strauss, 1994). Alphavirus genomes are: approximately 11.5 kb long, capped, polyadenylated, and infectious under appropriate transfection conditions. The nucleocapsid is composed of 240 molecules of the capsid protein arranged as a T=4 icosahedron, and is surrounded by a lipoprotein envelope (Paredes *et al.*, 1993). Protruding from the virion surface are 80 glycoprotein spikes, each of which is a trimer of virally encoded E1 and E2 glycoprotein heterodimers. The virions contain no host proteins.

Alphaviruses share replication strategies and genomic organization. The complete replicative cycle of alphaviruses occurs in the cytoplasm of infected cells. Expression from the alphavirus genome is segregated into two regions. The four enzymatic nonstructural proteins (nsP1-nsP4) are synthesized from the 5' two-thirds of the genome-length RNA and are required for RNA replication. Immediately following infection, the nsPs are produced by translation of parental genomes and catalyze the synthesis of a full-length negative-sense copy of the genome. This serves as a template for the synthesis of progeny plus-stranded genomes.

The negative-sense copy of the genome also serves as the template for the synthesis of subgenomic mRNA at approximately 10-fold molar excess relative to genomic RNA in infected cells (Schlesinger and Schlesinger, 1990). Synthesis of subgenomic 26S mRNA is initiated from the highly active internal 26S mRNA promoter, which is functional only on the negative-sense RNA. The subgenomic mRNA corresponds to the 3' one-third of the genome and encodes the alphavirus structural proteins.

Full-length, infectious cDNA clones of the RNA genome of VEE (Davis *et al.*, 1989) have been constructed, a panel of mutations which strongly attenuate the virus have been identified (Johnston and Smith, 1988; Davis *et al.*, 1990), and various constellations of these attenuating mutations have been inserted into the clones to
5 generate several live attenuated VEE vaccine candidates (Davis *et al.*, 1991; 1995b; Grieder *et al.*, 1995). The resulting vaccine candidates are avirulent and provide complete protection against lethal virus challenge in rodents, horses and nonhuman primates.

10 The alphavirus VRPs are propagation defective, single cycle vectors that contain a self-amplifying alphavirus RNA (replicon RNA) in which the structural protein genes of the virus are replaced by a heterologous antigen gene to be expressed. Alphavirus VRPs are typically made in cultured cells, referred to as packaging cells. Following introduction into mammalian cells, the replicon RNA is packaged into VRP
15 by supplying the structural proteins in "trans," i.e. the cells are co-transfected with both replicon RNA and one or more separate helper RNAs which together encode the full complement of alphavirus structural proteins. Importantly, only the replicon RNA is packaged into VRP, as the helper RNA(s) lack the *cis*-acting packaging sequence required for encapsidation. Thus, the VRPs are defective, in that they can only infect
20 target cells in culture or *in vivo*, where they express the heterologous antigen gene to high level, but they lack critical portions of the VEE genome (i.e., the VEE structural protein genes) necessary to produce virus particles which could spread to other cells.

Delivery of the replicon RNA into target cells (for vaccination) is facilitated by
25 the VRP following infection of the target cells. In the cytoplasm of the target cell, the replicon RNA is first translated to produce the viral replicase proteins necessary to initiate self-amplification and expression. The heterologous antigen gene is encoded by a subgenomic mRNA, abundantly transcribed from the replicon RNA, leading to high level expression of the heterologous antigen gene product. Since the VEE structural
30 protein genes are not encoded by the replicon RNA delivered to the target cell, progeny

virion particles are not assembled, thus limiting the replication to a single cycle within the infected target cell. Experimental VRP vaccines have been successful in vaccinating rodents against influenza virus, Lassa fever virus and Marburg virus (Pushko *et al.*, 1997; Hevey *et al.*, 1998). In nonhuman primates, VRP vaccines have demonstrated complete efficacy against lethal Marburg virus challenge (Hevey *et al.*, 1998), shown partial but significant protection against SIV infection and disease (Davis *et al.*, 2000) and induced an anti-HA response at a level consistent with protection of humans against influenza virus infection.

The alphavirus based replicon vector systems, and in particular the VEE-based systems, present several advantages in vaccination, including safety and high immunogenicity/efficacy. VEE is unique among the alphaviruses in that a live attenuated IND VEE vaccine, TC-83, (Kinney *et al.*, 1989; Kinney *et al.*, 1993) has been inoculated into approximately 8,000 humans. This allows direct safety and efficacy comparisons between human, nonhuman primate and rodent responses to the same VEE derivative. A large body of experience strongly suggests that the animal models generally reflect the human susceptibility and disease course, except that mice are far more susceptible to lethal VEE disease than humans or nonhuman primates. Furthermore, the VEE replicon vectors express high levels of the gene of interest in cell culture, and *in vivo* expression is targeted to lymphoid tissues, reflecting the natural tropism mediated by the VEE glycoproteins. Cells in the draining lymph node of VRP-inoculated mice contain detectable amounts of the desired gene product within hours of inoculation. This expression continues for up to five days.

To date, VRP vector vaccines have been used in over 2000 rodents and in 94 macaques at doses up to 5×10^8 i.u., with no indication of any clinical manifestations.

In work reported by Pushko *et al.* (1997), individual mice were immunized sequentially with Lassa virus N-VRP and influenza virus HA-VRP. Groups of mice, which received two inoculations of 3×10^4 or 3×10^6 i.u. of Lassa N-VRP followed by

two inoculations of 2×10^5 i.u. of HA-VRP, all responded with serum antibodies to both antigens. The level of anti-influenza antibody induced in these sequentially inoculated mice was equivalent to a control group, which received two inoculations of buffer followed by two inoculations of 2×10^5 i.u. of HA-VRP. All HA-VRP immunized mice were completely protected against influenza virus challenge. Furthermore, sequential immunization of mice with two inoculations of N-VRP prior to two inoculations of HA-VRP induced an immune response to both HA and N equivalent to immunization with either VRP construct alone. Primary and booster immunization with a VRP preparation expressing an immunogen from one pathogen did not interfere with the development of a protective response to subsequent primary immunization and boosting with VRP expressing an immunogen from a second pathogen, thus showing that the VRP-based system can be used to induce immunity to a variety of pathogens in the same individual over time.

Four macaques were inoculated subcutaneously at week 0 with 10^5 i.u. each of SIV-gp160-VRP (*env*) and SIV MA/CA-VRP (*gag*), boosted by the same route at week 7 with 10^7 i.u. of each VRP vaccine, and intravenously at weeks 12 and 20 with 5×10^8 i.u. of each VRP. Two control animals were inoculated with equivalent doses of HA-VRP (haemagglutinin, a glycoprotein from influenza virus), and two with the vehicle only. The four SIV-VRPs immunized monkeys received subcutaneously an additional dose of 2×10^7 i.u. of gp140-VRP at week 41, followed by a final boost of 2×10^7 i.u. each of gp140-VRP and MA/CA-VRP at week 49. Four weeks after the final immunization, all eight macaques were challenged intravenously with the pathogenic virus, SIVsmE660.

25

After these inoculations, three of four test macaques had measurable CTL-specific killing directed against both SIV *gag* and *env*, all four had gp160 IgG antibody by ELISA, and the three animals which harbored SIV-specific CTL also showed neutralizing antibody to SIVsmH-4.

30

Four of four vaccinated animals were protected against disease for at least 16 months following intravenous challenge with the pathogenic SIV swarm, while the two vehicle controls required euthanasia at week 10 and week 11, post challenge. In two of the vaccinees, plasma virus levels were below the limit of detection by branched chain
5 DNA assay. At 64 weeks post challenge, all four vaccinated animals showed no clinical signs of disease. One animal remained vDNA negative at 64 weeks.

The results of this highly pathogenic challenge demonstrated that the immune response induced by vaccination with SIV-VRP was effective in preventing early
10 mortality and increasing the ability to suppress challenge virus replication. The ability to control SIV replication and reduce viral load to undetectable levels was closely correlated with the strongest measurable antibody and cellular immune responses.

While these results are encouraging, the level of protection obtained would not
15 be acceptable for a human vaccine against HIV infection. Thus, there remains a need for a robust, effective and safe vaccine against HIV infection in humans. Development of a HIV vaccine comprising the complete, or immunogenic fragments of the, *gag* gene (Gag-VRP), an immunogenic portion of the *pol* gene (Pol-VRP), and the complete, or immunogenic fragments of the, *env* gene (Env-VRP), would increase the diversity of
20 available CTL epitopes substantially and thus address this need.

SUMMARY OF THE INVENTION

The present invention provides a composition comprising two or more isolated
25 nucleic acids selected from the group consisting of an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles
30 containing the *gag* gene product or the immunogenic fragment thereof and their release

from a cell, and an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity.

5

Also provided is a composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated
10 nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic
15 fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle.

20 In addition, the present invention provides a composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment
25 thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol*
30 gene product or immunogenic fragment thereof is modified to inhibit reverse

transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle, and further wherein the alphavirus replicon particles comprise a replicon RNA or at least one structural protein which comprises one or more attenuating mutations.

5

10. A method of making a population of alphavirus replicon particles of this invention is provided herein, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

10

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

15

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

20

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

25

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

30

(b) producing the alphavirus particles in the helper cell; and

- (c) collecting the alphavirus particles from the helper cells;
- B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particles, comprising in an alphavirus-
- 5 permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic
- 10 fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first
- 15 helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional
- 20 helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper
- 25 RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
- 30 (c) collecting the alphavirus particles from the helper cells;

- C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells; and
- D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second

helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles.

Also provided is a method of making a population of alphavirus replicon

5 particles, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

10

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

15

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

20

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

25

30

(b) producing the alphavirus particles in the helper cell; and

(c) collecting the alphavirus particles from the helper cells;

B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particle, comprising in an alphavirus-

5 permissive cell:

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

10

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

15

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

20

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper

25 RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or
30 more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

- C) (a) providing a third helper cell for producing a third population of
5 infectious, replication defective alphavirus particles, comprising in an alphavirus-
permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA
comprises an alphavirus packaging signal and a nucleic acid encoding a
10 *pol* gene product or an immunogenic fragment thereof of a human
immunodeficiency virus, wherein the *pol* gene product or immunogenic
fragment thereof is modified to inhibit reverse transcriptase activity, and
wherein the replicon RNA lacks sequences encoding alphavirus
structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first
15 helper RNA encoding at least one alphavirus structural protein and
furthermore not encoding at least one other alphavirus structural protein;
and
 - (iii) one or more additional helper RNA(s) separate from said
20 replicon RNA and separate from said first helper RNA, said additional
helper RNA(s) encoding at least one other alphavirus structural protein
not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging
signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper
25 RNAs produces an assembled alphavirus particle which is able to infect a cell, and is
unable to complete viral replication, and further wherein the third population contains
no detectable replication-competent alphavirus particles as determined by passage on
permissive cells in culture, and further wherein at least one of said replicon RNA, said
first helper RNA, and said one or more additional helper RNA(s) comprises one or
30 more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first
5 helper cell, the second population of alphavirus particles produced from the second
helper cell and the third population of alphavirus particles produced from the third
helper cell, thereby producing the population of alphavirus replicon particles.

Furthermore, the present invention provides a composition comprising two or
10 more isolated nucleic acids selected from the group consisting of an isolated nucleic
acid encoding an *env* gene product or an immunogenic fragment thereof of a human
immunodeficiency virus, an isolated nucleic acid encoding a *gag* gene product or an
immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag*
15 gene product or immunogenic fragment thereof is modified to inhibit formation of
virus-like particles containing the *gag* gene product or the immunogenic fragment
thereof and their release from a cell, and an isolated nucleic acid encoding a *pol* gene
product or an immunogenic fragment thereof of a human immunodeficiency virus,
wherein the *pol* gene product or immunogenic fragment thereof comprises a
20 modification resulting in deletion or inactivation of integrase, RNase H and reverse
transcriptase functions in the *pol* gene product or immunogenic fragment thereof.

In addition, the present invention provides a composition comprising a
population of alphavirus replicon particles comprising two or more isolated nucleic
acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env*
25 gene product or an immunogenic fragment thereof of a human immunodeficiency virus,
2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment
thereof of a human immunodeficiency virus, wherein the *gag* gene product or
immunogenic fragment thereof is modified to inhibit formation of virus-like particles
containing the *gag* gene product or the immunogenic fragment thereof and their release
30 from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an

- immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are
- 5 each contained within a separate alphavirus replicon particle.

- Also provided herein is a composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an
- 10 immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell,
- 15 and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each
- 20 contained within a separate alphavirus replicon particle, and further wherein the alphavirus replicon particles comprise a replicon RNA or at least one structural protein which comprises one or more attenuating mutations.

- In these embodiments, the *gag* gene product or immunogenic fragment thereof
- 25 can be modified by mutation of the second codon, whereby a glycine is changed to an alanine and the *pol* gene product or immunogenic fragment thereof can be modified by mutation of the nucleotide sequence encoding the active site motif, whereby YMDD is changed to YMAA or HMAA. In addition, the *pol* gene product or immunogenic fragment thereof is modified to produce only p51 of the *pol* gene product or
- 30 immunogenic fragment thereof.

The present invention provides a method of making a population of alphavirus replicon particles, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- 5 (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- 10 (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- 15 (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

- 20 wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- 25 (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

- B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particles, comprising in an alphavirus-
30 permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;
- C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA

- comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or
- 5 inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and
- 10 furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein
- 15 not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is
- 20 unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and
- 25 D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles.
- 30

An additional method of making a population of alphavirus replicon particles is provided, comprising:

- A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:
- 5 (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- 10 (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- 15 (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- 20 wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said
- 25 first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;
- 30 B) (a) providing a second helper cell for producing a second population of

infectious, replication defective alphavirus particle, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells;

C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- 5 (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and
10 wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and
15 furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein
20 not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is
25 unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- 30 (b) producing the alphavirus particles in the helper cell; and

(c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles.

In each of the methods above, the alphavirus replicon RNA of at least one of the first helper cell, the second helper cell and the third helper cell can comprise sequence encoding at least one alphavirus structural protein and the first helper RNA and the one or more additional helper RNA(s) in the at least one of the first helper cell, the second helper cell and the third helper cell, can encode at least one other alphavirus structural protein not encoded by the replicon RNA.

Furthermore, in the methods above which recite attenuating mutations, only at least one of the first population of alphavirus particles, the second population of alphavirus particles and the third population of alphavirus particles can comprise particles wherein at least one of the replicon RNA, the first helper RNA, and the one or more additional helper RNA(s) comprises one or more attenuating mutations.

The present invention further provides alphavirus particles produced by any of the methods of this invention.

The present invention further provides a method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the populations and/or compositions of this invention, in a pharmaceutically acceptable carrier.

Also provided herein is a method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an

immunogenic amount of the populations and/or compositions of this invention, in a pharmaceutically acceptable carrier.

Also provided by the present invention is an alphavirus replicon virosome
5 comprising an alphavirus replicon RNA encapsidated by a lipid bilayer comprising alphavirus glycoproteins, E1 and E2, which in one embodiment, can be Venezuelan Equine Encephalitis glycoproteins E1 and E2.

A method of producing an alphavirus replicon virosome is further provided,
10 comprising: a) combining alphavirus replicon RNA, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and b) gradually removing detergent, whereby alphavirus replicon virosomes are produced. Also provided is a virosome produced by this method.

Furthermore, the present invention provides a method of eliciting an immune
15 response in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon virosome of this invention in a pharmaceutically acceptable carrier.

The present invention additionally provides a method of treating or preventing
20 infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon virosome of this invention, wherein the virosome comprises alphavirus replicon RNA encoding one or more HIV immunogens.

25

In further embodiments, the present invention provides a composition a
population of alphavirus replicon virosomes comprising two or more isolated nucleic
acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env*
gene product or an immunogenic fragment thereof of a human immunodeficiency virus,
30 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment

thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon virosome.

10

Additionally provided herein is a composition comprising a population of alphavirus replicon virosomes comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in inactivation of reverse transcriptase activity in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon virosome.

25

A method of producing a population of alphavirus replicon virosomes is provided herein, comprising:

- A) (a) producing a first population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *env* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-

30

cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;

5

B) (a) producing a second population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *gag* gene product or immunogenic fragment thereof, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

10

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;

15

C) (a) producing a third population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding the *pol* gene product or immunogenic fragment thereof, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

20

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced; and

25

D) combining the first population of alphavirus replicon virosomes, the second population of alphavirus replicon virosomes and the third population of alphavirus replicon virosomes to produce the population of alphavirus replicon virosomes.

30

In addition, a method of producing a population of alphavirus replicon virosomes is provided, comprising:

- 5 A) (a) producing a first population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *env* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and
- b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;
- 10 B) (a) producing a second population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *gag* gene product or immunogenic fragment thereof, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles
- 15 containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and
- b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;
- 20 C) (a) producing a third population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding the *pol* gene product or immunogenic fragment thereof, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in inactivation of
- 25 reverse transcriptase activity in the *pol* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and
- b) gradually removing detergent, whereby alphavirus replicon virosomes are produced; and
- D) combining the first population of alphavirus replicon virosomes, the second
- 30 population of alphavirus replicon virosomes and the third population of alphavirus

replicon virosomes to produce the population of alphavirus replicon virosomes of claim 48.

Furthermore, the present invention provides a method of inducing an immune
5 response in a subject, comprising administering to the subject an immunogenic amount of the virosomes of this invention, in a pharmaceutically acceptable carrier.

Also provided is a method of treating or preventing infection by human
immunodeficiency virus in a subject, comprising administering to the subject an
10 immunogenic amount of the virosomes of this invention, in a pharmaceutically acceptable carrier.

Additionally provided by this invention is a composition comprising heparin
affinity-purified alphavirus replicon particles, wherein the alphavirus replicon particles
15 comprise at least one structural protein which comprises one or more attenuating mutations, as well as a method of preparing heparin affinity-purified alphavirus particles, comprising:

- a) producing alphavirus replicon particles, wherein the alphavirus replicon particles comprise a at least one structural protein which comprises one or
20 more attenuating mutations;
- b) loading the alphavirus replicon particles of step (a) in a heparin affinity chromatography column; and
- c) collecting the fraction from the column which contains the heparin affinity-purified alphavirus replicon particles.

25

In further embodiments, the present invention provides a method of producing VRP for use in a vaccine comprising:

- a) producing a plasmid encoding the nucleotide sequence of an alphavirus replicon RNA;
- 30 b) producing a plasmid encoding the nucleotide sequence of one or more

helper RNAs;

c) transcribing the plasmids of steps (a) and (b) into RNA *in vitro*;

d) electroporating the RNA of step (c) into a Vero cell line; and

5

e) purifying VRP from the Vero cell line of step (d) by heparin affinity chromatography. By this method, VRPs can be produced in large-scale.

- In additional embodiments, the present invention provides an isolated nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof. This nucleic acid can be present in a composition and in a vector.
- 10 Such a vector can be present in a cell. This nucleic acid can also be present in an alphavirus replicon particle.

- The present invention further provides a method of making an alphavirus replicon particle comprising nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, comprising
- 20 a) providing a helper cell for producing an infectious, defective alphavirus particle, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or
- 30

- inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cell.

In the method described above, at least one of the replicon RNA, the first helper RNA, and the one or more additional helper RNA(s) can comprises one or more attenuating mutations. The present invention additionally provides alphavirus replicon particle produced according to the above methods.

Further provided is a method of inducing an immune response in a subject, comprising administering to the subject an immunogenic amount of a composition comprising alphavirus replicon particles encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol*

gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof in a pharmaceutically acceptable carrier.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. DNA plasmid map of VEE replicon RNA expressing the HIV *gag* gene (p3-40.1.6). The plasmid is 12523 base pairs in length and encodes a single
10 polypeptide expressing the four non-structural genes nsP1-4, the Clade C *gag* gene and antibiotic resistance marker, Kanamycin KN(R). The plasmid contains two promoter regions, the T7 polymerase promoter and the 26S RNA promoter. The unique NotI restriction enzyme site used to linearize prior to *in vitro* transcription is also noted.

15 Figure 2. DNA plasmid map of the capsid helper construct (p3-13.2.2). The plasmid is 5076 base pairs in length and encodes the VEE capsid gene (C) and antibiotic resistance marker, Kanamycin KN(R). The plasmid contains two promoter regions, the T7 polymerase promoter and the 26S RNA promoter. The unique NotI restriction enzyme site used to linearize prior to *in vitro* transcription is also noted.

20

Figure 3. DNA plasmid map of the glycoprotein helper construct (p3-13.4.6). The plasmid is 6989 base pairs in length and encodes the VEE glycoprotein genes (E3, E2, 6K and E1) and antibiotic resistance marker, Kanamycin KN(R). The plasmid contains two promoter regions, the T7 polymerase promoter and the 26S RNA
25 promoter. The unique NotI restriction enzyme site used to linearize prior to *in vitro* transcription is also noted.

Figure 4. DNA plasmid map of VEE replicon RNA expressing HIV *pol* (p51) gene (p13-60.2.14). The plasmid is 12379 base pairs in length and encodes a single
30 polypeptide expressing the four non-structural genes, nsP1-4, the Clade C *pol* (p51)

gene and antibiotic resistance marker, Kanamycin KN(R). The plasmid contains two promoter regions, the T7 polymerase promoter and the 26S RNA promoter. The unique NotI restriction enzyme site used to linearize prior to *in vitro* transcription is also noted.

5

Figure 5. DNA plasmid map of VEE replicon RNA expressing HIV *env* gene (pERK-DU151env). The plasmid is 13584 base pairs in length and encodes a single polypeptide expressing the four non-structural genes, nsP1-4, the Clade C *env* gene and antibiotic resistance marker, Kanamycin KN(R). The plasmid contains two promoter regions, the T7 polymerase promoter and the 26S RNA promoter. The unique NotI restriction enzyme site used to linearize prior to *in vitro* transcription is also noted.

10

Figure 6. Western immunoblot, demonstrating the expression of HIV proteins in baby hamster kidney (BHK) cells infected with VRPs. The outer lanes of the panel are standard molecular weight markers. Lane 1 is the expression from VRPs encoding the p51 (*pol*) gene. Lane 2 is the expression from VRPs encoding the GP-160 (*env*) gene. Lane 3 is the expression from VRPs encoding the p55 (*gag*) gene. Arrows indicate proteins migrating with the apparent molecular weight of each respective protein.

15

20

Figure 7. Western immunoblot of cells infected with the Du151env VRP. At 18 hr post infection, the cells were lysed and the lysate run in a denaturing polyacrylamide gel. Proteins were transferred out of the gel onto a filter and the filter was probed with serum from subject Du151 using Western immunoblot methods. Lane 1, uninfected U87.CD4-CXCR4 cells. Lane 2, uninfected U87.CD4-CCR5 cells. Lane 3, infection of a mixed culture of U87.CD4-CXCR4 cells and BHK cells (mixtures were used as a positive control in case the U87 cells were refractory to infection by the VRP, which did not turn out to be the case). Lane 4, infected U87.CD4-CXCR4 cells. Lane 5, infected BHK cells. Lane 6, infection of a mixture of BHK cells and U87.CD4-CCR5 cells. Lane 7, infected U87.CD4-CCR5 cells. The positions of

25

30

molecular weight of markers run in the same gel are shown on the right, and the inferred positions of gp160 and gp120 are shown on the left.

Figure 8. Micrographs of U87.CD4-CCR and BHK cells used to examine expression and syncytium formation of DU151 envelope expressed from the VEE replicon. U87.CD4-CCR5 cells alone (panel 1), or a mixture of U87.CD4-CCR5 and BHK cells (Panel 2), BHK cells alone (Panel 3) and U87.CD4-CXCR4 cells (panel 4) were infected with DU151 env VRP at a multiplicity of infection of 3 i.u. per cell. At 18 hours post infection, the cells were examined using light microscopy for the presence of syncytia. The U87.CD4-CCR5 in panel 1 and 2 show clear syncytia, which was absent in the control cell types in the lower panels. In addition, no syncytia were seen in uninfected control cells or VRP-GFP infected cells (data not shown).

Figures 9A-C. Antigen-specific CTL response in mice to the HIV-1 Clade C VRP-gag vaccine. Eight BALB/c mice were immunized twice, first at day 0 and again at day 28 with 10^3 i.u. (Panel A) or 10^5 i.u. (panels B and C) VRP-gag. Eight days (Panels A and B) or 49 days (Panel C) post-boost, spleen cells were isolated and stimulated *in vitro* with vaccinia virus expressing HIV Gag for 1 week. Chromium release assays were performed using vaccinia-Gag infected target cells (diamond symbols) or control vaccinia alone-infected sc11 target cells (square symbols). Clear HIV Gag-specific lysis was detected in animals vaccinated with the VRP-gag vaccine.

Figure 10. Diagrammatic representation of the HIV-1 genome. Black bars indicate relative regions of the genome sequenced to generate phylogenetic sequence comparative data for Clade C *gag*, *pol* and *env* gene isolates.

Figure 11. Phylogenetic comparison of DU422 Clade C Gag isolate with referenced Clade C strains. Consensus clade A, B, D, Mal and SA strains are also shown. DU422 the vaccine strain had 95% amino acid sequence homology to the South African consensus Clade C sequence.

Figure 12. Phylogenetic comparison of DU151 Clade C isolate Env isolate with referenced Clade C strains. DU422 the vaccine strain had 93% amino acid sequence homology to the South African consensus Clade C sequence.

5 Figure 13. Phylogenetic comparison of DU151 Clade C isolate Pol isolate with referenced Clade C strains. DU422 the vaccine strain had 99% amino acid sequence homology to the South African consensus Clade C sequence.

10 Figure 14. DU422 HIV Gag expression as detected by immunofluorescence following electroporation with Gag replicon RNA. BHK cells were electroporated and subjected to immunofluorescence staining with an anti-Gag monoclonal antibody at 24 hours post-electroporation, to demonstrate expression of the Clade C protein.

15 Figure 15. Immunofluorescence detection of DU422 Gag protein expression in BHK cells. BHK cells were infected with VRP-Gag particles and subjected to immunofluorescence staining with an anti-Gag monoclonal antibody at 24 hours post-infection, to demonstrate expression of the Clade C protein.

DETAILED DESCRIPTION OF THE INVENTION

20

As used in the specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" can mean a single pharmaceutical carrier or mixtures of two or more such carriers.

25

The present invention is based on the discovery of a vaccine for the treatment and/or prevention of infection by HIV, comprising novel combinations of isolated nucleic acids encoding two or more distinct antigens which elicit an immune response in a subject which is effective in treating and/or preventing infection by HIV. In a particular embodiment, the nucleic acids encoding the antigens of the vaccine are

30

modified to enhance the immunogenicity of the antigen, improve the safety of the vaccine, or both.

As used herein, the term "isolated nucleic acid" means a nucleic acid separated or substantially free from at least some of the other components of the naturally occurring organism, for example, the cell structural components commonly found associated with nucleic acids in a cellular environment and/or other nucleic acids. The isolation of nucleic acids can be accomplished by well known techniques such as cell lysis or disruption of virus particles, followed by phenol plus chloroform extraction, followed by ethanol precipitation of the nucleic acids (Sambrook *et al.*, latest edition). The nucleic acids of this invention can be isolated according to methods well known in the art for isolating nucleic acids. Alternatively, the nucleic acids of the present invention can be synthesized according to standard protocols well described in the literature for synthesizing nucleic acids.

15

HIV-VRP Vaccines

The antigens of this invention can be gene products which are complete proteins or any fragment of a protein determined to be immunogenic by methods well known in the art. Modifications are made to the antigens of this invention to enhance immunogenicity and/or improve the safety of administration of a vaccine containing the antigen. Examples of such modifications are described in the Examples section herein. Furthermore, it is understood that, where desired, other modifications and changes (e.g., substitutions, deletions, additions) may be made in the amino acid sequence of the antigen of the present invention, which may not specifically impart enhanced immunogenicity or improved safety, yet still result in a protein or fragment which retains all of the functional characteristics by which the protein or fragment is defined. Such changes may occur in natural isolates, may be introduced by synthesis of the protein or fragment, or may be introduced into the amino acid sequence of the protein or fragment using site-specific mutagenesis of nucleic acid encoding the protein or

30

fragment, the procedures for which, such as mis-match polymerase chain reaction (PCR), are well known in the art.

The nucleic acids of this invention can be present in a vector and the vector of
5 this invention can be present in a cell. The vectors and cells of this invention can be in a composition comprising the cell or vector and a pharmaceutically acceptable carrier.

The vector of this invention can be an expression vector which contains all of the genetic components required for expression of the nucleic acids of this invention in
10 cells into which the vector has been introduced, as are well known in the art. For example, the expression vector of this invention can be a vector comprising the helper RNAs of this invention. Such an expression vector can be a commercial expression vector or it can be constructed in the laboratory according to standard molecular biology protocols. The expression vector can comprise viral nucleic acid including, but
15 not limited to, alphavirus, flavivirus, adenovirus, retrovirus and/or adeno-associated virus nucleic acid. The nucleic acid or vector of this invention can also be in a liposome or a delivery vehicle which can be taken up by a cell via receptor-mediated or other type of endocytosis.

20 In another embodiment, the nucleic acids of this invention can be present in a composition comprising a population of alphavirus replicon particles which comprise two or more distinct isolated nucleic acids of this invention and wherein the nucleic acids are each contained within a separate alphavirus replicon particle (herein referred to as a "VRP"). Thus, the expression vector of the present invention can be an
25 alphavirus replicon particle comprising a nucleic acid encoding an antigen of this invention.

In a particular embodiment, the present invention provides a composition comprising two or more isolated nucleic acids selected from the group consisting of an
30 isolated nucleic acid encoding an *env* gene product or an immunogenic fragment

thereof of a human immunodeficiency virus, an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of particles, e.g., virus-like particles, containing the *gag* gene product or the immunogenic fragment thereof, and their release from a cell, and an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity.

10 In a preferred embodiment, the invention provides alphavirus replicon particles (VRPs) that can be administered as an HIV vaccine. These HIV-VRPs are propagation defective, single cycle vectors that contain a self-amplifying RNA (replicon RNA), e.g., from VEE, in which the structural protein genes of the virus are replaced by a HIV-1 Clade C *gag* gene or any other HIV antigen to be expressed. Following introduction
15 into packaging (or helper) cells *in vitro*, the replicon RNA is packaged into VRPs by supplying the viral structural proteins in *trans* (helper RNAs).

The present invention further provides a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group
20 consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of particles, such as virus-like
25 particles, containing the *gag* gene product or the immunogenic fragment thereof, from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon
30 particle.

It is also contemplated that the compositions of this invention comprise alphavirus replicon particles in which either the replicon RNA or at least one structural protein comprises one or more attenuating mutations. Thus, the present invention additionally provides a population of alphavirus replicon particles comprising two or
5 more distinct types of such particles selected from the group consisting of 1) particles expressing a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) particles expressing a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment
10 thereof is modified to inhibit release of particles, such as virus-like particles, containing the *gag* gene product or the immunogenic fragment thereof, from a cell, and 3) particles expressing a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity; and
15 wherein the nucleic acids are each contained within a separate alphavirus replicon particle and further wherein the alphavirus replicon particles comprise a replicon RNA or at least one structural protein which comprises one or more attenuating mutations.

In a preferred embodiment, the population of alphavirus replicon particles
20 comprises particles expressing the nucleic acids encoding *pol*, *env*, and *gag* gene products. In this embodiment, vigorous antigen-specific cellular (e.g., CTL, NK cell and T-helper) and/or humoral (e.g., antibody) responses can be obtained when such particle populations are administered to a subject.

In the compositions described above, the *gag* gene product or immunogenic fragment thereof can be modified by mutation of the second codon, whereby a glycine is changed to an alanine. Alternatively, the *gag* gene product or immunogenic fragment thereof can be modified by any other means known in the art for inhibiting the release of particles containing the *gag* gene product or immunogenic fragment
30 thereof from a cell.

Furthermore, in the compositions of this invention, the *pol* gene product or immunogenic fragment thereof can be modified by mutation of the nucleotide sequence encoding the active site motif, whereby YMDD is changed to YMAA or HMAA (the latter providing a convenient site for cloning, see SEQ ID NO:16). The *pol* gene product or immunogenic fragment thereof can also be modified by any means known in the art for inhibiting reverse transcriptase activity.

The *pol* gene product or immunogenic fragment thereof of this invention may be further modified such that the coding sequences for integrase and RNase H are removed, inactivated and/or modified, e.g., by producing only the p51 region of the *pol* gene product. This modification has been shown in some studies to reduce the possibility of formation of replication competent alphavirus particles during production of alphavirus replicon particles comprising the *pol* gene product or immunogenic fragment thereof. This modification can be of the nucleic acid encoding the *pol* gene product or immunogenic fragment thereof according to methods known in the art. Thus, the particles and compositions of this invention can comprise nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof.

In the compositions of this invention, the *gag*, *env* or *pol* gene products or immunogenic fragments thereof can be from any HIV isolate or consensus sequence derived from HIV primary isolates now known or later identified, the isolation and characterization of which are well known in the art. Also, in the compositions of this invention, the *gag*, *env* or *pol* gene products or immunogenic fragments thereof can be produced from the same HIV isolate or HIV consensus sequence or from any combination of HIV isolates or HIV consensus sequences. In the Examples provided herein, the nucleic acid sequences encoding the *env*, *gag* and *pol* gene products of this

invention were selected based on a consensus sequence generated from primary isolates obtained from recent seroconvertors in Kwazulu/Natal in South Africa. Sequence analysis of these isolates identified them as subtype (or clade) C, and in preferred embodiments of the invention, the *env*, *gag* and *pol* genes are from Clade C isolates of HIV.

In preferred embodiments, each of the three HIV genes are derived from one or more of the South African isolates obtained from recent seroconvertors in Kwazulu/Natal as described herein (see Figures 11-13 for isolate names). In a further embodiment, the *gag* gene or gene fragment is from a *gag* sequence having 95% or greater amino acid identity with the South African consensus sequence for the *gag* gene. In a specific embodiment, the *gag* gene or fragment thereof is derived from HIV Subtype Clade C isolate DU422 and the *env* and *pol* genes or fragments thereof are derived from HIV isolate DU151.

The term "alphavirus" has its conventional meaning in the art and includes the various species of the alphavirus genus, such as Eastern Equine Encephalitis virus (EEE), Venezuelan Equine Encephalitis virus (VEE), Western Equine Encephalitis virus (WEE), Everglades virus, Mucambo virus, Pixuna virus, Sindbis virus, Semliki Forest virus, South African Arbovirus No. 86, Middleburg virus, Chikungunya virus, O=nyong-nyong virus, Ross River virus, Barmah Forest virus, Getah virus, Sagiya virus, Bebaru virus, Mayaro virus, Una virus, Aura virus, Whataroa virus, Babanki virus, Kyzylgach virus, Highlands J virus, Fort Morgan virus, Ndumu virus, Buggy Creek virus, as well as any specific strains of these alphaviruses (e.g., TR339; Girdwood) and any other virus classified by the International Committee on Taxonomy of Viruses (ICTV) as an alphavirus.

An "alphavirus replicon particle" as used herein is an infectious, replication defective, alphavirus particle which comprises alphavirus structural proteins and further comprises a replicon RNA. The replicon RNA comprises nucleic acid encoding the

alphavirus packaging segment, nucleic acid encoding alphavirus non-structural proteins and a heterologous nucleic acid sequence encoding an antigen of this invention. The non-structural proteins encoded by the replicon RNA may be such proteins as are required for replication and transcription. In a specific embodiment of this invention, the structure of the replicon RNA, starting at the 5' end, comprises the 5' untranslated region of the alphavirus RNA, the non-structural proteins (e.g., nsPs1-4) of the alphavirus, the 26S promoter (also known as the "subgenomic promoter"), the heterologous nucleic acid encoding an HIV antigen, and the 3' untranslated region of the alphavirus RNA. An example of a nucleic acid encoding alphavirus nonstructural proteins that can be incorporated into the embodiments of this invention is SEQ ID NO:2, which encodes the amino acid sequence of SEQ ID NO:3.

Although the alphavirus replicon RNA can comprise nucleic acid encoding one or two alphavirus structural proteins, the replicon RNA does not contain nucleic acid encoding all of the alphavirus structural proteins. The replicon RNA can lack nucleic acid encoding any alphavirus structural protein(s). Thus, the resulting alphavirus replicon particles of this invention are replication defective inasmuch as the replicon RNA does not encode all of the structural proteins required for encapsidation of the replicon RNA and assembly of an infectious virion.

As used herein, "alphavirus structural protein" or "structural protein" means the alphavirus proteins required for encapsidation of alphavirus replicon RNA and packaging of the encapsidated RNA into a virus particle. The alphavirus structural proteins include PE2, E2, E3, 6K and E1.

The alphavirus replicon particles of this invention can comprise replicon RNA from any of the alphaviruses of this invention. Furthermore, the alphavirus replicon particles of this invention can comprise alphavirus structural proteins from any of the alphaviruses of this invention. Thus, the replicon particles can be made up of replicon RNA and structural proteins from the same alphavirus or from different alphaviruses,

the latter of which would be chimeric alphavirus replicon particles (e.g., a particle comprising Sindbis virus replicon RNA and VEE structural proteins).

The alphavirus replicon particles of this invention can be made by employing a
5 helper cell for expressing an infectious, replication defective, alphavirus particle in an alphavirus-permissive cell. The helper cell includes (a) a first helper RNA encoding (i) at least one alphavirus structural protein, and (ii) not encoding at least one alphavirus structural protein; and (b) a second helper RNA separate from the first helper RNA, the second helper RNA (i) not encoding the at least one alphavirus structural protein
10 encoded by the first helper RNA, and (ii) encoding at least one alphavirus structural protein not encoded by the first helper RNA, such that all of the alphavirus structural proteins assemble together into alphavirus particles in the cell.

The alphavirus structural protein genes can be present on the helper RNAs of this invention in any combination. For example, the helper RNA of this invention can
15 encode the alphavirus capsid and E1, capsid and E2, E1 and E2, capsid only, E1 only, E2 only, etc. It is also contemplated that the alphavirus structural proteins are provided in *trans* from genes located on three separate RNA molecules within the helper cell.

In a preferred embodiment, the helper cell also includes a replicon RNA, which
20 encodes the alphavirus packaging segment and an inserted heterologous RNA. In the embodiment wherein the helper cell also includes a replicon RNA, the alphavirus packaging segment may be, and preferably is, deleted from both the first helper RNA and the second helper RNA. For example, in an embodiment wherein the helper cell includes a replicon RNA encoding the alphavirus packaging segment and an inserted
25 heterologous RNA, the first helper RNA encodes the alphavirus E1 glycoprotein and the alphavirus E2 glycoprotein, and the second helper RNA encodes the alphavirus capsid protein. In a preferred embodiment, the first helper RNA encodes the E3-E2-6k-E1 cassette from an alphavirus. In an alternative embodiment, the cassette encoded on the first helper RNA is referred to as the E3-E2-E1 cassette. A specific embodiment of
30 this aspect of the invention is diagrammed in Figure 3, and an exemplary nucleotide

sequence is SEQ ID NO:11. The replicon RNA, first helper RNA, and second helper RNA are all on separate molecules and are cotransfected, e.g., by electroporation, into the helper cell, which can be any alphavirus permissive cell, as is well known in the art.

5 In an alternative embodiment, the helper cell includes a replicon RNA encoding the alphavirus packaging segment and an inserted heterologous RNA and also includes the alphavirus capsid protein otherwise encoded by the second helper RNA. The first helper RNA encodes the alphavirus E1 glycoprotein and the alphavirus E2 glycoprotein. Thus, the replicon RNA and the first helper RNA are on separate
10 molecules, and the replicon RNA and the second helper RNA are on a single molecule.

The RNA encoding the structural proteins, i.e., the first helper RNA and the second helper RNA, can include one or more attenuating mutations. In a preferred
15 embodiment, either one or both of the first helper RNA and the second helper RNA include at least one attenuating mutation. The attenuating mutations provide the advantage that in the event of RNA recombination within the cell, the coming together of the structural and non-structural genes will produce a virus of decreased virulence.

20 The alphavirus replicon particles of this invention can be made by a) transfecting a helper cell as given above with a replication defective replicon RNA, b) producing the alphavirus particles in the transfected cell, and c) collecting the alphavirus particles from the cell. The replicon RNA encodes the alphavirus packaging segment and a heterologous RNA. The transfected helper cell further includes the first
25 helper RNA and second helper RNA as described above.

As described hereinabove, the structural proteins used to assemble the alphavirus replicon particles of this invention are distributed among one or more helper RNAs (i.e., a first helper RNA and a second helper RNA). As noted herein, one or
30 more structural protein genes may be located on the replicon RNA, provided that at

least one structural protein gene is deleted from the replicon RNA such that the replicon RNA and resulting alphavirus particle are replication defective. As used herein, the terms "deleted" or "deletion" mean either total deletion of the specified nucleic acid or the deletion of a sufficient portion of the specified nucleic acid to render the nucleic acid and/or its resultant gene product inoperative or nonfunctional, in accordance with standard usage. (See, e.g., U.S. Pat. No. 4,650,764 to Temin *et al.*) The term "replication defective" as used herein means that the replicon RNA cannot replicate in the host cell (i.e., produce infectious viral particles) in the absence of the helper RNA. The replicon RNA is replication defective inasmuch as the replicon RNA does not include all of the alphavirus structural protein genes required for replication, at least one of the required structural protein genes being deleted therefrom.

In one embodiment, the packaging segment or "encapsidation sequence" is deleted from at least the first helper RNA. In a preferred embodiment, the packaging segment is deleted from both the first helper RNA and the second helper RNA. In a specific embodiment, the second helper RNA is constructed from a VEE cDNA clone, deleting all non-structural proteins (i.e., nsPs1-4), the packaging signal, and the glycoprotein cassette (E3-E2-E1). An example of a plasmid encoding such a second helper RNA is provided in Figure 2, and an exemplary nucleotide sequence for such a second helper RNA is SEQ ID NO:8.

In the preferred embodiment wherein the packaging segment is deleted from both the first helper RNA and the second helper RNA, preferably the helper cell contains a replicon RNA in addition to the first helper RNA and the second helper RNA. The replicon RNA encodes the packaging segment and an inserted heterologous RNA encoding an HIV antigen or a fragment thereof. Typically, the inserted heterologous RNA encodes a gene product which is expressed by the target cell, and includes the promoter and regulatory segments necessary for the expression of that gene product in that cell.

In another preferred embodiment, the replicon RNA, the first helper RNA and the second helper RNA are provided on separate molecules such that a first molecule, i.e., the replicon RNA, encodes the packaging segment and the inserted heterologous RNA, a second molecule, i.e., the first helper RNA, encodes at least one but not all of the required alphavirus structural proteins, and a third molecule, i.e., the second helper RNA, encodes at least one but not all of the required alphavirus structural proteins. For example, in one preferred embodiment of the present invention, the helper cell includes a set of RNAs which include (a) a replicon RNA encoding an alphavirus packaging sequence and an inserted heterologous RNA, (b) a first helper RNA encoding the alphavirus E1 glycoprotein and the alphavirus E2 glycoprotein, and (c) a second helper RNA encoding the alphavirus capsid protein, so that the alphavirus E1 glycoprotein, the alphavirus E2 glycoprotein and the capsid protein assemble together into alphavirus particles containing the replicon RNA in the helper cell.

In an alternate embodiment, the replicon RNA and the first helper RNA are on separate molecules, and the replicon RNA and the second helper RNA are on a single molecule together, thereby providing a first molecule, i.e., the first helper RNA, encoding at least one but not all of the required alphavirus structural proteins, and a second molecule, i.e., the replicon RNA and second helper RNA, encoding the packaging segment, the inserted heterologous gene product and the structural protein(s) not encoded by the first helper. Thus, one or more structural protein(s) is encoded by the second helper RNA, but the second helper RNA is located on the second molecule together with the replicon RNA. For example, in one preferred embodiment of the present invention, the helper cell includes a set of RNAs including (a) a replicon RNA encoding an alphavirus packaging sequence, an inserted heterologous RNA, and an alphavirus capsid protein, and (b) a first helper RNA encoding the alphavirus E1 glycoprotein and the alphavirus E2 glycoprotein so that the alphavirus E1 glycoprotein, the alphavirus E2 glycoprotein and the capsid protein assemble together into alphavirus particles in the helper cell.

The present invention also contemplates alphavirus replicon particles which comprise replicon RNA encoding more than one heterologous gene product. For expression of more than one heterologous nucleic acid from a single replicon RNA, a promoter can be inserted upstream of each heterologous nucleic acid on the replicon RNA, such that the promoter regulates expression of the heterologous nucleic acid, resulting in the production of more than one antigen from a single replicon RNA. Another embodiment contemplates the insertion of an IRES sequence, such as the one from the picornavirus, EMC virus, between the heterologous genes downstream from a 26S promoter of the replicon, thus leading to translation of multiple antigens from a single replicon.

In one preferred embodiment of the present invention, the RNA encoding the alphavirus structural proteins, i.e., the capsid, E1 glycoprotein and/or E2 glycoprotein, contains at least one attenuating mutation. It is further contemplated that the RNA encoding the non-structural proteins can contain at least one attenuating mutation. The phrases "attenuating mutation" and "attenuating amino acid," as used herein, mean a nucleotide mutation or an amino acid coded for in view of such a mutation which result in a decreased probability of causing disease in its host (i.e., a loss of virulence), in accordance with standard terminology in the art, See, e.g., Davis *et al.* (1980). The mutation can be, for example, a substitution mutation or an in-frame deletion mutation. The phrase "attenuating mutation" excludes mutations which would be lethal to the virus. Thus, according to this embodiment, the E1 RNA and/or the E2 RNA and/or the capsid RNA can include at least one attenuating mutation. In a more preferred embodiment, the E1 RNA and/or the E2 RNA and/or the capsid RNA includes at least two, or multiple, attenuating mutations. The multiple attenuating mutations may be positioned in either the first helper RNA or in the second helper RNA, or they may be distributed randomly with one or more attenuating mutations being positioned in the first helper RNA and one or more attenuating mutations positioned in the second helper RNA. Appropriate attenuating mutations will be dependent upon the alphavirus used,

as is well known in the art.

For example, when the alphavirus is VEE, suitable attenuating mutations can be in codons at E2 amino acid position 76 which specify an attenuating amino acid,
5 preferably lysine, arginine, or histidine as E2 amino acid 76; codons at E2 amino acid position 120 which specify an attenuating amino acid, preferably lysine as E2 amino acid 120; codons at E2 amino acid position 209 which specify an attenuating amino acid, preferably lysine, arginine, or histidine as E2 amino acid 209; codons at E1 amino acid 272 which specify an attenuating mutation, preferably threonine or serine as E1
10 amino acid 272; codons at E1 amino acid 81 which specify an attenuating mutation, preferably isoleucine or leucine as E1 amino acid 81; and codons at E1 amino acid 253 which specify an attenuating mutation, preferably serine or threonine as E1 amino acid 253; and the combination mutation of the deletion of E3 codons 56-59 together with codons at E1 amino acid 253 which specify an attenuating mutation, as provided
15 herein. Other suitable attenuating mutations within the VEE genome will be known to those skilled in the art.

In an alternate embodiment, wherein the alphavirus is the South African Arbovirus No. 86 (S.A.A.R.86), suitable attenuating mutations can be, for example, in
20 codons at nsP1 amino acid position 538 which specify an attenuating amino acid, preferably isoleucine as nsP1 amino acid 538; codons at E2 amino acid position 304 which specify an attenuating amino acid, preferably threonine as E2 amino acid 304; codons at E2 amino acid position 314 which specify an attenuating amino acid, preferably lysine as E2 amino acid 314; codons at E2 amino acid position 376 which
25 specify an attenuating amino acid, preferably alanine as E2 amino acid 376; codons at E2 amino acid position 372 which specify an attenuating amino acid, preferably leucine as E2 amino acid 372; codons at nsP2 amino acid position 96 which specify an attenuating amino acid, preferably glycine as nsP2 amino acid 96; codons at nsP2 amino acid position 372 which specify an attenuating amino acid, preferably valine as
30 nsP2 amino acid 372; in combination, codons at E2 amino acid residues 304, 314, 372

and 376; codons at E2 amino acid position 378 which specify an attenuating amino acid, preferably leucine as E2 amino acid 378; codons at nsP2 amino acid residue 372 which specify an attenuating mutation, preferably valine as nsP2 amino acid 372; in combination, codons at nsP2 amino acid residues 96 and 372 attenuating substitution
5 mutations at nsP2 amino acid residues 96 and 372; codons at nsP2 amino acid residue 529 which specify an attenuating mutation, preferably leucine, at nsP2 amino acid residue 529; codons at nsP2 amino acid residue 571 which specify an attenuating mutation, preferably asparagine, at nsP2 amino acid residue 571; codons at nsP2 amino acid residue 682 which specify an attenuating mutation, preferably arginine, at nsP2
10 amino acid residue 682; codons at nsP2 amino acid residue 804 which specify an attenuating mutation, preferably arginine, at nsP2 amino acid residue 804; codons at nsP3 amino acid residue 22 which specify an attenuating mutation, preferably arginine, at nsP3 amino acid residue 22; and in combination, codons at nsP2 amino acid residues 529, 571, 682 and 804, and at nsP3 amino acid residue 22, specifying attenuating
15 amino acids at nsP2 amino acid residues 529, 571, 682 and 804 and at nsP3 amino acid residue 22. Other suitable attenuating mutations within the S.A.A.R.86 genome will be known to those skilled in the art.

The alphavirus capsid gene used to make alphavirus replicon particles can also
20 be subjected to site-directed mutagenesis. The altered capsid protein provides additional assurance that recombination to produce the virulent virus will not occur. The altered capsid protein gene which functions in particle assembly but not in autoproteolysis provides helper function for production of replicon particles, but does not allow for production of a viable recombinant. The capsid residues required for
25 proteolytic function are known (Strauss *et al.*, 1990).

Suitable attenuating mutations useful in embodiments wherein any of the alphaviruses of this invention are employed are known to or can be identified by those skilled in the art using routine protocols. Attenuating mutations may be introduced into
30 the RNA by performing site-directed mutagenesis on the cDNA which encodes the

RNA, in accordance with known procedures. See Kunkel (1985), the disclosure of which is incorporated herein by reference in its entirety. Alternatively, mutations may be introduced into the RNA by replacement of homologous restriction fragments in the cDNA which encodes for the RNA, in accordance with known procedures. The
5 identification of a particular mutation in an alphavirus as attenuating is done using routine experimentation according to methods well known in the art.

Preferably, the helper RNA of this invention includes a promoter. It is also preferred that the replicon RNA includes a promoter. Suitable promoters for inclusion
10 in the helper RNA and replicon RNA are well known in the art. One preferred promoter is the alphavirus 26S promoter, although many suitable promoters are available, as is well known in the art.

In the system wherein a first helper RNA, a second helper RNA, and a replicon
15 RNA are all on separate molecules, if the same promoter is used for all three RNAs, then a homologous sequence between the three molecules is provided. Thus, it is advantageous to employ different promoters on the first and second helper RNAs to provide further impediment to RNA recombination that might produce virulent virus. It is preferred that the selected promoter is operative with the non-structural proteins
20 encoded by the replicon RNA molecule.

The infectious, replication defective, alphavirus particles of this invention are prepared according to the methods disclosed herein in combination with techniques known to those skilled in the art. The methods include, for example, transfecting an
25 alphavirus-permissive cell with a replication defective replicon RNA including the alphavirus packaging segment and an inserted heterologous RNA, a first helper RNA encoding at least one alphavirus structural protein, and a second helper RNA encoding at least one alphavirus structural protein which is different from that encoded by the first helper RNA; producing the alphavirus particles in the transfected cell; and
30 collecting the alphavirus particles from the cell.

Methods for transfecting the alphavirus-permissive cell with the replicon RNA and helper RNAs can be achieved, for example, by (i) treating the cells with DEAE-dextran, (ii) by lipofection, by treating the cells with, for example, LIPOFECTIN, and (iii) by electroporation, with electroporation being a preferred means of achieving RNA uptake into the alphavirus-permissive cells. Examples of these techniques are well known in the art, see e.g., U.S. Pat. No. 5,185,440 to Davis *et al.*, and PCT Publication No. WO 92/10578 to Bioption AB, the disclosures of which are incorporated herein by reference in their entirety.

The steps of producing the infectious viral particles in the cells may also be carried out using conventional techniques. See e.g., U.S. Patent No. 5,185,440 to Davis *et al.*, PCT Publication No. WO 92/10578 to Bioption AB, and U.S. Patent No. 4,650,764 to Temin *et al.* (although Temin *et al.*, relates to retroviruses rather than alphaviruses). The infectious viral particles may be produced by standard cell culture growth techniques.

The steps of collecting the infectious alphavirus particles may also be carried out using conventional techniques. For example, the infectious particles may be collected by cell lysis, or collection of the supernatant of the cell culture, as is known in the art. See e.g., U.S. Patent No. 5,185,440 to Davis *et al.*, PCT Publication No. WO 92/10578 to Bioption AB, and U.S. Patent No. 4,650,764 to Temin *et al.* (although Temin *et al.* relates to retroviruses rather than alphaviruses). Other suitable techniques will be known to those skilled in the art. Optionally, the collected infectious alphavirus particles may be purified, if desired. Purification techniques for viruses are well known to those skilled in the art, and these are suitable for the purification of small batches of infectious alphavirus particles.

Thus, the present invention provides a method of making the populations of alphavirus replicon particles of this invention comprising:

A) (a) providing a first helper cell for producing a first population of infectious,

defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein;
 - and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells;
- B) (a) providing a second helper cell for producing a second population of
- infectious, defective alphavirus particles, comprising in an alphavirus-permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles

containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

- 5 (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- 10 (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

- 15 wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- 20 (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;
- C) providing a third helper cell for producing a third population of infectious, defective alphavirus particles, comprising in an alphavirus-permissive cell:

- 25 (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity or is modified to inactivate or delete integrase, RNase H and reverse
- 30 transcriptase functions in the *pol* gene product or immunogenic fragment

thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

(b) producing the alphavirus particles in the helper cell; and

(c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the populations of alphavirus replicon particles.

In a preferred embodiment, as noted above, the method provided also includes a mutation in the *pol* gene product or immunogenic fragment thereof resulting in inactivation or deletion of integrase and RNase H functions of the *pol* gene product or immunogenic fragment thereof. In a specific embodiment of this method, the region of the *pol* gene encoding the RNase H and integrase function of the *pol* gene product or immunogenic fragment thereof has been deleted.

A method of making the populations of alphavirus replicon particles of this invention, wherein the particles comprise at least one attenuating mutation, is also provided, comprising:

- 5 A) (a) providing a first helper cell for producing a first population of infectious, defective alphavirus particles, comprising in an alphavirus-permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences
10 encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - 15 (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging
20 signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on
25 permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells;
- 30 B) providing a second helper cell for producing a second population of infectious,

defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit release of particles, such as virus-like particles, containing the *gag* gene product or the immunogenic fragment thereof from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells;
- C) providing a third helper cell for producing a third population of infectious, defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity or is modified to inactivate or delete integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and
- D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second

helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the populations of alphavirus replicon particles of the present invention comprising at least one attenuating mutation.

5 In a preferred embodiment, as noted above, the method provided above can include a further mutation in the *pol* gene product or immunogenic fragment thereof resulting in inactivation or deletion of integrase and RNase H functions of the *pol* gene product or immunogenic fragment thereof. In a specific embodiment of this method, the region of the *pol* gene encoding the RNase H and integrase function of the *pol* gene
10 product or immunogenic fragment thereof has been deleted.

It is also contemplated regarding the method described above, that not all of the first, second and third populations of alphavirus particles do not all have to comprise an attenuating mutation. For example, the first population may comprise attenuating
15 mutations, but the second and third populations may not, etc.

The present invention further provides the compositions of the present invention which are produced by the methods of this invention.

20 The compositions and methods of this invention which incorporate attenuating mutations into the alphavirus replicon particles forming the composition and/or produced by the methods include purified compositions and methods of purification based on the presence of the attenuating mutations. In particular, certain attenuating mutations in the alphavirus structural proteins introduce heparin binding sites into these
25 proteins which are present on the surface of the alphavirus replicon particles. As an example, the V3014 E2 glycoprotein (SEQ ID NO:12 and SEQ ID NO:13) has a mutation in which a lysine is substituted for the glutamic acid at amino acid position 209. This mutation, which creates a more positively charged glycoprotein, increases the affinity of this protein for heparin. Thus, it is possible to purify such particles using
30 heparin affinity chromatography. Such chromatography can be performed using any of

several commercially available resins to which heparin has been bound. The source of heparin is variable; the commercially available resins currently use porcine heparin. The choice of resin will be based on its relative ease of use in a scaled-up, GMP-compliant process, e.g., price, column packing limitations, and potential for easy
5 sanitization. The use of heparin affinity chromatography results in a substantial purification of the VRPs with very little loss of material, and it is a scalable purification step. In a preferred embodiment, a heparin affinity chromatography step results in between an 8- to 27-fold reduction in total protein per ml, or from a 300- to 1000-fold reduction in total protein per VRP. Thus, the present invention provides heparin
10 affinity-purified alphavirus replicon particles containing attenuating mutations which are useful as clinical trial material and commercial product. The present invention also provides methods for preparing purified alphavirus replicon particles containing attenuating mutations comprising the use of heparin affinity chromatography, as described in the Examples provided herein. These particles can also be present in a
15 composition of this invention.

The alphavirus replicon particles of this invention can also be made in a cell free system. Such replicon particles are herein referred to as virosomes. In a specific embodiment of the method, such particles are constructed from a mixture containing
20 replicon RNA that does not encode all of the alphavirus structural proteins, purified glycoproteins E1 and E2, one or more non-cationic lipids, such as lecithin, and detergent. Detergent is slowly removed from the mixture to allow formation of lipid bilayers with incorporated RNA and glycoproteins.

In preferred embodiments of the methods of this invention, the glycoproteins E1 and E2 could be expressed in any recombinant protein expression system capable of glycosylation of mammalian proteins, such as stably transformed cell lines, for example CHO cells, or viral vector expression systems such as vaccinia, baculovirus, herpes virus, alphavirus or adenovirus. In a preferred embodiment, following expression of
30 the proteins, the E1 and E2 glycoproteins are purified from contaminating cellular

proteins in the expression supernatant. The purification of these glycoproteins can be achieved by affinity chromatographic column purification, for example using lectin-, heparin-, or antibody-affinity columns. This affinity purification step may be preceded by selective precipitation or selective extraction from the expression system supernatant

5 by methods including, but not limited to, ammonium sulfate precipitation or detergent extraction respectively. Final polishing steps of purification may include ion-exchange chromatography or buffer exchange, for example, and tangential flow methods to generate purified glycoproteins suitable for virosome assembly.

10 Thus, the present invention provides a method of producing alphavirus replicon virosomes, comprising: a) combining alphavirus replicon RNA, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and b) gradually removing detergent, whereby alphavirus replicon virosomes are produced. This method is described in more detail in the Examples section herein.

15 The present invention also provides alphavirus replicon virosomes comprising an alphavirus replicon RNA encapsidated by a lipid bilayer in which alphavirus glycoproteins are embedded. The replicon RNA can be from any alphavirus and the glycoproteins can be from any alphavirus. In a specific embodiment, the alphavirus

20 glycoproteins are VEE E1 and E2. The advantage of the alphavirus replicon virosomes is the ease of preparation, their stability, and their purity, since they are devoid of any cellular components being made in a cell free system.

The helper cells, RNAs and methods of the present invention are useful in *in vitro* expression systems, wherein the inserted heterologous RNA located on the

25 replicon RNA encodes a protein or peptide which is desirably produced *in vitro*. The helper cells, RNAs, methods, compositions and pharmaceutical formulations of the present invention are additionally useful in a method of administering a protein or peptide to a subject in need of the desired protein or peptide, as a method of treatment

30 or otherwise.

It is contemplated that the nucleic acids, vectors and alphavirus replicon particles of this invention can be administered to a subject to impart a therapeutic or beneficial effect. Therefore, the nucleic acids, vectors and particles of this invention can be present in a pharmaceutically acceptable carrier. By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material may be administered to a subject, along with the nucleic acid or vector of this invention, without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. The carrier would naturally be selected to minimize any degradation of the active ingredient and to minimize any adverse side effects in the subject, as would be well known to one of skill in the art (see, e.g., *Remington's Pharmaceutical Science*; latest edition).

Pharmaceutical formulations of this invention, such as vaccines, of the present invention can comprise an immunogenic amount of the alphavirus replicon particles as disclosed herein in combination with a pharmaceutically acceptable carrier. An "immunogenic amount" is an amount of the infectious alphavirus particles which is sufficient to evoke an immune response (humoral and/or cellular immune response) in the subject to which the pharmaceutical formulation is administered. An amount of from about 10^3 to about 10^7 replicon-containing particles, and preferably, about 10^4 to about 10^6 replicon-containing particles per dose is believed suitable, depending upon the age and species of the subject being treated. Exemplary pharmaceutically acceptable carriers include, but are not limited to, sterile pyrogen-free water and sterile pyrogen-free physiological saline solution.

Subjects which may be administered immunogenic amounts of the infectious, replication defective alphavirus particles of the present invention include, but are not limited to, human and animal (e.g., horse, donkey, mouse, hamster, monkey) subjects. Administration may be by any suitable means, such as intraperitoneal or intramuscular injection.

Pharmaceutical formulations for the present invention can include those suitable for parenteral (e.g., subcutaneous, intradermal, intramuscular, intravenous and intraarticular) administration. Alternatively, pharmaceutical formulations of the present invention may be suitable for administration to the mucous membranes of a subject (e.g., intranasal administration). The formulations may be conveniently prepared in unit dosage form and may be prepared by any of the methods well known in the art.

Thus, the present invention provides a method for delivering nucleic acids and vectors (e.g., alphavirus replicon particles; virosomes) encoding the antigens of this invention to a cell, comprising administering the nucleic acids or vectors to a cell under conditions whereby the nucleic acids are expressed, thereby delivering the antigens of this invention to the cell. The nucleic acids can be delivered as naked DNA or in a vector (which can be a viral vector) or other delivery vehicles and can be delivered to cells *in vivo* and/or *ex vivo* by a variety of mechanisms well known in the art (e.g., uptake of naked DNA, viral infection, liposome fusion, endocytosis and the like). The cell can be any cell which can take up and express exogenous nucleic acids.

Further provided herein is a method of inducing an immune response to an HIV antigen of this invention in a subject, comprising administering to the subject an immunogenic amount of the particles, virosomes and/or composition of this invention, in a pharmaceutically acceptable carrier.

A method of treating and/or preventing infection by HIV in a subject is also provided herein, comprising administering to the subject an effective amount of the particles, virosomes and/or compositions of this invention, in a pharmaceutically acceptable carrier.

The subject of this invention can be any animal in which an immune response can be induced or in which an infection by HIV can be treated and/or prevented. In a preferred embodiment, the subject of this invention is a mammal and most preferably is

a human.

Protocols and data regarding the testing of the compositions of this invention in animals and protocols for administration to humans are provided in the Examples
5 herein.

In a particular embodiment, the present invention provides an isolated nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the integrase, RNase H and reverse transcriptase
10 functions of the *pol* gene product or immunogenic fragment thereof have been inactivated or deleted. Such a modification has been shown in some studies to facilitate inhibition of the formation of replication competent alphavirus particles during production of alphavirus replicon particles comprising the *pol* gene product or immunogenic fragment thereof.

15

Also provided herein is a composition comprising the *pol*-expressing nucleic acid described above, a vector comprising the nucleic acid and a cell comprising the vector. The *pol*-expressing nucleic acid can also be present in an alphavirus replicon particle comprising the nucleic acid.

20

As noted above, the nucleic acid encoding the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in the inhibition of reverse transcriptase activity. In a preferred embodiment, a mutation is introduced at the active site motif that results in inhibition of reverse transcriptase activity. Such a mutation
25 may remove the DNA binding domain of the enzyme, for example. A mutation from YMDD to YMAA or HMAA at this motif is an example of such a mutation.

The present invention additionally provides a method of making an alphavirus replicon particle comprising nucleic acid encoding a *pol* gene product or immunogenic
30 fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or

immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions from the *pol* gene product or immunogenic fragment thereof, comprising

A) providing a helper cell for producing an infectious, defective alphavirus

5 particle, comprising in an alphavirus-permissive cell:

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic
10 fragment thereof is modified to delete or inactivate RNase H, integrase and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein;
15 and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein
20 not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper
25 RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

(B) producing the alphavirus particles in the helper cell; and

30 (C) collecting the alphavirus particles from the helper cell.

In the method provided above, at least one of the replicon RNA, the first helper RNA, and the one or more additional helper RNA(s) can comprise one or more attenuating mutations, as described herein.

5 In a specific embodiment of this method, a mutation is introduced at the active site motif in the *pol* gene product or immunogenic fragment thereof that results in inhibition of reverse transcriptase activity. Such a mutation may remove the DNA binding domain of the enzyme, for example. A mutation from YMDD to YMAA or HMAA at this motif is an example of such a mutation.

10

Also provided herein is an alphavirus replicon particle expressing the *pol* gene product or immunogenic fragment thereof, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene
15 product or immunogenic fragment thereof, produced according to any of the above methods.

In a further embodiment, the present invention provides a method of inducing an immune response in a subject, comprising administering to the subject an
20 immunogenic amount of a composition comprising an alphavirus particle comprising nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic
25 fragment thereof, in a pharmaceutically acceptable carrier.

Furthermore, the present invention provides a method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an effective amount of a composition comprising an alphavirus particle
30 comprising nucleic acid encoding a *pol* gene product or immunogenic fragment thereof

of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, in a pharmaceutically acceptable carrier.

5

In preferred embodiments of the methods of this invention, the subject is administered an effective amount of a population of alphavirus particles comprising particles expressing (1) nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or
10 immunogenic fragment thereof comprises a modification resulting in inactivation or deletion of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, (2) nucleic acid encoding a *gag* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit
15 release of *gag* gene product or the immunogenic fragment thereof from a cell, and (3) nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus in a pharmaceutically acceptable carrier.

In further preferred embodiments, the population of alphavirus particles
20 comprises particles expressing (1) nucleic acid encoding a *gag* gene sequence that has at least 92% identity with SEQ ID NO:4; (2) nucleic acid encoding a *pol* gene sequence that has at least 99% identity with SEQ ID NO:15; and (3) nucleic acid encoding an *env* gene sequence with at least 95% identity with SEQ ID NO:18. In a specific embodiment, the population of alphavirus particles comprises particles expressing (1)
25 nucleic acid of SEQ ID NO:4, (2) nucleic acid of SEQ ID NO:15, and (3) nucleic acid of SEQ ID NO:18.

EXAMPLES

30 The following examples are provided to illustrate the present invention, and

should not be construed as limiting thereof. In these examples, nm means nanometer, mL means milliliter, pfu/mL means plaque forming units/milliliter, VEE means Venezuelan Equine Encephalitis virus, EMC means encephalomyocarditis virus, BHK means baby hamster kidney cells, HIA means hemagglutinin gene, N means nucleocapsid, FACS means fluorescence activated cell sorter, and IRES means internal ribosome entry site. The expression "E2 amino acid (e.g., lys, thr, etc.) number" indicates the designated amino acid at the designated residue of the E2 gene, and is also used to refer to amino acids at specific residues in the E1 protein and in the E3 protein, respectively.

10

EXAMPLE 1

VEE Replicon Particles as Vaccines

Replicon particles for use as a vaccine are produced using the VEE-based vector system, originally developed from a full-length, infectious cDNA clone of the RNA genome of VEE (Figure 1 in Davis *et al.*, 1989). In this Example, one or more attenuating mutations (Johnston and Smith, 1988; Davis *et al.*, 1990) have been inserted into the clone to generate attenuated VEE vaccine vectors (Davis *et al.*, 1991; 1995; Grieder *et al.*, 1995).

20

As described herein, these constructs are genetically modified to create an RNA replicon (i.e., an RNA that self-amplifies and expresses), and one or more helper RNAs to allow packaging. The replicon RNA expresses an HIV gene, e.g., the Clade C HIV-1 *gag* gene. The replicon RNA is packaged into virus-like particles (herein referred to as "virus replicon particles" or "VRPs") that are infectious for only one cycle. During this cycle, the characteristics of the alphavirus-based vector result in very high levels of expression of the replicon RNA in cells to which the VRP is targeted, e.g., cells of the lymph node.

30

In the cytoplasm of the target cell, the replicon RNA is first translated to

- produce the viral replicase proteins necessary to initiate self-amplification and expression. In this Example, the HIV-1 Clade C *gag* gene is encoded by a subgenomic mRNA, abundantly transcribed from a negative-sense replicon RNA intermediate, leading to high-level expression of the HIV-1 Clade C *gag* gene product. Since the
- 5 VEE structural protein genes are not encoded by the replicon RNA, progeny virion particles are not assembled, thus limiting the replication to a single cycle within the infected target cell.

- Importantly, only the replicon RNA is packaged into VRPs, as the helper RNAs
- 10 lack the *cis*-acting packaging sequence required for encapsidation. The "split helper" or bipartite system (see Example 4) greatly reduces the chance for an intact genome being assembled by recombination, and as a back-up safety feature, one or more highly attenuating mutations, such as those contained in the glycoprotein genes in V3014 (Grieder *et al.*, 1995), are incorporated.

15

- Overall, the design of the VRPs incorporates several layered and redundant safety features. In addition to the above-described split helper system and attenuating mutations, over one-third of the genome of the virus has been removed, creating a defective genome which prevents spread from the initially infected target cell.
- 20 Nonetheless, if a statistically rare recombination event occurs to yield replication competent virus (RCV), the resulting virus would be a highly attenuated VEE strain.

EXAMPLE 2

Construction of VEE Replicon

25

- The VEE structural protein genes (C-PE2-6K-E1) are removed from a cDNA clone (pV4031) which contained two attenuating mutations (E2 lys 209, E1 thr 272), and a duplication of the 26S subgenomic RNA promoter sequence immediately downstream from the 3'-end of the E1 glycoprotein gene, followed by a multiple
- 30 cloning site as described in U.S. Pat. No. 5,505,947 to Johnston *et al.* The pV4031

plasmid DNA is digested to completion with *Apal* restriction enzyme, which cuts the VEE genomic sequence at nucleotide 7505 (numbered from the 5'-end of the genome sequence). A second recognition site for this enzyme is found in the duplicate 26S subgenomic promoter. Therefore, digestion of pV4031 with *Apal* produces two DNA fragments, one containing the VEE nonstructural genes (e.g. SEQ ID NO:2) and a single copy of the 26S subgenomic RNA promoter followed by a multiple cloning site, and a second smaller fragment containing a 26S subgenomic RNA promoter followed by the VEE structural genes. The large fragment is isolated and religated to produce the replicon, pVR2. In this example, as well as in the construction of the helper plasmids (Example 3), a kanamycin resistance gene (SEQ ID NO:6, encoding amino acid sequence as in SEQ ID NO:7) is present in the plasmids to aid in the cloning manipulations.

EXAMPLE 3

Construction of Helper Plasmids

The starting materials for the helper plasmids are four full-length cDNA clones: V3000, the virulent Trinidad donkey strain of VEE, three clones with attenuating mutations, pV3014 (E2 lys 209, E1 thr 272), V3519 (E2 lys 76, E2 lys 209, E1 thr 272) and V3526 (deletion of E3 56-59, E1 ser 253), which are in the genetic background of Trinidad donkey strain VEE. Several different helper plasmids have been made by using unique or rare restriction sites in the full-length cDNA clone to delete portions of the nonstructural protein region. The full-length clone is digested with one or two restriction enzymes, the larger DNA fragment is isolated and then religated to form a functional plasmid. *In vitro* RNA transcripts from these plasmids upon transfection of tissue culture cells would not encode a functional RNA replication complex, and also would not include an encapsidation signal. The helper constructs differ in the size of the nonstructural gene deletion. The helper constructs are designated by the attenuated mutant clone used in their construction, and by the percentage of the nonstructural region deleted. The following helper constructs were generated:

	V3014Δ520-7507(93%)
	V3519Δ520-7507(93%)
	V3526Δ520-7505(93%)
	V3014Δ520-6965(87%)
5	V3519Δ1687-7507(78%)
	V3014Δ2311-7505(70%)
	V3519Δ3958-7507(47%)
	V3526Δ520-7505(93%)
	V3014Δ3958-7505(47%)
10	V3519Δ1955-3359(19%)
	V3014Δ520-3954(46%)
	V3014Δ1955-3359(19%)
	V3014Δ1951-3359(19%)
	V3014Δ2311-3055(10%)
15	V3014Δ2307-3055(10%)

EXAMPLE 4

Construction of Bipartite RNA Helper Plasmids

- 20 A bipartite helper system is constructed as described herein. The V3014Δ520-7505(93%) helper is used to construct an additional deletion of the E2 and E1 glycoprotein genes by digestion with HpaI restriction enzyme and ligation, resulting in deletion of the sequence between nucleotide 8494 (in the E3 gene) and nucleotide 11,299 (near the 3'-end of the E1 gene). *In vitro* RNA transcripts of this glycoprotein
- 25 helper plasmid (presented graphically in Figure 2; an exemplary nucleotide sequence for such a plasmid is SEQ ID NO:8, including the nucleotide sequence (SEQ ID NO:9 and the amino acid sequence (SEQ ID NO:10 of the VEE capsid), when electroporated into BHK cells with a replicon RNA, are replicated and transcribed to give a mRNA encoding only the capsid protein of VEE.

The second member of the bipartite helper is constructed from the same original helper plasmid 3014Δ5207505(93%) by cleavage with Tth111I restriction enzyme (at nucleotide 7544) and SpeI restriction enzyme (at nucleotide 8389), resulting in deletion of the capsid gene, followed by insertion of a synthetic double-stranded oligonucleotide with Tth111I and SpeI termini. The inserted sequence restored the downstream portion of the 26S promoter and an ATG initiation codon followed by a Ser codon, such that the first amino acid residue of E3 (Ser) is the first codon following the inserted AUG. The resulting glycoprotein helper plasmid is presented graphically in Figure 3, and an exemplary nucleic acid sequence for such a plasmid is SEQ ID NO:11, encoding the VEE glycoproteins (E3-E2-6kD-E1), SEQ ID NO:12. The *in vitro* transcript of this plasmid, when transfected into a cell with replicon RNA, will produce the VEE glycoproteins (SEQ ID NO:13). Co-electroporation of both of these helper RNAs into a cell with replicon RNA results in production of infectious particles containing only replicon RNA.

15

Other than the 5' and 3' ends and the 26S promoters (40 nucleotides) of these helper RNAs, the only sequence in common between the capsid and glycoprotein helpers is the sequence from 8389 to 8494 (106 nucleotides)

20

EXAMPLE 5

VEE REPLICON PARTICLES EXPRESSING HIV GENES

The vaccines of this invention are exemplified by the use of a propagation defective, replicon particle vector system derived from an attenuated strain of Venezuelan equine encephalitis virus (VEE) to create a mixture of VEE replicon particles individually expressing HIV-1 *gag*, *pol*, or *env* genes. The three genes used in this Example were selected based on homology to consensus sequences generated from primary isolates obtained from recent seroconverters in Kwazulu/Natal. Plasma samples from approximately 20 recent seroconverters in the Durban/Hlabisa cohort and a similar number of HIV-positive, asymptomatic individuals were collected. HIV viral

30

RNA was isolated from the plasma, and the sequences of the *gag*, *pol* and *env* genes were analyzed. Two regions from each gene were amplified, and the resulting PCR products were sequenced (see Figure 10 for regions analyzed). A consensus sequence was derived for each gene, and the sequences of each isolate were compared to the
5 derived consensus. All isolates were found to be Subtype C of HIV, thus confirming the predominance of this subtype in South Africa.

A. CONSTRUCTION OF THE Gag-VRP VACCINE

Described herein is the design and manufacture of VEE replicon particles (VRPs) engineered to express the *gag* gene from a Subtype C isolate of HIV-1. The main purpose of this single antigen vaccine is to establish a safety profile for VRPs in healthy human subjects. Optimally, the HIV-Gag-VRPs will be formulated as a component of a trivalent vaccine, also containing HIV-Pol-VRP and HIV-gp160-VRP
15 (*env*) made in analogous procedures to the one described herein for HIV-Gag-VRPs.

In this Example, the VEE particles are based on the V3014 glycoprotein helper plasmid (Figure 3, SEQ ID NO:12 and SEQ ID NO:13), which harbors two highly attenuating mutations, one in E2 and the other in E1 (Grieder *et al.*, 1995). The V3014
20 glycoprotein helper RNA is able to package VRPs with significantly greater efficiency than the glycoprotein helper RNA derived from V3526 (Pushko *et al.*, 1997). Nonetheless, safety of the VRP vector system has not been compromised since detailed pathogenesis studies clearly have shown V3014 to be avirulent in adult mice by subcutaneous inoculation (Grieder *et al.*, 1995). V3014 was found to be significantly
25 impaired in its ability to reach and spread beyond the draining lymph node following subcutaneous inoculation. Unlike wild-type V3000, V3014 does not establish a viremia and does not reach the brain. In addition, on rare occasions when found, histopathological lesions in the periphery were much less severe than those induced by wild-type V3000 (Grieder *et al.*, 1995). Following inoculation with V3014, adult mice
30 are protected against lethal wild-type VEE infection.

The attenuated phenotype of V3014 also was observed in VEE challenge studies in horses. Animals inoculated subcutaneously with V3014 showed no significant leukopenia or febrile response compared to mock-vaccinated controls. In addition, results indicated that these animals were completely protected against virulent VEE (V3000) challenge.

Taken together, these data indicate that if the rare recombination event did occur during VRP assembly to yield RCV, the worst case scenario would be the generation of a highly attenuated strain of VEE.

10

B. SELECTION AND CLONING OF THE HETEROLOGOUS ANTIGEN

The exemplary HIV genes used in this invention, *gag*, *pol* and *env*, are derived from Subtype C (Clade C) viruses isolated from likely Phase III clinical trial sites in South Africa. The HIV infection rate in South Africa and its long established virology and public health infrastructure make this country an attractive choice for clinical testing of HIV vaccines. Focused sequencing and phylogenetic analysis of the *gag*, *pol*, and *env* genes of these isolates has allowed the selection of genes representative of the Clade C isolates circulating in this region of Africa.

20

1. HIV-1 Clade C *gag* gene

Two 400 bp regions of the *gag* gene were sequenced from approximately 30 plasma samples collected from HIV seropositive individuals in South Africa. A South African consensus sequence was then determined for the *gag* gene as well as a consensus sequence from the Los Alamos database for Subtype C virus. In addition, approximately 20 comparable sequences from Malawi were used, generated as part of another study, to confirm conclusions about sequence variation. Several isolates that were close to the South African consensus sequence were compared to other isolates in distance measurements. Among these 30 isolates, one was chosen as the source for the *gag* gene (SEQ ID NO:4; corresponding to the amino acid sequence in SEQ ID NO:5)

30

for the following reasons.

This isolate had greater than 95% amino acid identity to the South African consensus sequence, representing the approximate middle of the sequence diversity of all isolates. This isolate, known as DU422, came from a recent seroconverter, reflecting currently circulating strains and the transmitted phenotype. The phenotype of DU422 is NS1, CCR5(+), and CXCR4(-).

Prior to the insertion of the *gag* gene into the VEE replicon plasmid vector, the amino terminal myristylation ("myr") site of *gag* was removed to prevent the formation of Gag-containing virus-like particles. Restriction enzyme digests of the *gag* gene plasmid, the capsid helper plasmid, and the glycoprotein helper plasmid were performed to confirm the identity of the three vectors when compared to published maps of the parental plasmid pBR322, with the kanamycin resistance gene substituted for the ampicillin resistance gene. The confirmed plasmid maps of the VEE replicon plasmid containing the DU422 *gag* gene (p3-40.1.6), the capsid helper plasmid (p3-13.2.2), and the glycoprotein helper plasmid (p3-13.4.6) are presented in Figures 1, 2, and 3, respectively. The full nucleotide sequence of each of these plasmids is presented herein as SEQ ID NO:1, SEQ ID NO:8, and SEQ ID NO:11, respectively.

In Figures 6 and 15, expression of this HIV-1 Gag protein in BHK cells infected with VRPs expressing such a *gag* construct is demonstrated (Figure 6: Western blot, lane 3; Figure 15, immunofluorescence detection). The cells were infected at a multiplicity of infection (m.o.i.) of 3.5 infectious units (i.u.) per cell, and expression was measured 18 hours post-infection (p.i.). Cell lysates (from approximately 2×10^3 cells) were collected and fractionated either by a 4-12% gradient SDS-PAGE or by 10% SDS-PAGE. The fractionated polypeptides were transferred to PVDF membranes and probed with human HIV-1 positive serum.

2. HIV-1 Clade C *env* gene

A Clade C *env* gene (aka "gp160") from another HIV isolate, DU151, from a recent seroconverter was chosen based on its 92% amino acid identity to the South African consensus sequence for this gene, determined in an analogous method to the one described for the *gag* gene in Example 5.A.1. The phenotype of the DU151 isolate is NS1, CCR5(+), CXCR4(-). This gene was engineered into a VEE RNA replicon plasmid as shown in Figure 5, and the entire sequence of the plasmid is given at SEQ ID NO:17. The *env* gene construct used in this Example is SEQ ID NO:18.

In Figure 6, expression of this ENV protein (SEQ. ID. NO:19) in BHK cells infected with VRPs expressing this HIV *env* construct is demonstrated (Western blot, lane 2), showing that the protein expressed in the cells is of the correct size and is immunoreactive. In Figure 7, expression of this ENV protein in U87.CD4.CCR5 cells is shown. These cells process the ENV protein into two components, GP120 and GP41. In these cells, the expressed GP160 is fusogenic (see Figure 8).

3. HIV-1 Clade C *pol* gene

A Clade C *pol* gene from isolate DU151 was chosen based on its 99% amino acid identity with the South African consensus sequence. This gene was modified at the active site of the reverse transcriptase encoding sequence to inhibit its activity, and the p51 fragment of this modified gene (SEQ ID NO:15) was engineered into a VEE RNA replicon plasmid. The map of this *pol* plasmid is shown in Figure 4, and the nucleotide sequence of the plasmid is provided as SEQ ID NO:14. In Figure 6, expression of this POL p51 fragment (SEQ ID NO:16) in BHK cells is demonstrated (Western blot, lane 1), showing that the protein expressed in these cells is both the correct size and immunoreactive.

C. IMMUNOLOGICAL RESPONSE TO VRP-GAG VACCINE

Mice were injected subcutaneously in two doses, with 8-9 mice in each group. The mice were immunized once, then immunized a second time, with the same dose, 28 days later. Serum was collected the day prior to the first immunization, then at day 27 ("after 1st immunization") and at day 35 (after 2nd immunization).

The vigorous, antigen-specific humoral response of mice to the HIV-1 Clade C VRP-gag vaccine described in Example 5.A.1. is presented in Table 1. Details of this assay are described in Example 7A.1.

TABLE 1. Humoral Response to VRP-gag Vaccine

Dose:		Total Ab Titer (log ₁₀)
10 ³ i.u. dose:		
	after 1 st immunization	1.3 +/- 0.1
	after 2 nd immunization	2.8 +/- 1.1
10 ⁵ i.u. dose		
	after 1 st immunization	2.1 +/- 0.5
	after 2 nd immunization	4.1 +/- 0.6

The vigorous, antigen-specific CTL response in mice to the HIV-1 Clade C VRP-gag vaccine (Example 5.A.1) is presented in Figure 9. Details of this assay are described in Example 7A.3.

EXAMPLE 6

MANUFACTURING PROCESS FOR HIV VRP VACCINES

A. Manufacturing Process

5

Disclosed herein is a manufacturing process for VRP vaccines that is suitable for large-scale preparation of GMP-compliant (GMP = Good Manufacturing Practices) material for use in human clinical trials or for commercial manufacture. The process includes several steps and after each step (as appropriate), a set of "in process control" (IPC) assays or Release Tests (RT) is performed to confirm the successful completion of the step. The process steps and the accompanying IPC assay(s) or RTs (described in more detail in Example 6D.1 and 6D.2) are as follows:

15	Process Step Linearize 3 DNA plasmids <i>In vitro</i> RNA transcription Electroporation of certified Vero cell line Harvest culture fluids	IPC/RT IPC: Check for linearity IPC: Size, integrity and concentration IPC: Titration/Identity Test for replication-competent virus (RCV) RT:
	Pool the culture fluid	Mycoplasma Adventitious virus
20	Purification of bulk VRP by heparin affinity chromatography	PERT assay IPCs: Heparin residual assay BSA assay Bovine IgG assay

Filtration of bulk VRP

RT:

Test for RCV

Titration/Identity

Contaminating protein/DNA

Sterility

Endotoxin

Formulate, Fill, Release

RT:

Titration/Identity

Sterility

General Safety

B. Preparation of plasmid DNAs

- 5 Stock solutions of replicon plasmid DNA, capsid helper plasmid DNA and glycoprotein helper plasmid DNA are produced in *Escherichia coli* XL2 Blue cells (Stratagene, cat# 200150). All plasmids harbor the kanamycin resistance gene marker. The three plasmid DNAs were manufactured and purified by PureSyn, Inc. (Malvern, PA) under appropriate GLP/GMP procedures, with a complete Batch Record with full
- 10 traceability. Following fermentation and cell harvest, cell paste was lysed with base and plasmid DNAs were purified by ion pair chromatography on PolyFlo™ separation media.

- Prior to release by appropriate quality assurance/quality control oversight, each
- 15 lot of each plasmid DNA is analyzed to confirm identity, purity and quality (Table 2). An approved certificate of analysis for each DNA is then established for each plasmid DNA lot.

Table 2. Plasmid DNA Release Tests

Test	Method	Specification
DNA homogeneity	Agarose gel electrophoresis	>90% supercoiled
<i>E. coli</i> genomic DNA	Southern Blot	<.50 µg/mg plasmid
<i>E. coli</i> RNA	Agarose gel electrophoresis	No detectable bands
Endotoxin	Limulus Amoebocyte Lysate (LAL)	< 0.1 EU/mg
Total protein	Abs 260/280	1.8-1.9
Sterility	Bioburden assay, USP23	< 1 CFU
Identity	Restriction enzyme analysis	Matches map

To produce HIV-VRP vaccine for clinical use, both replicon and helper plasmids are linearized by digestion at the unique Not I site and used as templates for synthesis of run-off transcripts. The quality of the transcription products (i.e., the replicon and the two helper RNAs) is evaluated by agarose gel electrophoresis.

C. Characterization of the Vero cells

Vero cells are used in the production of HIV-VRPs (WHO Vero MCB P139, BioReliance Inc., Rockville, MD). Vials contained approximately 1×10^7 cells/mL in a cryoprotectant solution of 90% fetal bovine serum and 10% dimethyl sulfoxide. A Cell Certification Summary is provided with each lot. BioReliance Inc. has filed a Master File with the FDA regarding the WHO Vero MCB P139.

Vials of WHO Vero MCB P139 cells are expanded into flasks. Each of the flasks is then expanded again in order to prepare the Master Cell Bank (MCB). The Working Cell Bank (WCB) is prepared from the MCB. The MCB is tested for purity and identity. The WCB is tested for adventitious agents (detection of mycoplasma and viruses). Viability tests are performed on both the MCB and the WCB.

Tumorigenicity tests are performed once at the end of the production period.

D. Electroporation

10

Vero cells are cotransfected by electroporation with RNA mixtures comprising replicon RNA transcripts encoding HIV-gag, VEE capsid helper RNA transcripts, and VEE glycoprotein helper RNA transcripts. The transfected cells are transferred to tissue culture vessels and incubated in well-defined culture medium. Following harvest, the HIV-Gag-VRP is purified from pooled culture fluid supernatants by affinity column chromatography. Prior to formulation and filling, purified, bulk HIV-Gag-VRP is tested for the presence of RCV.

E. Final formulated product

20

The HIV-Gag-VRP vaccine is vialled at four different doses. The material is filtered (0.22 μ m) and added to vials at the appropriate concentration and volume, stoppered, quick-frozen and stored at -20° C.

F. Control tests of the Gag-VRP vaccine

1. In-Process Controls

Table 3 below summarizes the In-Process Controls performed during the manufacturing process of the HIV-Gag-VRP Vaccine.

30

Table 3. IPCs during the manufacture of HIV-Gag-VRP Vaccine

5	Test	Method	Target
	Check for linearity	Agarose Gel electrophoresis	Report
10	Size, integrity and concentration of RNAs	Agarose Gel electrophoresis	Report
	Titration/Identity	Indirect immunofluorescence assay (IFA), using standardized Gag-specific antibody preparation	Report
	Test for RCV	CPE Assay	Report
15	Heparin Residual	Chromogenic Inhibitory Assay	Report
	BSA residual	ELISA	Report
20	Bovine IgG Residual	ELISA	Report

2. Release tests

Tables 4 and 5 below summarize the release tests performed on the HIV-Gag-VRP Vaccine.

Table 4. Pool of the Culture Fluids

Test	Method	Target
Adventitious Virus (<i>in vivo</i>)	European guidelines	Negative
Adventitious Virus (<i>in vitro</i>)	5 cell lines	No growth
Mycoplasma	21CFR 610.30	No Growth
Reverse Transcriptase	PERT Assay	Negative

Table 5. Bulk VRP and Final Vial testing

Test	Method	Target Result
Replication competent virus (RCV)	Cytopathic effect (CPE) assay	Absence (in BHK cells, sensitivity is 1-10 pfu V3014)
VRP identity/ potency	Indirect immunofluorescence assay (IFA)	10 ⁶ to 10 ⁶ i.u. per mL
Cellular Protein Contaminant	Bio-Rad® DC protein assay	Total protein content per dose
Cellular DNA Contaminant	Southern Blot or PCR	< 10 ng per dose
Sterility	21 CFR § 610.12	Pass

	Endotoxin	LAL	< 5 EU/dose
5	General Safety	21 CFR § 610.11	Pass
	Particulates	USP	Pass
	Stability	IFA	10^6 to 10^8 i.u. per mL

10

EXAMPLE 7**PRECLINICAL STUDIES**

Pilot lots are manufactured following written procedures (SOPs and STMs) and according to the manufacturing scheme described in Example 6. These pilot lots are prepared and used for two major tasks. The first one is a preclinical immunogenicity evaluation, which includes studies to assess the immune response and the cell-mediated immune response in vaccinated animals. The second major task is a preclinical safety evaluation, which includes evaluations of system toxicity, hematopoietic and immune system toxicity, and local reactogenicity.

20

Finally, an *in situ* hybridization study is performed in mice in order to verify the *in vivo* expression of HIV-Gag-VRP gene product in lymphoid tissue.

A. Immunogenicity Studies

25

A.1 Humoral Immune Response in Mice

Three groups of five female BALB/c mice (4-6 weeks of age) are inoculated subcutaneously with 10^5 , 10^6 , or 10^7 i.u. of the HIV-Gag-VRP at three time points: on day 0, and at weeks 4 and 8. The fourth group, Control Group, receives the vehicle only. Immediately prior to inoculation, and at weeks 3, 5, 8 and 10 post-inoculation,

30

- blood samples are collected for humoral immune response evaluations. Gag protein-specific serum antibody titers and seroconversion rates are measured by ELISA (Caley *et al.*, 1997) against purified, recombinant Gag protein. The source of the antigen is the homologous Clade C *gag* gene expressed in insect or mammalian cells. Antigen specificity also is confirmed by immunoblot analysis. Anti-VEE responses are monitored by ELISA (Johnston and Smith, 1988).

A.2 Humoral Immune Response in Rabbits

- Three groups of five female New Zealand white rabbits are inoculated subcutaneously with 10^5 , 10^6 , or 10^7 i.u. of the HIV-Gag-VRP. The fourth group, Control Group, receives the vehicle only. Immediately prior to inoculation, and at weeks 3, 5, 8 and 10 post-inoculation, blood samples are collected for humoral immune response evaluations.

15

Humoral immune responses are evaluated as described in Section A.1.

A.3 Cell-Mediated Immune Response in Mice

- Three groups of five female BALB/c mice are inoculated subcutaneously with 10^5 , 10^6 , or 10^7 i.u. of the HIV-Gag-VRP at day 0 and day 28. The fourth group, Control Group, receives the vehicle only. Blood samples are collected at week 3 post-inoculation. Spleens are harvested for splenocyte collection on day 7 following the second inoculation for evaluation of cell-mediated immune responses.

25

- The cell-mediated immune response is evaluated by determining the ability of splenic T cells from immunized mice to proliferate *ex vivo* in the presence of either Gag protein or Gag peptide(s). The ability of splenic T and CD4+ T cells to produce interferon- γ and interleukin-4 respectively, is determined. Finally, the ability of cytotoxic T lymphocytes to lyse target cells that present murine major

30

histocompatibility complex class-I restricted epitopes for HIV-1 Clade C Gag protein is measured (see Betts *et al.*, 1997 for methods)

B. Safety Study

5

Three groups of six male and six female New Zealand white rabbits are inoculated subcutaneously with 10^4 , 10^6 , or 3×10^7 i.u. of the HIV-Gag-VRP. The fourth group, Control Group, receives the vehicle only. Animals receive four injections at week 0, week 3, week 6 and Week 9. Half of the animals are sacrificed two days after the last injection (week 9) and the other half at three weeks after the last injection (week 12). Similar studies are performed in mice with a high dose at 10^8 i.u. This level is 100 times the likely primate dose, based on efficacy studies in rhesus macaques.

In addition to system toxicity (record of mortality/morbidity, body temperature, body weight, food consumption and ophthalmic examinations), hematopoietic toxicity is evaluated by quantitating cellular components of peripheral blood, and immune system toxicity is assessed by histopathologic evaluation of the lymphoid organs. Local reactogenicity is evaluated by examining the injection sites grossly and microscopically to determine irritation potential. Serum samples are also tested for the presence of replication competent virus by blind passage in cell culture.

C. In Situ Hybridization Study in Mice

Three groups of five female BALB/c mice are inoculated subcutaneously with 10^5 , 10^6 , or 10^7 i.u. of the HIV-Gag-VRP. The fourth group, Control Group, receives the vehicle only. A single injection is performed in each group.

To verify expression of HIV-GAG-VRP in lymphoid tissue, the draining lymph nodes, spleen, and thymus of the mice are examined by *in situ* hybridization at 24 hours and 48 hours after the single inoculation.

EXAMPLE 8

Heparin Affinity Chromatography of VRPs

- 5 Generally, the majority of contaminating protein is non-VEE protein from the conditioned media. Heparin column capacity requirements for GMP manufacturing runs are therefore based on the volume of conditioned media, rather than the concentration of VRPs. Column parameters are optimized at room temperature, but variations in temperature do not greatly affect performance. The expected yields of
- 10 VRPs can range from 50% to > 90%.

While only minimal leaching of heparin from the columns has been detected, GMP requirements stipulate that a residual heparin assay be performed as an IPC test following the chromatography step.

15

A. Pharmacia HiTrap® Heparin

- Five mL columns of Pharmacia HiTrap® Heparin (cat no. 17-0407-01, Amersham Pharmacia Biotech), pre-equilibrated with 25 mM HEPES/0.25 M NaCl, pH
- 20 7.5, were loaded with HIV-Gag-VRPs produced in Vero cells. After column washing with the equilibration buffer, VRPs were eluted with a 15 column volume gradient from 0.25 – 1.0 M NaCl gradient in 25 mM HEPES, pH 7.5. The HIV-Gag-VRPs eluted at a conductivity of approximately 48 mS/cm. The wash step was optimized (based on the A_{280} peak) at a NaCl concentration between 0.25 M and 0.3 M.

25

B. Heparin Sepharose 6 Fast Flow® resin

- Heparin Sepharose 6 Fast Flow® resin (catalog no. 90-1000-2; Amersham Pharmacia Biotech) is supplied as a bulk resin which allows various size columns to be
- 30 packed as needed. Fast Flow® resins have the advantages of excellent flow

- characteristics and ability to be sanitized with sodium hydroxide solutions, which are particularly useful in a GMP manufacturing process. A 6 mL column was prepared by packing the Heparin Sepharose 6 Fast Flow® bulk resin in a BioRad® Econo-Column chromatography column, which was then pre-equilibrated with 25 mM HEPES/0.12 M NaCl, pH 7.5. VRPs were loaded onto the column, which was then washed with the equilibration buffer. Initial experiments indicated that the VRPs eluted at a lower conductivity (36 mS/cm) with this resin as compared to the HiTrap® Heparin, so the wash conditions were modified accordingly. The VRPs were eluted from the Fast Flow® resin with a 15 column volume gradient from 0.12 M to 1 M NaCl in 25 mM HEPES, pH 7.5.

EXAMPLE 9

Virosome Formation

- The feasibility of virosome formation is demonstrated in a series of experiments in which replicon RNA and RNA encoding the glycoprotein E1 and E2 genes (glycoprotein helper) were first transfected into BHK cells by electroporation. After 18-24 hours, cell supernatants were harvested and tested for the presence of virosomes as described briefly below.

Cell Culture

- BHK cells were used as a cell substrate and were maintained in growth medium (alpha-MEM (Life Technologies), supplemented with 10% Fetal Bovine Serum (Hyclone), 1x Glutamine (Life-Technologies)), in an atmosphere of 5% CO₂ at 37°C. Prior to electroporation, cells were detached from the cell culture vessel using 0.05% trypsin-0.53 mM EDTA solution (Life Technologies). Trypsin was neutralized with growth medium, and cells were washed twice with cold Phosphate-Buffered Saline (PBS, BioWhittaker) and resuspended at a concentration of 1.5×10^7 cells/ml.

RNA Transcription, Electroporation and Virosome Harvest

Plasmid DNA pVR-GFP (green fluorescent protein) was linearized using restriction endonuclease NotI (New England Biolabs) as recommended by the manufacturer. DNA was extracted with phenol:chloroform:iso-amyl alcohol (25:24:1, Gibco BRL) and precipitated with ethanol, following the addition of NH_4Ac to 2.5 M final concentration. RNA was synthesized in an *in vitro* transcription reaction using an mMessage mMachine® kit (Ambion) as recommended by the manufacturer. This RNA, without further purification, was used to transfect BHK cells. Helper RNA was prepared in a similar fashion. A BHK cell suspension in PBS (0.8 mL, 1.2×10^7 cells) was mixed with 10 μg of each RNA, and the mixture was electroporated. Electroporation settings for Gene-Pulser® (Bio-Rad Laboratories) were: 850 V, 25 μF , 3 pulses. Culture supernatant was collected at 18-24 hr post-electroporation and clarified by centrifugation for 10 min at 1000 rpm.

Titration of Virosomes

The presence of infectious virosome particles was demonstrated using an immunofluorescence assay to titer the virosomes by detecting the fluorescence of the GFP encoded by the replicon RNA in the virosomes. Serial dilutions of the cell culture supernatant were added to 12-well plates of BHK cells. Following an 18-24 hour incubation in an atmosphere of 5% CO_2 at 37°C, the medium was removed from each plate. Virosome infectious titer was then determined by counting the number of green-fluorescent single cells at a particular dilution, followed by a back-calculation to determine total infectious units (i.u.) per mL. A final titer of 440 i.u./mL was collected.

Confirmation of virosome identity

Three independent experimental methods were used to determine that the infectious particles were in fact virosomes, rather than replication competent viral

particles or naked RNA being carried over from the electroporated cells.

- i) The virosome-containing supernatant was passaged a second time by removing the cell supernatant from the 12-well plate used for titration and placing this supernatant onto a fresh monolayer of BHK cells. At 18-24 hours post-passage, the monolayer was examined under U/V fluorescence and found to contain 0 (zero) GFP-positive cells, indicating the infectious particles produced using this method can undergo only a single round of replication, a critical characteristic of a virosome.
- ii) To establish that the infectious titer detected following virosome packaging was not due to carry-over of RNA used in the electroporation, the supernatant was treated with RNase A (Invitrogen) at a concentration of 100 $\mu\text{g}/\mu\text{L}$ for 15 minutes at 37°C. The treated and untreated control supernatants were titrated according to the methods outlined above. The RNase-treated sample contained 400 i.u./mL and the control group had 440 i.u./mL, indicating that the RNase treatment had no significant effect on virosome titer.
- iii) To establish that the infectious particles were enveloped in the E1 and E2 glycoproteins, anti-VEE mouse serum was used to treat the cell supernatant in a neutralization assay. As a control, normal mouse serum was used to treat the virosome supernatant. In addition, VEE replicon particles expressing GFP were used in the assay, the infectivity of which is known to be inhibited by this serum.

25	Anti-VEE serum	Particle Titer (i.u./mL)	
		Normal Mouse	No serum
		Serum	
	Virosome Supernatant	20	440
	VRP-GFP	0	530
			890

30 The infectivity of the virosomes was inhibited similar to that of VRP-GFP,

indicating that the virosome particles were enveloped by the E1 and E2 glycoproteins.

These examples clearly demonstrate the ability to produce infectious virosome particles comprising replicon RNA enveloped with only the alphavirus E1 and E2 glycoproteins. Testing confirmed that these virosomes are infectious agents, but that they undergo only a single round of replication, as indicated by the inability to passage the agent. In addition, the agents contained the E1 and E2 glycoproteins, as evidenced by the ability to block infection with only VEE specific serum. Finally, the infectious RNA is protected from RNase enzymatic digestion, indicating an enveloped particle.

10

The natural lipid content in BHK cells is primarily non-cationic. Virosomes made in a completely cell free system can be made by using one or more non-cationic lipids, such as lecithin (phosphatidylcholine).

15

EXAMPLE 10

PHASE I CLINICAL PROTOCOL

Phase I Safety and Immunogenicity Trial of an HIV Subtype C Gag-VEE Replicon Particle Vaccine in HIV-1 Seronegative Human Subjects

A Phase I trial is conducted to evaluate the safety and immunogenicity of the HIV Gag-VRP prototype vaccine component in healthy seronegative adult volunteers. The doses are selected based on preclinical studies in rodents and nonhuman primates. The schedule mimics previous preclinical efficacy studies with the SIV model that demonstrated the capacity of SIV-VRP to induce SIV specific neutralizing antibodies and CTL.

Purpose: To evaluate the candidate vaccine component in an open-labeled, placebo-controlled study.

30

Subjects: Healthy adult volunteers without a history of identifiable high-risk behavior for HIV-1 infection as determined by a comprehensive screening questionnaire.

No. Subjects: 40

5

Route: Subcutaneous injection

Scheme: The volunteers are arranged in four groups, ten subjects per group. In each group, two subjects receive a placebo, while the other eight subjects receive either 10^4 , 10^6 , 10^7 , or 10^8 i.u. of HIV-Gag-VRPs. Subjects are vaccinated on day 0, day 30, and day 120.

10

Estimated Duration: Forty weeks

A. SELECTION of SUBJECTS

15

Subjects are healthy HIV-1 seronegative adults who fully comprehend the purpose and details of the study as described in the informed consent. Subjects whom either themselves or whose sexual partners have identifiable higher risk behavior for HIV-1 infection are not eligible. Higher risk behavior is determined by a prescreen series of questions designed to identify risk factors for HIV-1 infection. An assessment of absolute exclusion criteria using the self-administered and interview questions is conducted. Subsequently, investigators proceed with phlebotomy, history and physical examination, and final questions regarding sexual behavior and other practices. Eligibility determinations for the trial depend on results of laboratory tests and answers to these self-administered and interview questions.

20

The criteria used to define low risk behavior are as follows:

EITHER ALL OF THE FOLLOWING:

1. No newly acquired higher risk associated STD in the last six months
2. No possibly safe or unsafe sex with a known HIV+ individual or an active

25

- injection drug user in the past six months
- 3. No unsafe sexual activity
- 4. Possibly safe sexual activity with two or fewer partners within the last six months
- 5 5. No injection drug use

OR BOTH OF THE FOLLOWING:

- 1. Mutually monogamous relationship with a known or presumed HIV seronegative partner for the last six months
- 10 2. No injection drug use

A.1 Inclusion Criteria

Age: 18-60

- 15 Sex: Male or Female [*For females, negative pregnancy test at time of entry and assurance that adequate birth control measures will be used for one month prior to immunization and the duration of the study*]

Normal history and physical examination

Lower risk sexual behavior as defined above.

- 20 Normal complete blood count and differential defined as:

- Hematocrit 34% for women; 38% for men
- White count 3500 cells/mm³ with normal differential
- Total lymphocyte count 800 cells/mm³
- Absolute CD4 count 400 cells/mm³

- 25 - Platelets (150,000-550,000)

Normal ALT (~ 1.5 x institutional upper normal limit) and creatinine (1.6 mg/dl)

Normal urine dipstick with esterase and nitrite

Negative for hepatitis B surface antigen

- 30 Negative ELISA for HIV within eight weeks of immunization

Availability for follow-up for planned duration of the study (68 weeks)

A viable EBV transformed autologous B cell line

A.2 Exclusion Criteria

5

History of immunodeficiency, chronic illness, malignancy, autoimmune disease, or use of immunosuppressive medications

10 Medical or psychiatric condition or occupational responsibilities which preclude subject compliance with the protocol

Subjects with identifiable higher risk behavior for HIV infection as determined by screening questionnaire designed to identify risk factors for HIV infection; specific exclusions include:

- 15 History of injection drug use within the last 12 months prior to enrollment.
- Higher risk sexual behavior defined as one or more of the following behaviors:
1. A newly acquired higher risk associated STD within the past six months
 2. Possibly safe or unsafe sex with a known HIV+ individual in the past six months
 - 20 3. Possibly safe sexual activity with twelve or more partners in the past six months
 4. Unsafe sexual activity with four or more partners within the past six months.
- 25 Live attenuated vaccines within 60 days of study [NOTE: Medically indicated subunit or killed vaccines (e.g., influenza, pneumococcal) are not exclusionary, but should be given at least two weeks away from test article immunizations.]
- Use of experimental agents within 30 days prior to study
- Receipt of blood products or immunoglobulin in the past six months
- 30 Active syphilis [NOTE: If the serology is documented to be a false positive or

due to a remote (>six months) treated infection, the volunteer is eligible]

Active tuberculosis [NOTE: Volunteers with a positive PPD and a normal chest X-ray showing no evidence of TB and not requiring INH therapy are eligible.]

History of anaphylaxis or other serious adverse reactions to vaccines

5 Prior receipt of HIV vaccines or a placebo recipient in an HIV vaccine trial

Pregnant or lactating women

B. SAFETY and IMMUNOGENICITY MONITORING

10 Safety is evaluated by monitoring volunteers for adverse reactions during the course of the trial. Volunteers are followed for a total of 26 weeks post-final inoculation. The main toxicity associated with the subcutaneous injection in this study is that associated with subcutaneous injection of any immunogen, i.e., pain, redness and swelling at the injection site, as well as the possibility of fever, chills, aches and pains
15 and perhaps fatigue.

Safety monitoring includes periodic review of data from the trial with particular emphasis on monitoring for adverse reactions including the following evaluations:

Hematologic: CBC, differential, platelets

20 Hepatic/renal: ALT, creatinine, urinalysis

Neurologic: headache, paralysis, anxiety, confusion, weakness, tremors.

Systemic symptoms: fever, gastrointestinal complaints, myalgia, malaise, fatigue, headache, anaphylaxis, immune complex disease, and other hypersensitivity reactions

25 Local toxicity at the site of injection: e.g., pain, tenderness, erythema, regional lymphadenopathy, limitation of limb movement

The immunogenicity monitoring includes the following immunological assays, all utilizing HIV Subtype C based reagents:

Humoral responses:

HIV Subtype C Gag-specific ELISA

Anti-VEE ELISA

5 Cellular immune responses:

Standard cell-killing assay (i.e., chromium release) to measure CD8+ Gag-specific CTL activity

ELISPOT assay to measure IFN- γ

10 Mucosal immune responses:

Standardized assay for assessment of Gag-specific IgA

- Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference
- 15 into this application in order to more fully describe the state of the art to which this invention pertains.

REFERENCES

- 20 Barany F. 1985. Single-stranded hexameric linkers: a system for in-phase insertion mutagenesis and protein engineering. *Gene* 37(1-3):111-23.
- Betts, M. R., J. Krowka, C. Santamaria, K. Balsamo, F. Gao, G. Mulundu, C. Luo, N. N'Gandu, H. Sheppard, B. H. Hahn, S. Allen and J. A. Frelinger. 1997. Cross-clade
- 25 human immunodeficiency virus (HIV)-specific cytotoxic T-lymphocyte responses in HIV-infected Zambians. *J. Virol.* 71:8908-8911.
- Caley, I. J., M.R. Betts, D.M. Irlbeck, N. L. Davis, R. Swanstrom, J.A. Frelinger and R. E. Johnston. 1997. Humoral, mucosal and cellular immunity in response to an HIV-
- 30 1 vaccine candidate. *J. Virol.* 71:3031-3038.

Davis *et al.* 1980. In: Microbiology, 3d ed., p. 132.

Davis, N.L., L.V. Willis, J.F. Smith and R.E. Johnston. 1989. *In vitro* synthesis of infectious Venezuelan equine encephalitis virus RNA from a cDNA clone: Analysis of
5 a viable deletion mutant. *Virology* **171**:189-204.

Davis, N. L., L. V. Willis, J. F. Smith, G. Greenwald and R. E. Johnston. 1990. *In vitro* synthesis of infectious VEE virus RNA from a cDNA clone: Analysis of a viable deletion mutant and mutations affecting virulence. In: Vaccines 90, Cold Spring Harbor
10 Press, Cold Spring Harbor, NY. pp. 109-113.

Davis, N. L., N. Powell, G. F. Greenwald, L. V. Willis, B. J. Johnson, J. F. Smith and R. E. Johnston. 1991. Attenuating mutations in the E2 glycoprotein gene of Venezuelan equine encephalitis virus: Construction of single and multiple mutants in a
15 full-length cDNA clone. *Virology* **183**:20-31.

Davis, N. L., K. W. Brown, G. F. Greenwald, A. J. Zajac, V. L. Zacny, J. F. Smith and R. E. Johnston. 1995. Attenuated mutants of Venezuelan equine encephalitis virus containing lethal mutations in the PE2 cleavage signal combined with a second-site
20 suppressor mutation in E1. *Virology* **212**:102-110.

Davis, N. L., K. W. Brown and R. E. Johnston. 1996a. A viral vaccine vector that expresses foreign genes in lymph nodes and protects against mucosal challenge. *J. Virol.* **70**:3781-3787.

25 Davis, N. L., P. Pushko, K. W. Brown, P. C. Charles, I. J. Caley, M. Parker, G. Ludwig, J. F. Smith and R. E. Johnston. 1996b. Immunization against influenza with attenuated Venezuelan equine encephalitis virus vectors. In: Options for the Control of Influenza, III, L. E. Brown and A. W. Hampson, eds. Elsevier, Amsterdam. pp.803-
30 809.

- Davis, N. L., I. J. Caley, K. W. Brown, M.R. Betts, D.L. Irlbeck, K.M. McGrath, M.J. Connell, D.C. Montefiori, J.A. Frelinger, R. Swanstrom, P.R. Johnson and R. E. Johnston. 2000. Vaccination of macaques against pathogenic simian immunodeficiency virus with Venezuelan equine encephalitis virus replicon particles. *J. Virol.* 74:371-378
- 5 Grieder, F. B., N. L. Davis, J. F. Aronson, P. C. Charles, D. C. Sellon, K. Suzuki and R. E. Johnston. 1995. Specific restrictions in the progression of Venezuelan equine encephalitis virus induced disease resulting from single amino acid changes in the glycoproteins. *Virology* 206:994-1006.
- 10 Hevey, M., D. Negley, P. Pushko, J. Smith and A. Schmaljohn. 1998. Marburg virus vaccines based upon alphavirus replicons protect guinea pigs and nonhuman primates. *Virology* 251:28-37.
- 15 Hirsch, V., T.R. Fuerst, G. Sutter, M.W. Carroll, L. C. Yang, S. Goldstein *et al.* 1996. Patterns of viral replication correlate with outcome in SIV-infected macaques: effect of prior immunization with a trivalent SIV vaccine in modified vaccinia virus Ankara. *J. Virol.* 70:3741-3752.
- 20 Johnston, Robert E. and Jonathan F. Smith. 1988. Selection for accelerated penetration in cell culture co-selects for attenuated mutants of Venezuelan equine encephalitis virus. *Virology* 162:437-443.
- Johnston, R. E. and C. J. Peters. 1996. *Alphaviruses*. In: *Virology*, Third Edition, B. N. Fields, D. M. Knipe and P. M. Howley, eds., Raven Press, New York. pp. 843-898.
- 25 Kinney, R.M., B.J.B. Johnson, J.B. Welch, K.R. Tsuchiya and D.W. Trent. 1989. The full-length nucleotide sequences of the virulent Trinidad donkey strain of Venezuelan equine encephalitis virus and its attenuated vaccine derivative, strain TC-83. *Virology* 170:19-30.

- Kinney, R.M., G-J. Chang, K. R. Tsuchiya, J. M. Sneider, J. T. Roehrig, T. M. Woodward and D. W. Trent. 1993. Attenuation of Venezuelan equine encephalitis virus strain TC-83 is encoded by the 5'-noncoding region and the E2 envelope glycoprotein. *J. Virol.* 67:1269-1277.
- 5 Kunkel. 1985. *Proc. Natl. Acad. Sci. USA* 82:488.
- Paredes, A.M., D.T. Brown, R. Rothnagel, W. Chiu, R.J. Schoepp, R.E. Johnston and B.V.Prasad. 1993. Three-dimensional structure of a membrane-containing virus. *Proc.*
- 10 *Natl. Acad. Sci., USA* 90:9095-9099.
- Pushko, P., M. Parker, G. V. Ludwig, N. L. Davis, R. E. Johnston and J. F. Smith. 1997. Replicon-helper systems from attenuated Venezuelan equine encephalitis virus: expression of heterologous genes *in vitro* and immunization against heterologous
- 15 pathogens *in vivo*. *Virology* 239:389-401.
- Rosenberg, A.H., *et al.* 1987. Vectors for selective expression of cloned DNAs by T7 RNA polymerase. *Gene*. 56(1): p. 125-35.
- 20 Schlesinger, S. and M.J. Schlesinger. 1990. *Replication of Togaviridae and Flaviviridae*. In: Virology, Fields, B.N. and Knipe, D.M. (eds.) Raven Press. pp. 697-711.
- Strauss *et al.* 1990. *Seminars in Virology* 1:347.
- 25 Strauss, J.H. and E.G. Strauss. 1994. The alphaviruses: Gene expression, replication, and evolution. *Micro. Rev.* 58:491-562.
- Studier, F.W., *et al.* 1990. Use of T7 RNA polymerase to direct expression of cloned
- 30 genes. 1990. *Methods Enzymol.* 185:60-89.

What is claimed is:

1. A composition comprising two or more isolated nucleic acids selected from the group consisting of an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity.
2. A composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle.
3. A composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag*

gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle, and further wherein the alphavirus replicon particles comprise a replicon RNA or at least one structural protein which comprises one or more attenuating mutations.

4. A method of making the population of alphavirus replicon particles of claim 2 comprising:
 - A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:
 - (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the

helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains

no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles of claim 2.

5. The method of claim 4, wherein the alphavirus replicon RNA of at least one of the first helper cell, the second helper cell and the third helper cell comprises sequence encoding at least one alphavirus structural protein and wherein the first helper RNA and the one or more additional helper RNA(s) in the at least one of the first helper cell, the second helper cell and the third helper cell, encodes at least one other alphavirus structural protein not encoded by said replicon RNA.

6. A method of making the population of alphavirus replicon particles of claim 3, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein;

and

- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particle, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and

furthermore not encoding at least one other alphavirus structural protein;
and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof is modified to inhibit reverse transcriptase activity, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and

furthermore not encoding at least one other alphavirus structural protein;
and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles of claim 3.

7. The method of claim 6, wherein the alphavirus replicon RNA of at least one of the first helper cell, the second helper cell and the third helper cell comprises sequence encoding at least one alphavirus structural protein and wherein the first helper RNA and the one or more additional helper RNA(s) in the at least one of the first helper cell, the second helper cell and the third helper cell, encodes at least one other alphavirus structural protein not encoded by said replicon RNA.

8. The method of claim 6, wherein only at least one of the first population of alphavirus particles, the second population of alphavirus particles and the third population of alphavirus particles comprises particles wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations.
9. A population of alphavirus replicon particles produced by the method of claim 4.
10. A population of alphavirus replicon particles produced by the method of claim 6.
11. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 1 in a pharmaceutically acceptable carrier.
12. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 2 in a pharmaceutically acceptable carrier.
13. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 3 in a pharmaceutically acceptable carrier.
14. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 9 in a pharmaceutically acceptable carrier.
15. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the

population of claim 10 in a pharmaceutically acceptable carrier.

16. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 1 in a pharmaceutically acceptable carrier.

17. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 2 in a pharmaceutically acceptable carrier.

18. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 3 in a pharmaceutically acceptable carrier.

19. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 9 in a pharmaceutically acceptable carrier.

20. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 10 in a pharmaceutically acceptable carrier.

21. A composition comprising two or more isolated nucleic acids selected from the group consisting of an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof

of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof.

22. A composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle.

23. A composition comprising a population of alphavirus replicon particles comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a

modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon particle, and further wherein the alphavirus replicon particles comprise a replicon RNA or at least one structural protein which comprises one or more attenuating mutations.

24. A method of making the population of alphavirus replicon particles of claim 22, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and

(c) collecting the alphavirus particles from the helper cells;

C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

(b) producing the alphavirus particles in the helper cell; and

(c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles of claim 22.

25. The method of claim 24, wherein the alphavirus replicon RNA of at least one of the first helper cell, the second helper cell and the third helper cell comprises sequence encoding at least one alphavirus structural protein and wherein the first helper RNA and the one or more additional helper RNA(s) in the at least one of the first helper cell, the second helper cell and the third helper cell, encodes at least one other alphavirus structural protein not encoded by said replicon RNA.

26. A method of making the population of alphavirus replicon particles of claim 23, comprising:

A) (a) providing a first helper cell for producing a first population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:

(i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

(ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and

(iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the first population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells;

B) (a) providing a second helper cell for producing a second population of infectious, replication defective alphavirus particle, comprising in an alphavirus-permissive cell:

- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein

- not encoded by said first helper RNA;
- and with at least one of said helper RNAs lacking an alphavirus packaging signal;
- wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the second population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;
- (b) producing the alphavirus particles in the helper cell; and
 - (c) collecting the alphavirus particles from the helper cells;
- C) (a) providing a third helper cell for producing a third population of infectious, replication defective alphavirus particles, comprising in an alphavirus-permissive cell:
- (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;
 - (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
 - (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional

helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;
and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the third population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture, and further wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cells; and

D) combining the first population of alphavirus particles produced from the first helper cell, the second population of alphavirus particles produced from the second helper cell and the third population of alphavirus particles produced from the third helper cell, thereby producing the population of alphavirus replicon particles of claim 23.

27. The method of claim 26, wherein the alphavirus replicon RNA of at least one of the first helper cell, the second helper cell and the third helper cell comprises sequence encoding at least one alphavirus structural protein and wherein the first helper RNA and the one or more additional helper RNA(s) in the at least one of the first helper cell, the second helper cell and the third helper cell, encodes at least one other alphavirus structural protein not encoded by said replicon RNA.

28. The method of claim 26, wherein only at least one of the first population of alphavirus particles, the second population of alphavirus particles and the third population of alphavirus particles comprises particles wherein at least one of said

replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations.

29. A population of alphavirus replicon particles produced by the method of claim 24.

30. A population of alphavirus replicon particles produced by the method of claim 26.

31. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 21 in a pharmaceutically acceptable carrier.

32. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 22 in a pharmaceutically acceptable carrier.

33. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 23 in a pharmaceutically acceptable carrier.

34. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 29 in a pharmaceutically acceptable carrier.

35. A method of inducing an immune response to human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 30 in a pharmaceutically acceptable carrier.

36. A method of treating or preventing infection by human immunodeficiency

virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 21 in a pharmaceutically acceptable carrier.

37. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 22 in a pharmaceutically acceptable carrier.

38. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 23 in a pharmaceutically acceptable carrier.

39. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 29 in a pharmaceutically acceptable carrier.

40. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the population of claim 30 in a pharmaceutically acceptable carrier.

41. An alphavirus replicon virosome comprising an alphavirus replicon RNA encapsidated by a lipid bilayer comprising alphavirus glycoproteins, E1 and E2.

42. The virosome of claim 41, wherein the alphavirus glycoproteins are Venezuelan Equine Encephalitis glycoproteins E1 and E2.

43. A method of producing the alphavirus replicon virosome of claim 41, comprising:

a) combining alphavirus replicon RNA, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced.

44. An alphavirus replicon virosome produced from the method of claim 43.

45. A method of eliciting an immune response in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon virosome of claim 41 in a pharmaceutically acceptable carrier.

46. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon virosome of claim 41, wherein the virosome comprises alphavirus replicon RNA encoding one or more HIV immunogens.

47. A composition comprising a population of alphavirus replicon virosomes comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon virosome.

48. A composition comprising a population of alphavirus replicon virosomes

comprising two or more isolated nucleic acids selected from the group consisting of 1) an isolated nucleic acid encoding an *env* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, 2) an isolated nucleic acid encoding a *gag* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, and 3) an isolated nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in inactivation of reverse transcriptase activity in the *pol* gene product or immunogenic fragment thereof, and wherein the nucleic acids are each contained within a separate alphavirus replicon virosome.

49. A method of producing the population of alphavirus replicon virosomes of claim 47, comprising:

A) (a) producing a first population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *env* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;

B) (a) producing a second population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *gag* gene product or immunogenic fragment thereof, wherein the *gag* gene product or immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes

are produced;

C) (a) producing a third population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding the *pol* gene product or immunogenic fragment thereof, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced; and

D) combining the first population of alphavirus replicon virosomes, the second population of alphavirus replicon virosomes and the third population of alphavirus replicon virosomes to produce the population of alphavirus replicon virosomes of claim 47.

50. A method of producing the population of alphavirus replicon virosomes of claim 48, comprising:

A) (a) producing a first population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *env* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;

B) (a) producing a second population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding and *gag* gene product or immunogenic fragment thereof, wherein the *gag* gene product or

immunogenic fragment thereof is modified to inhibit formation of virus-like particles containing the *gag* gene product or the immunogenic fragment thereof and their release from a cell, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and

b) gradually removing detergent, whereby alphavirus replicon virosomes are produced;

- C) (a) producing a third population of alphavirus replicon virosomes by combining alphavirus replicon RNA comprising nucleic acid encoding the *pol* gene product or immunogenic fragment thereof, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in inactivation of reverse transcriptase activity in the *pol* gene product or immunogenic fragment thereof, alphavirus glycoproteins E1 and E2, non-cationic lipids and detergent; and
- b) gradually removing detergent, whereby alphavirus replicon virosomes are produced; and
- D) combining the first population of alphavirus replicon virosomes, the second population of alphavirus replicon virosomes and the third population of alphavirus replicon virosomes to produce the population of alphavirus replicon virosomes of claim 48.

51. A method of eliciting an immune response in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 47, in a pharmaceutically acceptable carrier.

52. A method of eliciting an immune response in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 48, in a pharmaceutically acceptable carrier.

53. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 47, in a pharmaceutically acceptable carrier.

54. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 47, in a pharmaceutically acceptable carrier.

55. A composition comprising heparin affinity-purified alphavirus replicon particles, wherein the alphavirus replicon particles comprise at least one structural protein which comprises one or more attenuating mutations.

56. A method of preparing the heparin affinity-purified alphavirus particles of claim 55, comprising:

- a) producing alphavirus replicon particles, wherein the alphavirus replicon particles comprise at least one structural protein which comprises one or more attenuating mutations;
- b) loading the alphavirus replicon particles of step (a) in a heparin affinity chromatography column; and
- c) collecting the fraction from the column which contains the heparin affinity-purified alphavirus replicon particles.

57. A composition produced by the method of claim 56.

58. A method of producing VRP for use in a vaccine comprising:

- a) producing a plasmid encoding the nucleotide sequence of an alphavirus replicon RNA;
- b) producing a plasmid encoding the nucleotide sequence of one or more helper RNAs;
- c) transcribing the plasmids of steps (a) and (b) into RNA *in vitro*;
- d) electroporating the RNA of step (c) into a Vero cell line; and
- e) purifying VRP from the Vero cell line of step (d) by heparin affinity chromatography.

59. The method of claim 58, wherein the VRP is produced in large-scale.
60. VRP produced by the method of claim 59.
61. An isolated nucleic acid encoding a *pol* gene product or immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof.
62. A composition comprising the nucleic acid of claim 61.
63. A vector comprising the nucleic acid of claim 61.
64. A cell comprising the vector of claim 63.
65. An alphavirus replicon particle comprising the nucleic acid of claim 61.
66. A method of making the alphavirus replicon particle of claim 65, comprising
- a) providing a helper cell for producing an infectious, defective alphavirus particle, comprising in an alphavirus-permissive cell:
 - (i) an alphavirus replicon RNA, wherein the replicon RNA comprises an alphavirus packaging signal and a nucleic acid encoding a *pol* gene product or an immunogenic fragment thereof of a human immunodeficiency virus, wherein the *pol* gene product or immunogenic fragment thereof comprises a modification resulting in deletion or inactivation of integrase, RNase H and reverse transcriptase functions in the *pol* gene product or immunogenic fragment thereof, and wherein the replicon RNA lacks sequences encoding alphavirus structural proteins;

- (ii) a first helper RNA separate from said replicon RNA, said first helper RNA encoding at least one alphavirus structural protein and furthermore not encoding at least one other alphavirus structural protein; and
- (iii) one or more additional helper RNA(s) separate from said replicon RNA and separate from said first helper RNA, said additional helper RNA(s) encoding at least one other alphavirus structural protein not encoded by said first helper RNA;

and with at least one of said helper RNAs lacking an alphavirus packaging signal;

wherein the combined expression of the alphavirus replicon RNA and the helper RNAs produces an assembled alphavirus particle which is able to infect a cell, and is unable to complete viral replication, and further wherein the population contains no detectable replication-competent alphavirus particles as determined by passage on permissive cells in culture;

- (b) producing the alphavirus particles in the helper cell; and
- (c) collecting the alphavirus particles from the helper cell.


67. The method of claim 66, wherein at least one of said replicon RNA, said first helper RNA, and said one or more additional helper RNA(s) comprises one or more attenuating mutations.

68. An alphavirus replicon particle produced according to the method of claim 66.


69. An alphavirus replicon particle produced according to the method of claim 67.


70. A method of inducing an immune response in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 62 in a pharmaceutically acceptable carrier.


71. A method of inducing an immune response in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon particle of claim 65 in a pharmaceutically acceptable carrier.
72. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the composition of claim 62 in a pharmaceutically acceptable carrier.
73. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of the alphavirus replicon particle of claim 65 in a pharmaceutically acceptable carrier.
74. A method of inducing an immune response in a subject, comprising administering to the subject an immunogenic amount of a composition comprising the alphavirus replicon particles of claim 65 in a pharmaceutically acceptable carrier.
75. A method of treating or preventing infection by human immunodeficiency virus in a subject, comprising administering to the subject an immunogenic amount of a composition comprising the alphavirus replicon particles of claim 65 in a pharmaceutically acceptable carrier.


**FIG. 1**

2/15


**FIG. 2**

**FIG. 3**

**FIG. 4**

**FIG. 5**

6/15

**FIG. 6**

7/15








FIG. 7

**FIG. 8**

9/15

**FIG. 9A****FIG. 9B****FIG. 9C**




FIG. 10




FIG. 11




FIG. 12






FIG. 13

**FIG. 14**

**FIG. 15**

SEQUENCE LISTING

<110> AlphaVax, Inc. and University of North Carolina

Olmsted, Robert
Keith, Paula
Drygan, Sergey
Daley, Ian
Maughan, Maureen
Johnston, Robert
Davis, Nancy
Swanstrom, Ronald

<120> ALPHAVIRUS VECTORS AND VIROSOMES WITH MODIFIED HIV GENES FOR USE AS VACCINES

<130> 01113.0001P1

<150> 60/216,995

<151> 2000-07-07

<160> 19

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 12523

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 1

atggggcggcg	catgagagaa	gccagacca	attacctacc	caaaatggag	aaagtccacg	60
ttgacatcga	ggaagacagc	ccattcctca	gagcttttga	gcggagcttc	cgcagtttg	120
aggtagaagc	caagcaggtc	actgataatg	accatgctaa	tgccagagcg	ttttcgcatc	180
tggtttcaaa	actgatccaa	acggaggtgg	acccatccga	cacgatcctt	gacattggaa	240
gtggcggccg	ccgcagaatg	tattctaagc	acaagtatca	ttgtatctgt	ccgatgagat	300
gtggggaaga	tccggacaga	ttgtataagt	atgcaactaa	gctgaagaaa	aaactgaagg	360
aaataaactga	taagggaattg	gacaagaaaa	tgaaggagct	cgccgcgctc	atgagcgacc	420
ctgacctgga	aactgagact	atgtgcctcc	acgacgacga	gtcgtgtcgc	tacgaagggc	480
aagtgcgtgt	ttaccaggat	gtatacgcgg	ttgacggacc	gacaagcttc	tatcaccaag	540
ccaataaggg	agtttagatc	gcctactgga	taggctttga	caccacccct	tttatgttta	600
agaacttggc	tggagcatat	ccatcatact	ctaccaactg	ggccgacgaa	accgtgttaa	660
cggctcgtaa	cataggcccta	tgacgctctg	acggttatgga	gcggtcacgt	agagggatgt	720
ccattcttag	aaagaagtat	ttgaaaccaa	ccaacaatgt	tcattctctc	gttggtctga	780
ccatctacca	cyagaagagg	gacttactga	ggagctggca	cctgcgltct	gtatttcaet	840
tacgtggcaa	gcaaaattac	acatgtcggt	gtgagactat	agttagtgtg	gacgggtacg	900
tcgttaaaag	aatagctatc	agtcacggcc	tgtatgggaa	gccttcaggc	tatgctgcta	960
cagtgacacc	cgagggatcc	ttgtgtctga	aagtgcagaa	cacattaaac	ggggagaggg	1020
tcctctttcc	cggtgtgcacg	tatgtgccag	ctacatttgt	tgaccaaatg	actggcctac	1080
tggcaacaga	tgtcagtgcg	gacgacgcgc	aaaaactgct	ggttggcgct	aaccagcgta	1140

tagtgcgtcaa	cggctgcgacc	cagagaaaaca	ccaataccatt	gaaaaattac	cttttgcccg	1200
tagtgcccca	ggcatttgct	agggtgggcaa	aggaatataa	ggaagatcaa	gaagatgaaa	1260
ggccactagg	actacgagat	agacaggttag	tcactgggggt	ttgttggggt	ttttagaagc	1320
acaagataac	atctatttat	aagcgcccg	ataccacaaac	catcatcaaa	gtgaacagcg	1380
atttccactc	atttcgtctg	cccaggatag	gcagtaaac	attggagatc	ggggtgagaa	1440
caagaattcag	gaaaaattgt	gaggagcaca	agcgcgcgtc	acctctcat	accgcgcagg	1500
acgtacaaga	agctaatgct	gcagccgatg	aggctaagga	gggtgcgtgaa	gcgcaggagtg	1560
tgcgcgcagc	tctaccactc	ttggcagctg	atgttgaggga	gcccaactctg	gaagccagtg	1620
tcgacttgat	gttacaagag	gctggggccg	gctcagtgga	gacacctcgt	ggcttgataa	1680
agggtaccag	ctacgctggc	gaggacaaga	tcggctctta	cgctgtgctt	tctccgcagg	1740
ctgtactcaa	gagtgaaaaa	ttatcttgca	tcaccctctc	cgctgaaaca	gtcatagtga	1800
taacacactc	tggccgaaaa	gggcgttatg	cgttggaacc	ataccatggt	aaagtatgct	1860
tgcagagggg	acatgcaata	ccgctccagg	accttcaagc	tcctgagtga	agtgccacga	1920
ttgtgtacaa	cgaacgtgag	ttcgtaaaca	ggtaacctga	ccatattgccc	acacatggag	1980
gagcgctgaa	cactgatgaa	gaattattaca	aaactgtcaa	gccacgcgag	cacgacggcg	2040
aataacctgta	cgacatcgac	aggaacaacgt	gcgtcaagaa	agaactagtc	actggcgtag	2100
ggctcacagg	cgagctgggt	gatcctccct	tcactgaatt	cgctcacgag	agctctgagaa	2160
cacgaccacg	cgctccttac	caagtaccaa	ccataggggt	gtatggcggt	ccaggatcag	2220
gcaagctcgt	catctataa	agcgagtcga	acgcaaaaaga	tctagtgggt	agcgcocaag	2280
aagaaaaactg	tgcagaatt	ataaggcgag	tcaagaaat	gaaagggtcg	gagctcaatg	2340
ccagaactgt	ggactcagtg	ctcttgaatg	gatgcaaaaa	ccccgtagag	acctgtgata	2400
ttga cgaagc	ttttgctgtg	catgcaggta	ctctcagagc	gctcatagcc	attataaagc	2460
ctaaaaaggg	agtgctctgc	ggggatccca	aacagtgcgg	tttttttaac	atgatgtgcc	2520
tgaagatgtca	ttttaaccac	gagatttgca	cacaagctct	ccacaaaagc	actctctgcc	2580
gttgactcaa	atctgtgact	ctcgtcgtct	caaccttgct	ttacgacaaa	aaaatagaaa	2640
cgcagatccc	gaaagagact	agaattgtga	ttgacactac	cggcagtagc	agacctagc	2700
aggagcatct	catctcact	tgtttcagag	gggtgggtgaa	gcagttgcaa	atagattaca	2760
aaggcacaaga	aataatgagc	gcagctgcct	ctcaagggtc	gacccgtaaa	gggtgtgatg	2820
ccgttcggtg	caagggtgaat	gaaaaactcc	tgtaacgacc	cacctcagaa	catgtgaagc	2880
tctactcgac	ccgcacggag	gacgcgcatg	tgtggaaaac	actagccggc	gaccatcgga	2940
taaaaacact	gactgccaa	tacctgggga	atcttcaactg	cacgatagag	gagtgccaag	3000
cagagcatga	tgccatcatg	aggcacatct	tggagagacc	ggacccctacc	gacgtctccc	3060
agaa taaggg	aaacgtgtgt	tgggccaagg	cttttagtgcc	gggtgctgaag	acogctggca	3120
tagacatgac	cactgaaaca	tggaaacactg	tggattattt	tgaacccgag	aaagctcaet	3180
cagcagagat	gaatttgaa	caactatgct	tgaagttctt	tggactcgat	ctggaatccg	3240
gtctattttc	tgcaccact	gttcogttat	ccattaggaa	taactactgg	gataactccc	3300
cgctgcctaa	catgtacggg	ctgaataaag	aagtgttcg	tcagctctct	cgcaggtaac	3360
cacaactgcc	tcgggcagtt	gccactggaa	gagcttatga	catgaacact	gagctactgc	3420
gcgaattatga	tcgcgcata	aacctagtac	ctgtaaacag	aagactgcct	catgctttag	3480
tcttccccca	taatgaacac	ccacagagtg	aettttcttc	atctgcagc	aaattgaagg	3540
gcgaagactgt	cctgttggtc	ggggaaaagt	tgtccgtccc	aggcaaaagt	gttgactcgtg	3600
tgtcagaccc	gcctgaggtc	accttcagag	ctcggctgga	tttaggcctg	ccaggtgatg	3660
tgcgccaaata	tgacataata	tttgttaatg	tgaaggaccc	atataaatac	catcaactct	3720
agcaagtgtga	agacatgcc	attaagctta	gcactgtgac	caagaaagct	tgctctgcact	3780
tgaatcccg	cggaaacctg	gtcagcatag	gttatggtta	cgctgacagg	gccagcgaaa	3840
gcacatctgg	tgctatagcg	cggcagttca	agttttcccg	ggatgcacaa	cgcgaatcct	3900
catctgaaga	gacggaagtt	ctgtttgtat	tcatgggta	cgatgcgaag	cccgctagct	3960
acaa tccctta	caagctttca	tcaaccttga	ccaacattta	tacaggttcc	agactccacc	4020
aagcgggatg	tgcacctcca	tatcatgtgg	tgcgagggga	tattgccacg	gccacgcgaag	4080
gagtgattat	aaatcgtcgt	aacagcaaa	gacaacctgt	cggagggggt	tgcggagcgc	4140
tgtataagaa	gttcccgcaa	agcttcgalt	tacagccgat	cgaagttaga	aaagcgcgac	4200
tggtcacaa	tgcagctaaa	catatcattc	atgcgctagg	accaaacttc	aacaagttgt	4260
cggaggttga	aggtgcacaa	cagttggcag	aggettattg	gtccatcgct	aagattgtca	4320
acgataacaa	ttacagatca	gtagcgattc	catctgttgc	accgcgacac	tttctcgga	4380
acaagatcgc	actaacccaa	tcatgtgaac	atcttgcagc	agcttttagac	accactgatg	4440

cagatgtagc	catatactgc	agggacaaga	aatgggaaat	gactctcaag	gaagcagtg	4500
ctagaggaga	agcagtgagg	gagatatgca	tatcogaaga	ctcttcagtg	acagaaactg	4560
atgcagagct	gttgagggtg	catcogaaga	gtctcttggt	tggaaagaa	gtctcagaga	4620
caagcgtagg	caaaactttc	tcatatttgg	aaggagcaaa	gtttccaccg	gcggccaaag	4680
atatagcaga	ataataatg	atgtggcccg	ttgcaaagg	ggccaatgag	caggtagtga	4740
tgtatatctc	cggagaaaag	atgagcagta	ttaggtcgaa	atgcccgcgc	gaagagctgg	4800
aagctccacc	accactatgc	acgtctgcct	gcttctgcat	ccatgcccgc	actccagaaa	4860
agctacagcg	ctaaagacc	tcaogtcagg	acaaattatc	tgtgtgtcga	tcctttccat	4920
tgccgaagta	tagaatcaat	ggtgtgcaga	agatcccaat	ctcccagcct	atatgttctt	4980
caccgaaggt	gcctgcgtat	atcatcccaa	ggaagtatct	cgtggaaaca	ccaocggtag	5040
acgagacatc	ggagccatcg	gcagagaaac	aatccacaga	ggggacacct	gaacaaaccac	5100
catctataac	cgaggatgag	accaggagta	gaacgcctga	gcgatcatc	atcgaagagg	5160
aagaagaggga	tagcataagt	ttgctgtcag	atggcccagc	ccaccaggtg	ctgcaagtgc	5220
aggcagacat	tcacggggcg	ccctctgtat	ctagctcatc	ctggctccat	ccctatgcat	5280
ccgactttga	tgtggacagt	ttatccatc	ttgacacact	ggaggagagt	acogtgacca	5340
gcgggggcaac	gtcagccgag	actaatctct	acttcgcaaa	gagtagtgag	tttctggcgc	5400
gacccgtgccc	tgccgctcga	acagtattca	ggaacccctc	acatcccgct	ccgctgacaa	5460
gaacacgcgt	acttgcaccc	agcagggcct	gctcgagaac	cagcctagtt	tcaccccgcg	5520
caggcgtgaa	taggggtgatc	actagagagg	agctcgaggc	gcttaaccgc	taacgcaact	5580
ctagcaggtc	ggctctcgaga	accagcctgg	ttcccaacc	gccaggcgta	aatagggtga	5640
ttacagagaga	ggagttttgag	gcgttcgtag	cacaacaaca	atgacgggtt	gatgggggtg	5700
catatcatct	ttctctccag	accgggtcaag	ggcatttaca	acaaaataca	gtaaaggcaa	5760
cggtgctcat	cgaagtgtgt	ttggagagga	ccgaatttga	gatttctgat	gcocccgcgc	5820
tcgaacaaga	aaaagcaaaa	ttactacgca	agaaattaca	gttaaatccc	acacctgcta	5880
acagaagcag	ataccagatc	aggaaggttg	agacacatga	agccataaca	gctagacgta	5940
ttctgcgaag	ctcagggcat	tatttgaagg	cagaaggaaa	agtgaggtgc	taccgaaacc	6000
tgcatctctg	tcctttgtat	tcactatag	tgaacgcgtc	cttttcaacg	cccaaggtgc	6060
cagtggaagc	ctgtaacgcc	atgttgaaag	agaactttcc	gactgtggct	ttctactgta	6120
ttattccaga	gtacagatgc	tatttggaca	tggttgacgg	agcttcatgc	tgcttagaca	6180
ctgcacgttt	ttgcctcgca	aagctgcgca	gctttccaaa	gaacacctcc	tatttgggac	6240
ccacaatacg	atcgccagtg	ccttcagcga	tcagaaacac	gctcagaac	gtcctggcag	6300
ctgcacaaaa	aagaatttgc	aatgtcacgc	aaatgagaga	attgccccta	ttggattcgg	6360
cgcccttttaa	tgtggaatgc	ttcaagaaat	atgcgtgtaa	taatgaatat	tgggaaaagt	6420
ttaaagaaaa	cccatcagg	cttactgaag	aaaacgtgtg	aaattacatt	accaaattaa	6480
aaggacaaaa	agctgctgct	ctttttcgga	agacacataa	tttgaatatg	ttcaggagca	6540
taccaatgga	caggtttgta	atggacttaa	agagagacgt	gaaagtgtat	ccaggaaaca	6600
aacatactga	agaacggccc	aaggtacagg	tgatccagcg	tgccgatccg	atgcacaaa	6660
cgatctctgt	cggaaatccac	cgagagctgg	ttaggagatt	aaatggcgct	ctgctccoga	6720
acatctcatc	actgtttgat	atgtcggctg	aagactttga	cgctattata	gcogagcact	6780
tcagcctgg	ggatgtgtgt	ctggaaactg	acatcgctgc	gtttgataaa	agtgaggagc	6840
agcccatggc	ttcgaacggc	ttaatgattc	tggaaagact	aggtgtggac	cgagagctgt	6900
tcagcctgat	tgagcggcgt	ttcggcgaaa	ttcatcaat	acatttgccc	ataaataact	6960
aatttaaaat	cggaagcatg	atgaaatctg	gaatgttctc	caactgtgtt	gtgaacacag	7020
tcattaaact	tgtaatcgca	agcagagtgt	tgagagaaa	gctaaacoga	tcaccatgtg	7080
cagcatctcat	tggagatgac	aatatcgtga	aggaggtcaa	atccggacaaa	ttaatggcag	7140
acaggtggcgc	cacctgggtg	aatatggag	tcaagattat	agatgctgtg	gtggggcaga	7200
aagcgcctcta	ttctctggga	gggtttattt	tgtgtgaact	cgtgacgggc	acagcgtgoc	7260
gtgtggcaga	ccccataaaa	aggctgttta	agcttggcaa	acctctggca	gcagacgagt	7320
aaactgatga	tgacagagga	aggccattgc	atgaagagtc	aaacgcctgg	aaccaggtgg	7380
gtatctcttc	agagcgtgtg	aaggcagtag	aatcaagtag	tgaacacgta	ggaacttcca	7440
tcatagttat	ggccatgact	actctagcta	gcaggtgtta	atcatcagc	tacctgagag	7500
gggcccctat	aactctctac	ggctaaacgt	aatggactac	gacatagctc	agtcgcgcaa	7560
gtatggctgog	agagcgtcaa	tattaagagg	ggaaaaata	gataaatggg	aaaagattag	7620
gttaaaggcca	gggggaaaga	aacattatat	gttaaaacac	atagtatggg	cgagcgggga	7680
gctggaaaga	tttgcaacta	accctggcct	tttagaaca	tcagaaggat	gtaaacaaat	7740

aatgaacag	ctacaaccag	ctctccagac	aggaacagag	gaacttaaat	cattatacaa	7800
cacagatgca	actctctatt	gtgtacatga	aaagatagaa	gtacgagaca	ccaaggaagc	7860
cttagataag	atagagggaag	aacaaaacaa	atgtcagcaa	aaaacgcagc	aggcaaaagc	7920
ggctcagcgg	aaagtccagtc	aaaattatcc	tatagtgcag	aatctccaag	ggcgaatggc	7980
acatcaagcc	atatcaacct	gaaccttgaa	tgcattgggt	aaagtaatag	agaanaagcg	8040
tttttagccca	gaggtataac	ccatgtttac	agcattatca	gaaggagcca	ccccacaaga	8100
ttttaaacac	atgtttaataa	cagtgggggg	acaccaaagc	gccatgcmaa	tgttaaaaga	8160
tactattaat	gaagaggcgtg	cagaatggga	tagattacat	ccagtccatg	cggggccctat	8220
tgcaccagcg	cagatgagag	aaccaagggg	aagtgcacata	gcagggaacta	ctagtaccct	8280
tcaggaaacaa	atagcatgga	tgacaagtaa	cccaactatt	ccagtgggag	acatctataa	8340
aagatggata	attctggggg	taataaaaa	agtgagaatg	tatagcccg	tcagcatbttt	8400
ggacataaaga	caagggccaa	aggaaccott	tcgagaactat	gtagatcggt	tccttaaaac	8460
tttaagagct	gaacaagcta	cacaagaagt	aaaaaattgg	atgacagaca	ccttggttagt	8520
ccaaaaatg	aaccagatg	gtaagaccat	tttgagagca	ttaggaccag	gggctacatt	8580
agaagaaatg	atgacagcat	gtcaaggggt	gggaggacct	ggccacaaga	caagagtatt	8640
gggtcaggca	atagctcaaa	caaacagctg	aaacataatg	atgcagagaa	gcaattttaa	8700
aggccctaga	agaaatgtta	aatgttttaa	ctgtggcaag	gaagggccaa	tagccagaaa	8760
ttgcagagcc	cctaggaaaa	aaggctgttg	gaaatgtgga	aaagaaggac	accaaatgaa	8820
agactgcact	gagaggcagg	ctaatttttt	agggaataat	tggccttccc	acaaggggag	8880
gccagggaat	ttccttcaga	acagaccaga	gccaaacgac	ccaccagcag	agagcttcag	8940
gttcgaagag	acaaccoccc	ctcogaacaa	ggagccgata	gaaagggaa	ccttaacctc	9000
cctcaaatca	ctcttggcca	gogaccocct	gtctcaataa	gagtttaatt	aagtaacagt	9060
acagcagcaa	ttggcgaagt	gcttaccatg	aactcogcgc	gattggcatg	ccgctttaaa	9120
atbttttttt	tattttttct	ttcttttccg	aatcggtatt	tgtttttaat	atbttcaaaa	9180
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	gggaagagcg	cgcccgccgc	9240
ctgggtacac	ttttgtggc	gttcgcgacg	cgaggctgga	tggccttccc	cattatgatt	9300
cttctcgctt	cggcgccatg	cgggatgccc	gcgttgccag	ccatgtctgc	caggcaggtg	9360
gatgacgacc	atcagggaca	gcttcaagga	tgcctcgccg	ctcttaccag	cctaactctg	9420
atcattggac	cgtctgatgt	caaggcgatt	tatgcgcgct	cggcgagcac	atggaaacgg	9480
ttggcatgga	ttgtaggcgc	cgcctatcac	cttgtctcgc	tcccgcggtt	gggtcgcggt	9540
gcabtgagcc	ggccacccct	gaacctgaatg	gaagccggcg	gcacctcgct	aacgggttca	9600
ccactccaag	aattggagcc	aatcaattct	tggggagaa	tgtgaatgoc	caaaccaacc	9660
cttgccagaa	catatccatc	gcgtccgcga	tctccagcag	cgcgacgccc	cgcactcogg	9720
gcagcgcttg	gtcctggcca	cgggtgcgca	tgatctgtct	cctgtcgttg	aggacccgcg	9780
taggctggcg	gggttgccct	actggttagc	agaaatgaat	acogatacgc	gagcgaaact	9840
gaagcgcactg	ctgcctcaaa	acgtctcgca	cctgagcaac	aaatgaatg	gtcttcgggt	9900
tcctgttttc	gtaaagtctg	gaaacgcgga	agtgcagccc	ctgcaccatt	atgttcogga	9960
ttctgatcgc	aggatgtctg	tggctacccct	gtggaacacc	tacatctgta	tttaacgaagc	10020
gttgccattg	accctgagtg	atbttttctc	ggcccgccgc	catccatacc	gcgaagtgtt	10080
tacctcaca	acgttcagtg	aacccggcat	gttctatc	agtaaccogt	atcgtgagca	10140
tcctctctgc	tttcatcggt	atcattacc	ccatgaacag	aaatccccc	tacacggagg	10200
catcagtgac	caaacaggaa	aaaaccgcc	ttaacatg	ccgctttatc	agaagccaga	10260
cattacagct	cttgaggaaa	ctcaacgagc	tggacggcga	tgaacaggca	gacatctgtg	10320
aatcgcttca	cgaccacgct	gatgagcttt	acccgagctg	ctcgcgcgtt	ttcggtagtg	10380
acggtgaaaa	cctctgacac	atgcagctcc	cggagacggt	cacagcttgt	ctgtagcggg	10440
ttcccgggag	cagacaagcc	cgtcaggcgc	cgtcagcggg	tgttggcggg	tgtcggggcg	10500
cagccatgac	cagctcaagt	agcgatagcg	gagtgatac	tggcttaact	atgcgcgac	10560
agagcagatt	gtacatgagag	tgcaccattg	cgggtgtaaa	taccgcacag	atgcgtgaag	10620
agaaataacc	gcacatgcgc	ctcttcgcgt	tcctcgctca	ctgactcgct	atgcgtggct	10680
gttcggctgc	ggcgagcggg	atcagctcac	tcaaaaggcg	taatacgggt	atccacagaa	10740
tcagggggata	accagggaaa	gaacatgtga	gcaaaaggcc	agcaaaagcg	caggacaggt	10800
aaaaagcccg	cgttgcctgc	gtttttccat	aggctccgcc	ccctgcagca	gcatacaca	10860
aatcgacgct	caagtccagag	gtggcgaaac	ccgacaggac	tataaagata	ccaggcgctt	10920
ccccctggaa	ctctccctgt	gcgctctctc	gtctccagcc	tgcgccttac	cggtacacgt	10980
tcgcctcttc	tcctctcggg	aagcgtggcg	ctcttctcata	gtctacgctg	taggtatctc	11040


```

agttcgggtg aggtcgttgc ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc 11100
gacccgtcgc ccttaccggg taactatcgt cttgagtgca acccggttaag aacagactta 11160
tcgccactcg cagcagccac tggtaacagg attagcagag cgagggtatgt aggcgggtgt 11220
acaggtctct tgaagtggtg gcttaactac ggctacacta gaaggacagt atttggtatc 11280
tcgcgtctgc tgaagccagt taccttcgga aaaagagttg gtatgctctg atccggcaaa 11340
caaacacacg ctggtagcgg tgggtttttt gtttgcaagc agcagattac gcgcagaaaa 11400
aaaggtatcc aagaagatcc ttgatcttt tctacgggtg ctgacgctca gtggaaacgaa 11460
aactcacgtt aagggaattt ggtcatgaac aataaaactg tctgcttaca taacacgtaa 11520
tacaaagggg gttatgagcc atattcaacg ggaacgctct tgctcagagg cgcgattaaa 11580
ttccaacatg gatgctgatt tatatgggta taaatgggct cgcgataatg tcgggcaatc 11640
aggtgcgaca atctatcgat tgtatgggaa gcccgatgag ccagagtgtg ttctgaaaca 11700
tggcaagggt agcgttgcca atgatgttac agatgagatg gtcagactaa actggctgac 11760
ggaatttatg cctcttcoga ccatcaagca ttttatcgtt actcgtgatg atgcatgggt 11820
actcaccact gogatcccg ggaacacagc attccaggtt ttagaagaat atcctgattc 11880
aggtgaaatt attgttgatg cgcctggcag gtctcctgag cgggtgcatt cgattcctgt 11940
ttgtaattgt ccttttaaca gogatcgcgt atttcgtctc gctcaggcgc aatcagcaat 12000
gaataacggg ttggttgatg cagatgattt tgatgacgag cgtaatggct ggctgttgta 12060
acaagctcgg aaagaaatgc ataagctttt gccattctca cgggattcag tgcctactca 12120
tggtgatttc tcaactgata accttatttt tgacgagggg aaattaatag gtgtatttga 12180
tgttgagcga gtccgaatcg cagaccgata ccaggatctt gccatctcat ggaactgcct 12240
cggttgagtt tctcctcat tacagaaaag gctttttcaa aaatatggta tggataatcc 12300
ttgatgaatt aaatgcagat ttcatttgat gctcagatg ttttctaaag aattcctatg 12360
tttgacagct tatcatcgat aagctttaat gcggtagttt atcacagtta aattgctaac 12420
gcagtcagggc accggtgatg aaatctaaca atgcctcat cgtcatcttc ggcacccgtca 12480
ccctgtagtc tgtctagagg atccctaata cgaactcaata tag 12523

```

<210> 2

<211> 7479

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<221> CDS

<222> (1)...(7479)

<400> 2

```

atg gag aaa gtt cac gtt gac atc gag gaa gac agc cca ttc ctc aga      48
Met Glu Lys Val His Val Asp Ile Glu Glu Asp Ser Pro Phe Leu Arg
1          5          10          15

```

```

gct ttg cag cgg agc ttc cgg cag ttt gag gta gaa gcc aag cag gtc      96
Ala Leu Gln Arg Ser Phe Pro Gln Phe Glu Val Glu Ala Lys Gln Val
20          25          30

```

```

act gat aat gac cat gct aat gcc aga gcg ttt tcg cat ctg gct tca      144
Thr Asp Asn Asp His Ala Asn Ala Arg Ala Phe Ser His Leu Ala Ser
35          40          45

```

```

aaa ctg atc gaa acg gag gtg gac cca tcc gac acg atc ctt gac att      192
Lys Leu Ile Glu Thr Glu Val Asp Pro Ser Asp Thr Ile Leu Asp Ile
50          55          60

```

gga agt gcg ccc gcc cgc aga atg tat tct aag cac aag tat cat tgt Gly Ser Ala Pro Ala Arg Arg Met Tyr Ser Lys His Lys Tyr His Cys 65 70 75 80	240
atc tgt ccg atg aga tgt gcg gaa gat ccg gac aga ttg tat aag tat Ile Cys Pro Met Arg Cys Ala Glu Asp Pro Asp Arg Leu Tyr Lys Tyr 85 90 95	288
gca act aag ctg aag aaa aac tgt aag gaa ata act gat aag gaa ttg Ala Thr Lys Leu Lys Lys Asn Cys Lys Glu Ile Thr Asp Lys Glu Leu 100 105 110	336
gac aag aaa atg aag gag ctc gcc gcc gtc atg agc gac cct gac ctg Asp Lys Lys Met Lys Glu Leu Ala Ala Val Met Ser Asp Pro Asp Leu 115 120 125	384
gaa act gag act atg tgc ctc cac gac gac gag tgc tgt cgc tac gaa Glu Thr Glu Thr Met Cys Lys His Asp Asp Glu Ser Cys Arg Tyr Glu 130 135 140	432
ggg caa gtc gct gtt tac cag gat gta tac gcg gtt gac gga ccg aca Gly Gln Val Ala Val Tyr Gln Asp Val Tyr Ala Val Asp Gly Pro Thr 145 150 155 160	480
agt ctc tat cac caa gcc aat aag gga gtt aga gtc gcc tac tgg ata Ser Leu Tyr His Gln Ala Asn Lys Gly Val Arg Val Ala Tyr Trp Ile 165 170 175	528
ggc ttt gac acc acc cct ttt atg ttt aag aac ttg gct gga gca tat Gly Phe Asp Thr Thr Pro Phe Met Phe Lys Asn Leu Ala Gly Ala Tyr 180 185 190	576
cca tca tac tct acc aac tgg gcc gac gaa acc gtg tta acg gct cgt Pro Ser Tyr Ser Thr Asn Trp Ala Asp Glu Thr Val Leu Thr Ala Arg 195 200 205	624
aac ata ggc cta tgc agc tct gac gtt atg gag cgg tca cgt aga ggg Asn Ile Gly Leu Cys Ser Ser Asp Val Met Glu Arg Ser Arg Arg Gly 210 215 220	672
atg tcc att ctt aga aag aag tat ttg aaa cca tcc aac aat gtt cta Met Ser Ile Leu Arg Lys Lys Tyr Leu Lys Pro Ser Asn Asn Val Leu 225 230 235 240	720
ttc tct gtt ggc tgc acc atc tac cac gag aag agg gac tta ctg agg Phe Ser Val Gly Ser Thr Ile Tyr His Glu Lys Arg Asp Leu Leu Arg 245 250 255	768
agc tgg cac ctg ccg tct gta ttt cac tta cgt gcc aag caa aat tac Ser Trp His Leu Pro Ser Val Phe His Leu Arg Gly Lys Gln Asn Tyr 260 265 270	816
aca tgt cgg tgt gag act ata gtt agt tgc gac ggg tac gtc gtt aaa Thr Cys Arg Cys Glu Thr Ile Val Ser Cys Asp Gly Tyr Val Val Lys 275 280 285	864

aga ata gct atc agt cca ggc ctg tat ggg aag cct tca ggc tat gct Arg Ile Ala Ile Ser Pro Gly Leu Tyr Gly Lys Pro Ser Gly Tyr Ala 290 295 300	912
gct acg atg cac cgc gag gga ttc ttg tgc tgc aaa gtg aca gac aca Ala Thr Met His Arg Glu Gly Phe Leu Cys Cys Lys Val Thr Asp Thr 305 310 315 320	960
tta aac ggg gag agg gtc tct ttt ccc gtg tgc acg tat gtg cca gct Leu Asn Gly Glu Arg Val Ser Phe Pro Val Cys Thr Tyr Val Pro Ala 325 330 335	1008
aca ttg tgt gac caa atg act ggc ata ctg gca aca gat gtc agt gcg Thr Leu Cys Asp Gln Met Thr Gly Ile Leu Ala Thr Asp Val Ser Ala 340 345 350	1056
gac gac gcg caa aaa ctg ctg gtt ggg ctc aac cag cgt ata gtc gtc Asp Asp Ala Gln Lys Leu Leu Val Gly Leu Asn Gln Arg Ile Val Val 355 360 365	1104
aac ggt cgc acc cag aga aac acc aat acc atg aaa aat tac ctt ttg Asn Gly Arg Thr Gln Arg Asn Thr Asn Thr Met Lys Asn Tyr Leu Leu 370 375 380	1152
ccc gta gtg gcc cag gca ttt gct agg tgg gca aag gaa tat aag gaa Pro Val Val Ala Gln Ala Phe Ala Arg Trp Ala Lys Glu Tyr Lys Glu 385 390 395 400	1200
gat caa gaa gat gaa agg cca cta gga cta cga gat aga cag tta gtc Asp Gln Glu Asp Glu Arg Pro Leu Gly Leu Arg Asp Arg Gln Leu Val 405 410 415	1248
atg ggg tgt tgt tgg gct ttt aga agg cac aag ata aca tct att tat Met Gly Cys Trp Ala Phe Arg Arg His Lys Ile Thr Ser Ile Tyr 420 425 430	1296
aag cgc ccg gat acc caa acc atc atc aaa gtg aac agc gat ttc cac Lys Arg Pro Asp Thr Gln Thr Ile Ile Lys Val Asn Ser Asp Phe His 435 440 445	1344
tca ttc gtg ctg ccc agg ata ggc agt aac aca ttg gag atc ggg ctg Ser Phe Val Leu Pro Arg Ile Gly Ser Asn Thr Leu Glu Ile Gly Leu 450 455 460	1392
aga aca aga atc agg aaa atg tta gag gag cac aag gag ccg tca cct Arg Thr Arg Ile Arg Lys Met Leu Glu Glu His Lys Glu Pro Ser Pro 465 470 475 480	1440
ctc att acc gcc gag gac gta caa gaa gct aag tgc gca gcc gat gag Leu Ile Thr Ala Glu Asp Val Gln Glu Ala Lys Cys Ala Ala Asp Glu 485 490 495	1488
gct aag gag gtg cgt gaa gcc gag gag ttg cgc gca gct cta cca cct Ala Lys Glu Val Arg Glu Ala Glu Glu Leu Arg Ala Ala Leu Pro Pro 500 505 510	1536

ttg gca gct gat gtt gag gag ccc act ctg gaa gcc gat gtc gac ttg Leu Ala Ala Asp Val Glu Glu Pro Thr Leu Glu Ala Asp Val Asp Leu 515 520 525	1584
atg tta caa gag gct ggg gcc ggc tca gtg gag aca cct cgt ggc ttg Met Leu Gln Glu Ala Gly Ala Gly Ser Val Glu Thr Pro Arg Gly Leu 530 535 540	1632
ata aag gtt acc agc tac gct ggc gag gac aag atc ggc tct tac gct Ile Lys Val Thr Ser Tyr Ala Gly Glu Asp Lys Ile Gly Ser Tyr Ala 545 550 555 560	1680
gtg ctt tct ccg cag gct gta ctc aag agt gaa aaa tta tct tgc atc Val Leu Ser Pro Gln Ala Val Leu Lys Ser Glu Lys Leu Ser Cys Ile 565 570 575	1728
cac cct ctc gct gaa caa gtc ata gtg ata aca cac tct ggc cga aaa His Pro Leu Ala Glu Gln Val Ile Val Ile Thr His Ser Gly Arg Lys 580 585 590	1776
ggg cgt tat gcc gtg gaa cca tac cat ggt aaa gta gtg gtg cca gag Gly Arg Tyr Ala Val Glu Pro Tyr His Gly Lys Val Val Val Pro Glu 595 600 605	1824
gga cat gca ata ccc gtc cag gac ttt caa gct ctg agt gaa agt gcc Gly His Ala Ile Pro Val Gln Asp Phe Gln Ala Leu Ser Glu Ser Ala 610 615 620	1872
acc att gtg tac aac gaa cgt gag ttc gta aac agg tac ctg cac cat Thr Ile Val Tyr Asn Glu Arg Glu Phe Val Asn Arg Tyr Leu His His 625 630 635 640	1920
att gcc aca cat gga gga gcg ctg aac act gat gaa gaa tat tac aaa Ile Ala Thr His Gly Gly Ala Leu Asn Thr Asp Glu Glu Tyr Tyr Lys 645 650 655	1968
act gtc aag ccc agc gag cac gac ggc gaa tac ctg tac gac atc gac Thr Val Lys Pro Ser Glu His Asp Gly Glu Tyr Leu Tyr Asp Ile Asp 660 665 670	2016
agg aaa cag tgc gtc aag aaa gaa cta gtc act ggg cta ggg ctc aca Arg Lys Gln Cys Val Lys Lys Glu Leu Val Thr Gly Leu Gly Leu Thr 675 680 685	2064
ggc gag ctg gtg gat cct ccc ttc cat gaa ttc gcc tac gag agt ctg Gly Glu Leu Val Asp Pro Pro Phe His Glu Phe Ala Tyr Glu Ser Leu 690 695 700	2112
aga aca cga cca gcc gct cct tac caa gta cca acc ata ggg gtg tat Arg Thr Arg Pro Ala Ala Pro Tyr Gln Val Pro Thr Ile Gly Val Tyr 705 710 715 720	2160
ggc gtg cca gga tca ggc aag tct ggc atc att aaa agc gca gtc acc Gly Val Pro Gly Ser Gly Lys Ser Gly Ile Ile Lys Ser Ala Val Thr 725 730 735	2208

aaa aaa gat cta gtg gtg agc gcc aag aaa gaa aac tgt gca gaa att Lys Lys Asp Leu Val Val Ser Ala Lys Lys Lys Glu Asn Cys Ala Glu Ile 740 745 750	2256
ata agg gac gtc aag aaa atg aaa ggg ctg gac gtc aat gcc aga act Ile Arg Asp Val Lys Lys Met Lys Gly Leu Asp Val Asn Ala Arg Thr 755 760 765	2304
gtg gac tca gtg ctc ttg aat gga tgc aaa cac ccc gta gag acc ctg Val Asp Ser Val Leu Leu Asn Gly Cys Lys His Pro Val Glu Thr Leu 770 775 780	2352
tat att gac gaa gct ttt gct tgt cat gca ggt act ctc aga gcg ctc Tyr Ile Asp Glu Ala Phe Ala Cys His Ala Gly Thr Leu Arg Ala Leu 785 790 795 800	2400
ata gcc att ata aga cct aaa aag gca gtg ctc tgc ggg gat ccc aaa Ile Ala Ile Ile Arg Pro Lys Lys Ala Val Leu Cys Gly Asp Pro Lys 805 810 815	2448
cag tgc ggt ttt ttt aac atg atg tgc ctg aaa gtg cat ttt aac cac Gln Cys Gly Phe Phe Asn Met Met Cys Leu Lys Val His Phe Asn His 820 825 830	2496
gag att tgc aca caa gtc ttc cac aaa agc atc tct cgc cgt tgc act Glu Ile Cys Thr Gln Val Phe His Lys Ser Ile Ser Arg Arg Cys Thr 835 840 845	2544
aaa tct gtg act tcg gtc gtc tca acc ttg ttt tac gac aaa aaa atg Lys Ser Val Thr Ser Val Val Ser Thr Leu Phe Tyr Asp Lys Lys Met 850 855 860	2592
aga acg acg aat ccg aaa gag act aag att gtg att gac act acc ggc Arg Thr Thr Asn Pro Lys Glu Thr Lys Ile Val Ile Asp Thr Thr Gly 865 870 875 880	2640
agt acc aaa cct aag cag gac gat ctc att ctc act tgt ttc aga ggg Ser Thr Lys Pro Lys Gln Asp Asp Leu Ile Leu Thr Cys Phe Arg Gly 885 890 895	2688
tgg gtg aag cag ttg caa ata gat tac aaa ggc aac gaa ata atg acg Trp Val Lys Gln Leu Gln Ile Asp Tyr Lys Gly Asn Glu Ile Met Thr 900 905 910	2736
gca gct gcc tct caa ggg ctg acc cgt aaa ggt gtg tat gcc gtt cgg Ala Ala Ala Ser Gln Gly Leu Thr Arg Lys Gly Val Tyr Ala Val Arg 915 920 925	2784
tac aag gtg aat gaa aat cct ctg tac gca ccc acc tca gaa cat gtg Tyr Lys Val Asn Glu Asn Pro Leu Tyr Ala Pro Thr Ser Glu His Val 930 935 940	2832
aac gtc cta ctg acc cgc acg gag gac cgc atc gtg tgg aaa aca cta Asn Val Leu Leu Thr Arg Thr Glu Asp Arg Ile Val Trp Lys Thr Leu 945 950 955 960	2880

gcc ggc gac cca tgg ata aaa aca ctg act gcc aag tac cct ggg aat Ala Gly Asp Pro Trp Ile Lys Thr Leu Thr Ala Lys Tyr Pro Gly Asn 965 970 975	2928
ttc act gcc acg ata gag gag tgg caa gca gag cat gat gcc atc atg Phe Thr Ala Thr Ile Glu Glu Trp Gln Ala Glu His Asp Ala Ile Met 980 985 990	2976
agg cac atc ttg gag aga ccg gac cct acc gac gtc ttc cag aat aag Arg His Ile Leu Glu Arg Pro Asp Pro Thr Asp Val Phe Gln Asn Lys 995 1000 1005	3024
gca aac gtg tgt tgg gcc aag gct tta gtg ccg gtg ctg aag acc gct Ala Asn Val Cys Trp Ala Lys Ala Leu Val Pro Val Leu Lys Thr Ala 1010 1015 1020	3072
ggc ata gac atg acc act gaa caa tgg aac act gtg gat tat ttt gaa Gly Ile Asp Met Thr Thr Glu Gln Trp Asn Thr Val Asp Tyr Phe Glu 1025 1030 1035 1040	3120
acg gac aaa gct cac tca gca gag ata gta ttg aac caa cta tgc gtg Thr Asp Lys Ala His Ser Ala Glu Ile Val Leu Asn Gln Leu Cys Val 1045 1050 1055	3168
agg ttc ttt gga ctc gat ctg gac tcc ggt cta ttt tct gca ccc act Arg Phe Phe Gly Leu Asp Leu Asp Ser Gly Leu Phe Ser Ala Pro Thr 1060 1065 1070	3216
gtt ccg tta tcc att agg aat aat cac tgg gat aac tcc ccg tog cct Val Pro Leu Ser Ile Arg Asn Asn His Trp Asp Asn Ser Pro Ser Pro 1075 1080 1085	3264
aac atg tac ggg ctg aat aaa gaa gtg gtc cgt cag ctc tct cgc agg Asn Met Tyr Gly Leu Asn Lys Glu Val Val Arg Gln Leu Ser Arg Arg 1090 1095 1100	3312
tac cca caa ctg cct cgg gca gtt gcc act gga aga gtc tat gac atg Tyr Pro Gln Leu Pro Arg Ala Val Ala Thr Gly Arg Val Tyr Asp Met 1105 1110 1115 1120	3360
aac act ggt aca ctg cgc aat tat gat ccg cgc ata aac cta gta cct Asn Thr Gly Thr Leu Arg Asn Tyr Asp Pro Arg Ile Asn Leu Val Pro 1125 1130 1135	3408
gta aac aga aga ctg cct cat gct tta gtc ctc cac cat aat gaa cac Val Asn Arg Arg Leu Pro His Ala Leu Val Leu His His Asn Glu His 1140 1145 1150	3456
cca cag agt gac ttt tct tca ttc gtc agc aaa ttg aag ggc aga act Pro Gln Ser Asp Phe Ser Ser Phe Val Ser Lys Leu Lys Gly Arg Thr 1155 1160 1165	3504
gtc ctg gtg gtc ggg gaa aag ttg tcc gtc cca ggc aaa atg gtt gac Val Leu Val Val Gly Glu Lys Leu Ser Val Pro Gly Lys Met Val Asp 1170 1175 1180	3552

tggttg tca gac cgg cct gag gct acc ttc aga gct cgg ctg gat tta	3600
Trp Leu Ser Asp Arg Pro Glu Ala Thr Phe Arg Ala Arg Leu Asp Leu	
1185 1190 1195 1200	
ggc atc cca ggt gat gtg ccc aaa tat gac ata ata ttt gtt aat gtg	3648
Gly Ile Pro Gly Asp Val Pro Lys Tyr Asp Ile Ile Phe Val Asn Val	
1205 1210 1215	
agg acc cca tat aaa tac cat cac tat cag cag tgt gaa gac cat gcc	3696
Arg Thr Pro Tyr Lys Tyr His His Tyr Gln Gln Cys Glu Asp His Ala	
1220 1225 1230	
att aag ctt agc atg ttg acc aag aaa gct tgt ctg cat ctg aat ccc	3744
Ile Lys Leu Ser Met Leu Thr Lys Lys Ala Cys Leu His Leu Asn Pro	
1235 1240 1245	
ggc gga acc tgt gtc agc ata ggt tat ggt tac gct gac agg gcc agc	3792
Gly Gly Thr Cys Val Ser Ile Gly Tyr Gly Tyr Ala Asp Arg Ala Ser	
1250 1255 1260	
gaa agc atc att ggt gct ata gcg cgg cag ttc aag ttt tcc cgg gta	3840
Glu Ser Ile Ile Gly Ala Ile Ala Arg Gln Phe Lys Phe Ser Arg Val	
1265 1270 1275 1280	
tgc aaa ccg aaa tcc tca ctt gaa gag acg gaa gtt ctg ttt gta ttc	3888
Cys Lys Pro Lys Ser Ser Leu Glu Glu Thr Glu Val Leu Phe Val Phe	
1285 1290 1295	
att ggg tac gat cgc aag gcc cgt acg cac aat cct tac aag ctt tca	3936
Ile Gly Tyr Asp Arg Lys Ala Arg Thr His Asn Pro Tyr Lys Leu Ser	
1300 1305 1310	
tca acc ttg acc aac att tat aca ggt tcc aga ctg cac gaa gcc gga	3984
Ser Thr Leu Thr Asn Ile Tyr Thr Gly Ser Arg Leu His Glu Ala Gly	
1315 1320 1325	
tgt gca ccc tca tat cat gtg gtg cga ggg gat att gcc acg gcc acc	4032
Cys Ala Pro Ser Tyr His Val Val Arg Gly Asp Ile Ala Thr Ala Thr	
1330 1335 1340	
gaa gga gtg att ata aat gct gct aac agc aaa gga caa cct gcc gga	4080
Glu Gly Val Ile Ile Asn Ala Ala Asn Ser Lys Gly Gln Pro Gly Gly	
1345 1350 1355 1360	
ggg gtg tgc gga gcg ctg tat aag aag ttc cgg gaa agc ttc gat tta	4128
Gly Val Cys Gly Ala Leu Tyr Lys Lys Phe Pro Glu Ser Phe Asp Leu	
1365 1370 1375	
cag ccg atc gaa gta gga aaa gcg cga ctg gtc aaa ggt gca gct aaa	4176
Gln Pro Ile Glu Val Gly Lys Ala Arg Leu Val Lys Gly Ala Ala Lys	
1380 1385 1390	
cat atc att cat gcc gta gga cca aac ttc aac aaa gtt tgg gag gtt	4224
His Ile Ile His Ala Val Gly Pro Asn Phe Asn Lys Val Ser Glu Val	
1395 1400 1405	

gaa ggt gac aaa cag ttg gca gag gct tat gag tcc atc gct aag att Glu Gly Asp Lys Gln Leu Ala Glu Ala Tyr Glu Ser Ile Ala Lys Ile 1410 1415 1420	4272
gtc aac gat aac aat tac aag tca gta gcg att cca ctg ttg tcc acc Val Asn Asp Asn Asn Tyr Lys Ser Val Ala Ile Pro Leu Leu Ser Thr 1425 1430 1435 1440	4320
ggc atc ttt tcc ggg aac aaa gat cga cta acc caa tca ttg aac cat Gly Ile Phe Ser Gly Asn Lys Asp Arg Leu Thr Gln Ser Leu Asn His 1445 1450 1455	4368
ttg ctg aca gct tta gac acc act gat gca gat gta gcc ata tac tgc Leu Leu Thr Ala Leu Asp Thr Thr Asp Ala Asp Val Ala Ile Tyr Cys 1460 1465 1470	4416
agg gac aag aaa tgg gaa atg act ctc aag gaa gca gtg gct agg aga Arg Asp Lys Lys Trp Glu Met Thr Leu Lys Glu Ala Val Ala Arg Arg 1475 1480 1485	4464
gaa gca gtg gag gag ata tgc ata tcc gac gac tct tca gtg aca gaa Glu Ala Val Glu Glu Ile Cys Ile Ser Asp Asp Ser Ser Val Thr Glu 1490 1495 1500	4512
cct gat gca gag ctg gtg agg gtg cat cgg aag agt tct ttg gct gga Pro Asp Ala Glu Leu Val Arg Val His Pro Lys Ser Ser Leu Ala Gly 1505 1510 1515 1520	4560
agg aag ggc tac agc aca agc gat ggc aaa act ttc tca tat ttg gaa Arg Lys Gly Tyr Ser Thr Ser Asp Gly Lys Thr Phe Ser Tyr Leu Glu 1525 1530 1535	4608
ggg acc aag ttt cac cag gcg gcc aag gat ata gca gaa att aat gcc Gly Thr Lys Phe His Gln Ala Ala Lys Asp Ile Ala Glu Ile Asn Ala 1540 1545 1550	4656
atg tgg ccc gtt gca acg gag gcc aat gag cag gta tgc atg tat atc Met Trp Pro Val Ala Thr Glu Ala Asn Glu Gln Val Cys Met Tyr Ile 1555 1560 1565	4704
ctc gga gaa agc atg agc agt att agg tgg aaa tgc ccc gtc gaa gag Leu Gly Glu Ser Met Ser Ser Ile Arg Ser Lys Cys Pro Val Glu Glu 1570 1575 1580	4752
tgg gaa gcc tcc aca cca cct agc acg ctg cct tgc ttg tgc atc cat Ser Glu Ala Ser Thr Pro Pro Ser Thr Leu Pro Cys Leu Cys Ile His 1585 1590 1595 1600	4800
gcc atg act cca gaa aga gta cag cgc cta aaa gcc tca cgt cca gaa Ala Met Thr Pro Glu Arg Val Gln Arg Leu Lys Ala Ser Arg Pro Glu 1605 1610 1615	4848
caa att act gtg tgc tca tcc ttt cca ttg cgg aag tat aga atc act Gln Ile Thr Val Cys Ser Ser Phe Pro Leu Pro Lys Tyr Arg Ile Thr 1620 1625 1630	4896

ggt gtg cag aag atc caa tgc tcc cag cct ata ttg ttc tca cag aaa Gly Val Gln Lys Ile Gln Cys Ser Gln Pro Ile Leu Phe Ser Pro Lys 1635 1640 1645	4944
gtg cct gcg tat att cat cca agg aag tat ctc gtg gaa aca cca cag Val Pro Ala Tyr Ile His Pro Arg Lys Tyr Leu Val Glu Thr Pro Pro 1650 1655 1660	4992
gta gac gag act ccg gag cca tgc gca gag aac caa tcc aca gag ggg Val Asp Glu Thr Pro Glu Pro Ser Ala Glu Asn Gln Ser Thr Glu Gly 1665 1670 1675 1680	5040
aca cct gaa caa cca cca ctt ata acc gag gat gag acc agg act aga Thr Pro Glu Gln Pro Pro Leu Ile Thr Glu Asp Glu Thr Arg Thr Arg 1685 1690 1695	5088
acg cct gag ccg atc atc atc gaa gag gaa gaa gag gat agc ata agt Thr Pro Glu Pro Ile Ile Ile Glu Glu Glu Glu Asp Ser Ile Ser 1700 1705 1710	5136
ttg ctg tca gat ggc ccg acc cac cag gtg ctg caa gtc gag gca gac Leu Leu Ser Asp Gly Pro Thr His Gln Val Leu Gln Val Glu Ala Asp 1715 1720 1725	5184
att cac ggg ccg ccc tct gta tct agc tca tcc tgg tcc att cct cat Ile His Gly Pro Pro Ser Val Ser Ser Ser Ser Trp Ser Ile Pro His 1730 1735 1740	5232
gca tcc gac ttt gat gtg gac agt tta tcc ata ctt gac acc ctg gag Ala Ser Asp Phe Asp Val Asp Ser Leu Ser Ile Leu Asp Thr Leu Glu 1745 1750 1755 1760	5280
gga gct agc gtg acc agc ggg gca acg tca gcc gag act aac tct tac Gly Ala Ser Val Thr Ser Gly Ala Thr Ser Ala Glu Thr Asn Ser Tyr 1765 1770 1775	5328
ttc gca aag agt atg gag ttt ctg gcg cga ccg gtg cct gcg cct cga Phe Ala Lys Ser Met Glu Phe Leu Ala Arg Pro Val Pro Ala Pro Arg 1780 1785 1790	5376
aca gta ttc agg aac cct cca cat ccc gct ccg cgc aca aga aca ccg Thr Val Phe Arg Asn Pro Pro His Pro Ala Pro Arg Thr Arg Thr Pro 1795 1800 1805	5424
tca ctt gca ccc agc agg gcc tgc tgc aga acc agc cta gtt tcc acc Ser Leu Ala Pro Ser Arg Ala Cys Ser Arg Thr Ser Leu Val Ser Thr 1810 1815 1820	5472
ccg cca ggc gtg aat agg gtg atc act aga gag gag ctc gag gcg ctt Pro Pro Gly Val Asn Arg Val Ile Thr Arg Glu Glu Leu Glu Ala Leu 1825 1830 1835 1840	5520
acc ccg tca cgc act cct agc agg tgc gtc tgc aga acc agc ctg gtc Thr Pro Ser Arg Thr Pro Ser Arg Ser Val Ser Arg Thr Ser Leu Val 1845 1850 1855	5568

tcc aac ccg cca ggc gta aat agg	gtg att aca aga gag gag ttt gag	5616
Ser Asn Pro Gly Val Asn Arg	Val Ile Thr Arg Glu Glu Phe Glu	
1860	1865 1870	
gcg ttc gta gca caa caa caa tga	cgg ttt gat gcg ggt gca tac atc	5664
Ala Phe Val Ala Gln Gln Gln *	Arg Phe Asp Ala Gly Ala Tyr Ile	
1875	1880 1885	
ttt tcc tcc gac acc ggt caa ggg cat tta caa caa aaa tca gta agg		5712
Phe Ser Ser Asp Thr Gly Gln Gly His Leu Gln Gln Lys Ser Val Arg		
1890	1895 1900	
caa acg gtg cta tcc gaa gtg gtg ttg gag agg acc gaa ttg gag att		5760
Gln Thr Val Leu Ser Glu Val Val Leu Glu Arg Thr Glu Leu Glu Ile		
1905	1910 1915	
tcg tat gcc ccg cgc ctc gac caa gaa aaa gaa gaa tta cta cgc aag		5808
Ser Tyr Ala Pro Arg Leu Asp Gln Glu Lys Glu Glu Leu Leu Arg Lys		
1920	1925 1930 1935	
aaa tta cag tta aat ccc aca cct gct aac aga agc aga tac cag tcc		5856
Lys Leu Gln Leu Asn Pro Thr Pro Ala Asn Arg Ser Arg Tyr Gln Ser		
1940	1945 1950	
agg aag gtg gag aac atg aaa gcc ata aca gct aga cgt att ctg caa		5904
Arg Lys Val Glu Asn Met Lys Ala Ile Thr Ala Arg Arg Ile Leu Gln		
1955	1960 1965	
ggc cta ggg cat tat ttg aag gca gaa gga aaa gtg gag tgc tac cga		5952
Gly Leu Gly His Tyr Leu Lys Ala Glu Gly Lys Val Glu Cys Tyr Arg		
1970	1975 1980	
acc ctg cat cct gtt cct ttg tat tca tct agt gtg aac cgt gcc ttt		6000
Thr Leu His Pro Val Pro Leu Tyr Ser Ser Ser Val Asn Arg Ala Phe		
1985	1990 1995	
tca agc ccc aag gtc gca gtg gaa gcc tgt aac gcc atg ttg aaa gag		6048
Ser Ser Pro Lys Val Ala Val Glu Ala Cys Asn Ala Met Leu Lys Glu		
2000	2005 2010 2015	
aac ttt ccg act gtg gct tct tac tgt att att cca gag tac gat gcc		6096
Asn Phe Pro Thr Val Ala Ser Tyr Cys Ile Ile Pro Glu Tyr Asp Ala		
2020	2025 2030	
tat ttg gac atg gtt gac gga gct tca tgc tgc tta gac act gcc agt		6144
Tyr Leu Asp Met Val Asp Gly Ala Ser Cys Cys Leu Asp Thr Ala Ser		
2035	2040 2045	
ttt tgc cct gca aag ctg cgc agc ttt cca aag aaa cac tcc tat ttg		6192
Phe Cys Pro Ala Lys Leu Arg Ser Phe Pro Lys Lys His Ser Tyr Leu		
2050	2055 2060	
gaa ccc aca ata cga tog gca gtg cct tca gcg atc cag aac acg ctc		6240
Glu Pro Thr Ile Arg Ser Ala Val Pro Ser Ala Ile Gln Asn Thr Leu		
2065	2070 2075	

cag aac gtc ctg gca gct gcc aca aaa aga aat tgc aat gtc acg caa Gln Asn Val Leu Ala Ala Thr Lys Arg Asn Cys Asn Val Thr Gln 2080 2085 2090 2095	6288
atg aga gaa ttg ccc gta ttg gat tgc gcg gcc ttt aat gtg gaa tgc Met Arg Glu Leu Pro Val Leu Asp Ser Ala Ala Phe Asn Val Glu Cys 2100 2105 2110	6336
ttc aag aaa tat gcg tgt aat aat gaa tat tgg gaa acg ttt aaa gaa Phe Lys Lys Tyr Ala Cys Asn Asn Glu Tyr Trp Glu Thr Phe Lys Glu 2115 2120 2125	6384
aac ccc atc agg ctt act gaa gaa aac gtg gta aat tac att acc aaa Asn Pro Ile Arg Leu Thr Glu Glu Asn Val Val Asn Tyr Ile Thr Lys 2130 2135 2140	6432
tta aaa gga cca aaa gct gct gct ctt ttt gcg aag aca cat aat ttg Leu Lys Gly Pro Lys Ala Ala Ala Leu Phe Ala Lys Thr His Asn Leu 2145 2150 2155	6480
aat atg ttg cag gac ata cca atg gac agg ttt gta atg gac tta aag Asn Met Leu Gln Asp Ile Pro Met Asp Arg Phe Val Met Asp Leu Lys 2160 2165 2170 2175	6528
aga gac gtg aaa gtg act cca gga aca aaa cat act gaa gaa cgg ccc Arg Asp Val Lys Val Thr Pro Gly Thr Lys His Thr Glu Glu Arg Pro 2180 2185 2190	6576
aag gta cag gtg atc cag gct gcc gat cgg cta gca aca gcg tat ctg Lys Val Gln Val Ile Gln Ala Ala Asp Pro Leu Ala Thr Ala Tyr Leu 2195 2200 2205	6624
tgc gga atc cac cga gag ctg gtt agg aga tta aat gcg gtc ctg ctt Cys Gly Ile His Arg Glu Leu Val Arg Arg Leu Asn Ala Val Leu Leu 2210 2215 2220	6672
ccg aac att cat aca ctg ttt gat atg tgc gct gaa gac ttt gac gct Pro Asn Ile His Thr Leu Phe Asp Met Ser Ala Glu Asp Phe Asp Ala 2225 2230 2235	6720
att ata gcc gag cac ttc cag cct ggg gat tgt gtt ctg gaa act gac Ile Ile Ala Glu His Phe Gln Pro Gly Asp Cys Val Leu Glu Thr Asp 2240 2245 2250 2255	6768
atc gcg tgc ttt gat aaa agt gag gac gac gcc atg gct ctg acc gcg Ile Ala Ser Phe Asp Lys Ser Glu Asp Asp Ala Met Ala Leu Thr Ala 2260 2265 2270	6816
tta atg att ctg gaa gac tta ggt gtg gac gca gag ctg ttg acg ctg Leu Met Ile Leu Glu Asp Leu Gly Val Asp Ala Glu Leu Thr Leu 2275 2280 2285	6864
att gag gcg gct ttc ggc gaa att tca tca ata cat ttg ccc act aaa Ile Glu Ala Ala Phe Gly Glu Ile Ser Ser Ile His Leu Pro Thr Lys 2290 2295 2300	6912

act aaa ttt aaa ttc gga gcc atg atg aaa tct gga atg ttc ctc aca Thr Lys Phe Lys Phe Gly Ala Met Met Lys Ser Gly Met Phe Leu Thr 2305 2310 2315	6960
ctg ttt gtg aac aca gtc att aac att gta atc gca agc aga gtg ttg Leu Phe Val Asn Thr Val Ile Asn Ile Val Ile Ala Ser Arg Val Leu 2320 2325 2330 2335	7008
aga gaa cgg cta acc gga tca cca tgt gca gca ttc att gga gat gac Arg Glu Arg Leu Thr Gly Ser Pro Cys Ala Ala Phe Ile Gly Asp Asp 2340 2345 2350	7056
aat atc gtg aaa gga gtc aaa tcg gac aaa tta atg gca gac agg tgc Asn Ile Val Lys Gly Val Lys Ser Asp Lys Leu Met Ala Asp Arg Cys 2355 2360 2365	7104
gcc acc tgg ttg aat atg gaa gtc aag att ata gat gct gtg gtg ggc Ala Thr Trp Leu Asn Met Glu Val Lys Ile Ile Asp Ala Val Val Gly 2370 2375 2380	7152
gag aaa gcg ccc tat ttc tgt gga ggg ttt att ttg tgt gac tcc gtg Glu Lys Ala Pro Tyr Phe Cys Gly Gly Phe Ile Leu Cys Asp Ser Val 2385 2390 2395	7200
acc gcc aca gcg tgc cgt gtg gca gac ccc cta aaa agg ctg ttt aag Thr Gly Thr Ala Cys Arg Val Ala Asp Pro Leu Lys Arg Leu Phe Lys 2400 2405 2410 2415	7248
ctt gcc aaa cct ctg gca gca gac gat gaa cat gat gat gac agg aga Leu Gly Lys Pro Leu Ala Ala Asp Asp Glu His Asp Asp Arg Arg 2420 2425 2430	7296
agg gca ttg cat gaa gag tca aca cgc tgg aac cga gtg ggt att ctt Arg Ala Leu His Glu Glu Ser Thr Arg Trp Asn Arg Val Gly Ile Leu 2435 2440 2445	7344
tca gag ctg tgc aag gca gta gaa tca agg tat gaa acc gta gga act Ser Glu Leu Cys Lys Ala Val Glu Ser Arg Tyr Glu Thr Val Gly Thr 2450 2455 2460	7392
tcc atc ata gtt atg gcc atg act act cta gct agc agt gtt aaa tca Ser Ile Ile Val Met Ala Met Thr Thr Leu Ala Ser Ser Val Lys Ser 2465 2470 2475	7440
ttc agc tac ctg aga ggg gcc cct ata act ctc tac ggc Phe Ser Tyr Leu Arg Gly Ala Pro Ile Thr Leu Tyr Gly 2480 2485 2490	7479
<210> 3	
<211> 2492	
<212> PRT	
<213> Artificial Sequence	

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 3

```

Met Glu Lys Val His Val Asp Ile Glu Glu Asp Ser Pro Phe Leu Arg
 1           5           10           15
Ala Leu Gln Arg Ser Phe Pro Gln Phe Glu Val Glu Ala Lys Gln Val
 20           25           30
Thr Asp Asn Asp His Ala Asn Ala Arg Ala Phe Ser His Leu Ala Ser
 35           40           45
Lys Leu Ile Glu Thr Glu Val Asp Pro Ser Asp Thr Ile Leu Asp Ile
 50           55           60
Gly Ser Ala Pro Ala Arg Arg Met Tyr Ser Lys His Lys Tyr His Cys
 65           70           75           80
Ile Cys Pro Met Arg Cys Ala Glu Asp Pro Asp Arg Leu Tyr Lys Tyr
 85           90           95
Ala Thr Lys Leu Lys Lys Asn Cys Lys Glu Ile Thr Asp Lys Glu Leu
100          105          110
Asp Lys Lys Met Lys Glu Leu Ala Ala Val Met Ser Asp Pro Asp Leu
115          120          125
Glu Thr Glu Thr Met Cys Leu His Asp Asp Glu Ser Cys Arg Tyr Glu
130          135          140
Gly Gln Val Ala Val Tyr Gln Asp Val Tyr Ala Val Asp Gly Pro Thr
145          150          155          160
Ser Leu Tyr His Gln Ala Asn Lys Gly Val Arg Val Ala Tyr Trp Ile
165          170          175
Gly Phe Asp Thr Thr Pro Phe Met Phe Lys Asn Leu Ala Gly Ala Tyr
180          185          190
Pro Ser Tyr Ser Thr Asn Trp Ala Asp Glu Thr Val Leu Thr Ala Arg
195          200          205
Asn Ile Gly Leu Cys Ser Ser Asp Val Met Glu Arg Ser Arg Arg Gly
210          215          220
Met Ser Ile Leu Arg Lys Lys Tyr Leu Lys Pro Ser Asn Asn Val Leu
225          230          235          240
Phe Ser Val Gly Ser Thr Ile Tyr His Glu Lys Arg Asp Leu Leu Arg
245          250          255
Ser Trp His Leu Pro Ser Val Phe His Leu Arg Gly Lys Gln Asn Tyr
260          265          270
Thr Cys Arg Cys Glu Thr Ile Val Ser Cys Asp Gly Tyr Val Val Lys
275          280          285
Arg Ile Ala Ile Ser Pro Gly Leu Tyr Gly Lys Pro Ser Gly Tyr Ala
290          295          300
Ala Thr Met His Arg Glu Gly Phe Leu Cys Cys Lys Val Thr Asp Thr
305          310          315          320
Leu Asn Gly Glu Arg Val Ser Phe Pro Val Cys Thr Tyr Val Pro Ala
325          330          335
Thr Leu Cys Asp Gln Met Thr Gly Ile Leu Ala Thr Asp Val Ser Ala
340          345          350
Asp Asp Ala Gln Lys Leu Leu Val Gly Leu Asn Gln Arg Ile Val Val
355          360          365
Asn Gly Arg Thr Gln Arg Asn Thr Asn Thr Met Lys Asn Tyr Leu Leu
370          375          380
Pro Val Val Ala Gln Ala Phe Ala Arg Trp Ala Lys Glu Tyr Lys Glu
385          390          395          400

```

Asp Gln Glu Asp Glu Arg Pro Leu Gly Leu Arg Asp Arg Gln Leu Val
 405 410 415
 Met Gly Cys Cys Trp Ala Phe Arg Arg His Lys Ile Thr Ser Ile Tyr
 420 425 430
 Lys Arg Pro Asp Thr Gln Thr Ile Ile Lys Val Asn Ser Asp Phe His
 435 440 445
 Ser Phe Val Leu Pro Arg Ile Gly Ser Asn Thr Leu Glu Ile Gly Leu
 450 455 460
 Arg Thr Arg Ile Arg Lys Met Leu Glu Glu His Lys Glu Pro Ser Pro
 465 470 475 480
 Leu Ile Thr Ala Glu Asp Val Gln Glu Ala Lys Cys Ala Ala Asp Glu
 485 490 495
 Ala Lys Glu Val Arg Glu Ala Glu Glu Leu Arg Ala Ala Leu Pro Pro
 500 505 510
 Leu Ala Ala Asp Val Glu Glu Pro Thr Leu Glu Ala Asp Val Asp Leu
 515 520 525
 Met Leu Gln Glu Ala Gly Ala Gly Ser Val Glu Thr Pro Arg Gly Leu
 530 535 540
 Ile Lys Val Thr Ser Tyr Ala Gly Glu Asp Lys Ile Gly Ser Tyr Ala
 545 550 555 560
 Val Leu Ser Pro Gln Ala Val Leu Lys Ser Glu Lys Leu Ser Cys Ile
 565 570 575
 His Pro Leu Ala Glu Gln Val Ile Val Ile Thr His Ser Gly Arg Lys
 580 585 590
 Gly Arg Tyr Ala Val Glu Pro Tyr His Gly Lys Val Val Val Pro Glu
 595 600 605
 Gly His Ala Ile Pro Val Gln Asp Phe Gln Ala Leu Ser Glu Ser Ala
 610 615 620
 Thr Ile Val Tyr Asn Glu Arg Glu Phe Val Asn Arg Tyr Leu His His
 625 630 635 640
 Ile Ala Thr His Gly Gly Ala Leu Asn Thr Asp Glu Glu Tyr Tyr Lys
 645 650 655
 Thr Val Lys Pro Ser Glu His Asp Gly Glu Tyr Leu Tyr Asp Ile Asp
 660 665 670
 Arg Lys Gln Cys Val Lys Lys Glu Leu Val Thr Gly Leu Gly Leu Thr
 675 680 685
 Gly Glu Leu Val Asp Pro Pro Phe His Glu Phe Ala Tyr Glu Ser Leu
 690 695 700
 Arg Thr Arg Pro Ala Ala Pro Tyr Gln Val Pro Thr Ile Gly Val Tyr
 705 710 715 720
 Gly Val Pro Gly Ser Gly Lys Ser Gly Ile Ile Lys Ser Ala Val Thr
 725 730 735
 Lys Lys Asp Leu Val Val Ser Ala Lys Lys Glu Asn Cys Ala Glu Ile
 740 745 750
 Ile Arg Asp Val Lys Lys Met Lys Gly Leu Asp Val Asn Ala Arg Thr
 755 760 765
 Val Asp Ser Val Leu Leu Asn Gly Cys Lys His Pro Val Glu Thr Leu
 770 775 780
 Tyr Ile Asp Glu Ala Phe Ala Cys His Ala Gly Thr Leu Arg Ala Leu
 785 790 795 800
 Ile Ala Ile Ile Arg Pro Lys Lys Ala Val Leu Cys Gly Asp Pro Lys
 805 810 815
 Gln Cys Gly Phe Phe Asn Met Met Cys Leu Lys Val His Phe Asn His
 820 825 830

Glu Ile Cys Thr Gln Val Phe His Lys Ser Ile Ser Arg Arg Cys Thr
 835 840 845
 Lys Ser Val Thr Ser Val Val Ser Thr Leu Phe Tyr Asp Lys Lys Met
 850 855 860
 Arg Thr Thr Asn Pro Lys Glu Thr Lys Ile Val Ile Asp Thr Thr Gly
 865 870 875 880
 Ser Thr Lys Pro Lys Gln Asp Asp Leu Ile Leu Thr Cys Phe Arg Gly
 885 890 895
 Trp Val Lys Gln Leu Gln Ile Asp Tyr Lys Lys Asn Glu Ile Met Thr
 900 905 910
 Ala Ala Ala Ser Gln Gly Leu Thr Arg Lys Gly Val Tyr Ala Val Arg
 915 920 925
 Tyr Lys Val Asn Glu Asn Pro Leu Tyr Ala Pro Thr Ser Glu His Val
 930 935 940
 Asn Val Leu Leu Thr Arg Thr Glu Asp Arg Ile Val Trp Lys Thr Leu
 945 950 955 960
 Ala Gly Asp Pro Trp Ile Lys Thr Leu Thr Ala Lys Tyr Pro Gly Asn
 965 970 975
 Phe Thr Ala Thr Ile Glu Glu Trp Gln Ala Glu His Asp Ala Ile Met
 980 985 990
 Arg His Ile Leu Glu Arg Pro Asp Pro Thr Asp Val Phe Gln Asn Lys
 995 1000 1005
 Ala Asn Val Cys Trp Ala Lys Ala Leu Val Pro Val Leu Lys Thr Ala
 1010 1015 1020
 Gly Ile Asp Met Thr Thr Glu Gln Trp Asn Thr Val Asp Tyr Phe Glu
 1025 1030 1035 1040
 Thr Asp Lys Ala His Ser Ala Glu Ile Val Leu Asn Gln Leu Cys Val
 1045 1050 1055
 Arg Phe Phe Gly Leu Asp Leu Asp Ser Gly Leu Phe Ser Ala Pro Thr
 1060 1065 1070
 Val Pro Leu Ser Ile Arg Asn Asn His Trp Asp Asn Ser Pro Ser Pro
 1075 1080 1085
 Asn Met Tyr Gly Leu Asn Lys Glu Val Val Arg Gln Leu Ser Arg Arg
 1090 1095 1100
 Tyr Pro Gln Leu Pro Arg Ala Val Ala Thr Gly Arg Val Tyr Asp Met
 1105 1110 1115 1120
 Asn Thr Gly Thr Leu Arg Asn Tyr Asp Pro Arg Ile Asn Leu Val Pro
 1125 1130 1135
 Val Asn Arg Arg Leu Pro His Ala Leu Val Leu His His Asn Glu His
 1140 1145 1150
 Pro Gln Ser Asp Phe Ser Ser Phe Val Ser Lys Leu Lys Gly Arg Thr
 1155 1160 1165
 Val Leu Val Val Gly Glu Lys Leu Ser Val Pro Gly Lys Met Val Asp
 1170 1175 1180
 Trp Leu Ser Asp Arg Pro Glu Ala Thr Phe Arg Ala Arg Leu Asp Leu
 1185 1190 1195 1200
 Gly Ile Pro Gly Asp Val Pro Lys Tyr Asp Ile Ile Phe Val Asn Val
 1205 1210 1215
 Arg Thr Pro Tyr Lys Tyr His His Tyr Gln Gln Cys Glu Asp His Ala
 1220 1225 1230
 Ile Lys Leu Ser Met Leu Thr Lys Lys Ala Cys Leu His Leu Asn Pro
 1235 1240 1245
 Gly Gly Thr Cys Val Ser Ile Gly Tyr Gly Tyr Ala Asp Arg Ala Ser
 1250 1255 1260

Glu Ser Ile Ile Gly Ala Ile Ala Arg Gln Phe Lys Phe Ser Arg Val
 1265 1270 1275 1280
 Cys Lys Pro Lys Ser Ser Leu Glu Glu Thr Glu Val Leu Phe Val Phe
 1285 1290 1295
 Ile Gly Tyr Asp Arg Lys Ala Arg Thr His Asn Pro Tyr Lys Leu Ser
 1300 1305 1310
 Ser Thr Leu Thr Asn Ile Tyr Thr Gly Ser Arg Leu His Glu Ala Gly
 1315 1320 1325
 Cys Ala Pro Ser Tyr His Val Val Arg Gly Asp Ile Ala Thr Ala Thr
 1330 1335 1340
 Glu Gly Val Ile Ile Asn Ala Ala Asn Ser Lys Gly Gln Pro Gly Gly
 1345 1350 1355 1360
 Gly Val Cys Gly Ala Leu Tyr Lys Lys Phe Pro Glu Ser Phe Asp Leu
 1365 1370 1375
 Gln Pro Ile Glu Val Gly Lys Ala Arg Leu Val Lys Gly Ala Ala Lys
 1380 1385 1390
 His Ile Ile His Ala Val Gly Pro Asn Phe Asn Lys Val Ser Glu Val
 1395 1400 1405
 Glu Gly Asp Lys Gln Leu Ala Glu Ala Tyr Glu Ser Ile Ala Lys Ile
 1410 1415 1420
 Val Asn Asp Asn Asn Tyr Lys Ser Val Ala Ile Pro Leu Leu Ser Thr
 1425 1430 1435 1440
 Gly Ile Phe Ser Gly Asn Lys Asp Arg Leu Thr Gln Ser Leu Asn His
 1445 1450 1455
 Leu Leu Thr Ala Leu Asp Thr Thr Asp Ala Asp Val Ala Ile Tyr Cys
 1460 1465 1470
 Arg Asp Lys Lys Trp Glu Met Thr Leu Lys Glu Ala Val Ala Arg Arg
 1475 1480 1485
 Glu Ala Val Glu Glu Ile Cys Ile Ser Asp Asp Ser Ser Val Thr Glu
 1490 1495 1500
 Pro Asp Ala Glu Leu Val Arg Val His Pro Lys Ser Ser Leu Ala Gly
 1505 1510 1515 1520
 Arg Lys Gly Tyr Ser Thr Ser Asp Gly Lys Thr Phe Ser Tyr Leu Glu
 1525 1530 1535
 Gly Thr Lys Phe His Gln Ala Ala Lys Asp Ile Ala Glu Ile Asn Ala
 1540 1545 1550
 Met Trp Pro Val Ala Thr Glu Ala Asn Glu Gln Val Cys Met Tyr Ile
 1555 1560 1565
 Leu Gly Glu Ser Met Ser Ser Ile Arg Ser Lys Cys Pro Val Glu Glu
 1570 1575 1580
 Ser Glu Ala Ser Thr Pro Pro Ser Thr Leu Pro Cys Leu Cys Ile His
 1585 1590 1595 1600
 Ala Met Thr Pro Glu Arg Val Gln Arg Leu Lys Ala Ser Arg Pro Glu
 1605 1610 1615
 Gln Ile Thr Val Cys Ser Ser Phe Pro Leu Pro Lys Tyr Arg Ile Thr
 1620 1625 1630
 Gly Val Gln Lys Ile Gln Cys Ser Gln Pro Ile Leu Phe Ser Pro Lys
 1635 1640 1645
 Val Pro Ala Tyr Ile His Pro Arg Lys Tyr Leu Val Glu Thr Pro Pro
 1650 1655 1660
 Val Asp Glu Thr Pro Glu Pro Ser Ala Glu Asn Gln Ser Thr Glu Gly
 1665 1670 1675 1680
 Thr Pro Glu Gln Pro Pro Leu Ile Thr Glu Asp Glu Thr Arg Thr Arg
 1685 1690 1695

Thr Pro Glu Pro Ile Ile Ile Glu Glu Glu Glu Glu Asp Ser Ile Ser
 1700 1705 1710
 Leu Leu Ser Asp Gly Pro Thr His Gln Val Leu Gln Val Glu Ala Asp
 1715 1720 1725
 Ile His Gly Pro Pro Ser Val Ser Ser Ser Trp Ser Ile Pro His
 1730 1735 1740
 Ala Ser Asp Phe Asp Val Asp Ser Leu Ser Ile Leu Asp Thr Leu Glu
 1745 1750 1755 1760
 Gly Ala Ser Val Thr Ser Gly Ala Thr Ser Ala Glu Thr Asn Ser Tyr
 1765 1770 1775
 Phe Ala Lys Ser Met Glu Phe Leu Ala Arg Pro Val Pro Ala Pro Arg
 1780 1785 1790
 Thr Val Phe Arg Asn Pro Pro His Pro Ala Pro Arg Thr Arg Thr Pro
 1795 1800 1805
 Ser Leu Ala Pro Ser Arg Ala Cys Ser Arg Thr Ser Leu Val Ser Thr
 1810 1815 1820
 Pro Pro Gly Val Asn Arg Val Ile Thr Arg Glu Glu Leu Glu Ala Leu
 1825 1830 1835 1840
 Thr Pro Ser Arg Thr Pro Ser Arg Ser Val Ser Arg Thr Ser Leu Val
 1845 1850 1855
 Ser Asn Pro Pro Gly Val Asn Arg Val Ile Thr Arg Glu Glu Phe Glu
 1860 1865 1870
 Ala Phe Val Ala Gln Gln Gln Arg Phe Asp Ala Gly Ala Tyr Ile Phe
 1875 1880 1885
 Ser Ser Asp Thr Gly Gln Gly His Leu Gln Gln Lys Ser Val Arg Gln
 1890 1895 1900
 Thr Val Leu Ser Glu Val Val Leu Glu Arg Thr Glu Leu Glu Ile Ser
 1905 1910 1915 1920
 Tyr Ala Pro Arg Leu Asp Gln Glu Lys Glu Glu Leu Leu Arg Lys Lys
 1925 1930 1935
 Leu Gln Leu Asn Pro Thr Pro Ala Asn Arg Ser Arg Tyr Gln Ser Arg
 1940 1945 1950
 Lys Val Glu Asn Met Lys Ala Ile Thr Ala Arg Arg Ile Leu Gln Gly
 1955 1960 1965
 Leu Gly His Tyr Leu Lys Ala Glu Gly Lys Val Glu Cys Tyr Arg Thr
 1970 1975 1980
 Leu His Pro Val Pro Leu Tyr Ser Ser Ser Val Asn Arg Ala Phe Ser
 1985 1990 1995 2000
 Ser Pro Lys Val Ala Val Glu Ala Cys Asn Ala Met Leu Lys Glu Asn
 2005 2010 2015
 Phe Pro Thr Val Ala Ser Tyr Cys Ile Ile Pro Glu Tyr Asp Ala Tyr
 2020 2025 2030
 Leu Asp Met Val Asp Gly Ala Ser Cys Cys Leu Asp Thr Ala Ser Phe
 2035 2040 2045
 Cys Pro Ala Lys Leu Arg Ser Phe Pro Lys Lys His Ser Tyr Leu Glu
 2050 2055 2060
 Pro Thr Ile Arg Ser Ala Val Pro Ser Ala Ile Gln Asn Thr Leu Gln
 2065 2070 2075 2080
 Asn Val Leu Ala Ala Thr Lys Arg Asn Cys Asn Val Thr Gln Met
 2085 2090 2095
 Arg Glu Leu Pro Val Leu Asp Ser Ala Ala Phe Asn Val Glu Cys Phe
 2100 2105 2110
 Lys Lys Tyr Ala Cys Asn Asn Glu Tyr Trp Glu Thr Phe Lys Glu Asn
 2115 2120 2125

22

Pro Ile Arg Leu Thr Glu Glu Asn Val Val Asn Tyr Ile Thr Lys Leu
 2130 2135 2140
 Lys Gly Pro Lys Ala Ala Ala Leu Phe Ala Lys Thr His Asn Leu Asn
 2145 2150 2155 2160
 Met Leu Gln Asp Ile Pro Met Asp Arg Phe Val Met Asp Leu Lys Arg
 2165 2170 2175
 Asp Val Lys Val Thr Pro Gly Thr Lys His Thr Glu Glu Arg Pro Lys
 2180 2185 2190
 Val Gln Val Ile Gln Ala Ala Asp Pro Leu Ala Thr Ala Tyr Leu Cys
 2195 2200 2205
 Gly Ile His Arg Glu Leu Val Arg Arg Leu Asn Ala Val Leu Leu Pro
 2210 2215 2220
 Asn Ile His Thr Leu Phe Asp Met Ser Ala Glu Asp Phe Asp Ala Ile
 2225 2230 2235 2240
 Ile Ala Glu His Phe Gln Pro Gly Asp Cys Val Leu Glu Thr Asp Ile
 2245 2250 2255
 Ala Ser Phe Asp Lys Ser Glu Asp Asp Ala Met Ala Leu Thr Ala Leu
 2260 2265 2270
 Met Ile Leu Glu Asp Leu Gly Val Asp Ala Glu Leu Leu Thr Leu Ile
 2275 2280 2285
 Glu Ala Ala Phe Gly Glu Ile Ser Ser Ile His Leu Pro Thr Lys Thr
 2290 2295 2300
 Lys Phe Lys Phe Gly Ala Met Met Lys Ser Gly Met Phe Leu Thr Leu
 2305 2310 2315 2320
 Phe Val Asn Thr Val Ile Asn Ile Val Ile Ala Ser Arg Val Leu Arg
 2325 2330 2335
 Glu Arg Leu Thr Gly Ser Pro Cys Ala Ala Phe Ile Gly Asp Asp Asn
 2340 2345 2350
 Ile Val Lys Gly Val Lys Ser Asp Lys Leu Met Ala Asp Arg Cys Ala
 2355 2360 2365
 Thr Trp Leu Asn Met Glu Val Lys Ile Ile Asp Ala Val Val Gly Glu
 2370 2375 2380
 Lys Ala Pro Tyr Phe Cys Gly Gly Phe Ile Leu Cys Asp Ser Val Thr
 2385 2390 2395 2400
 Gly Thr Ala Cys Arg Val Ala Asp Pro Leu Lys Arg Leu Phe Lys Leu
 2405 2410 2415
 Gly Lys Pro Leu Ala Ala Asp Asp Glu His Asp Asp Asp Arg Arg Arg
 2420 2425 2430
 Ala Leu His Glu Glu Ser Thr Arg Trp Asn Arg Val Gly Ile Leu Ser
 2435 2440 2445
 Glu Leu Cys Lys Ala Val Glu Ser Arg Tyr Glu Thr Val Gly Thr Ser
 2450 2455 2460
 Ile Ile Val Met Ala Met Thr Thr Leu Ala Ser Ser Val Lys Ser Phe
 2465 2470 2475 2480
 Ser Tyr Leu Arg Gly Ala Pro Ile Thr Leu Tyr Gly
 2485 2490

<210> 4

<211> 1476

<212> DNA

<213> Artificial Sequence

<220>

 <223> Description of Artificial Sequence; Note =
 synthetic construct

<221> CDS

<222> (1)... (1476)

<400> 4

atg gct gcg aga gcg tca ata tta aga ggg gaa aaa tta gat aaa tgg	48
Met Ala Ala Arg Ala Ser Ile Leu Arg Gly Glu Lys Leu Asp Lys Trp	
1 5 10 15	

gaa aag att agg tta agg cca ggg gga aag aaa cat tat atg tta aaa	96
Glu Lys Ile Arg Leu Arg Pro Gly Gly Lys Lys His Tyr Met Leu Lys	
20 25 30	

cac ata gta tgg gcg agc agg gag ctg gaa aga ttt gca ctt aac cct	144
His Ile Val Trp Ala Ser Arg Glu Leu Glu Arg Phe Ala Leu Asn Pro	
35 40 45	

ggc ctt tta gaa aca tca gaa gga tgt aaa caa ata atg aaa cag cta	192
Gly Leu Leu Glu Thr Ser Glu Gly Cys Lys Gln Ile Met Lys Gln Leu	
50 55 60	

caa cca gct ctg cag aca gga aca gag gaa ctt aaa tca tta tac aac	240
Gln Pro Ala Leu Gln Thr Gly Thr Glu Glu Leu Lys Ser Leu Tyr Asn	
65 70 75 80	

aca gta gca act ctc tat tgt gta cat gaa aag ata gaa gta cga gac	288
Thr Val Ala Thr Leu Tyr Cys Val His Glu Lys Ile Glu Val Arg Asp	
85 90 95	

acc aag gaa gcc tta gat aag ata gag gaa gaa caa aac aaa tgt cag	336
Thr Lys Glu Ala Leu Asp Lys Ile Glu Glu Glu Gln Asn Lys Cys Gln	
100 105 110	

caa aaa acg cag cag gca aaa gcg gct gac ggg aaa gtc agt caa aat	384
Gln Lys Thr Gln Gln Ala Lys Ala Ala Asp Gly Lys Val Ser Gln Asn	
115 120 125	

tat cct ata gtg cag aat ctc caa ggg caa atg gta cat caa gcc ata	432
Tyr Pro Ile Val Gln Asn Leu Gln Gly Gln Met Val His Gln Ala Ile	
130 135 140	

tca cct aga acc ttg aat gca tgg gta aaa gta ata gaa gaa aag gct	480
Ser Pro Arg Thr Leu Asn Ala Trp Val Lys Val Ile Glu Glu Lys Ala	
145 150 155 160	

ttt agc cca gag gta ata ccc atg ttt aca gca tta tca gaa gga gcc	528
Phe Ser Pro Glu Val Ile Pro Met Phe Thr Ala Leu Ser Glu Gly Ala	
165 170 175	

acc cca caa gat tta aac acc atg tta aat aca gtg ggg gga cac caa	576
Thr Pro Gln Asp Leu Asn Thr Met Leu Asn Thr Val Gly Gly His Gln	
180 185 190	

gca gcc atg caa atg tta aaa gat act att aat gaa gag gct gca gaa	624
Ala Ala Met Gln Met Leu Lys Asp Thr Ile Asn Glu Glu Ala Ala Glu	
195 200 205	

tgg gat aga tta cat cca gtc cat gcg ggg cct att gca cca ggc cag Trp Asp Arg Leu His Pro Val His Ala Gly Pro Ile Ala Pro Gly Gln 210 215 220	672
atg aga gaa cca agg gga agt gac ata gca gga act act agt acc ctt Met Arg Glu Pro Arg Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu 225 230 235 240	720
cag gaa caa ata gca tgg atg aca agt aac cca cct att cca gtg gga Gln Glu Gln Ile Ala Trp Met Thr Ser Asn Pro Pro Ile Pro Val Gly 245 250 255	768
gac atc tat aaa aga tgg ata att ctg ggg tta aat aaa ata gtg aga Asp Ile Tyr Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg 260 265 270	816
atg tat agc cgg gtc agc att ttg gac ata aga caa ggg cca aag gaa Met Tyr Ser Pro Val Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu 275 280 285	864
ccc ttt cga gac tat gta gat cgg ttc ttt aaa act tta aga gct gaa Pro Phe Arg Asp Tyr Val Asp Arg Phe Phe Lys Thr Leu Arg Ala Glu 290 295 300	912
caa gct aca'caa gaa gta aaa aat tgg atg aca gac acc ttg tta gtc Gln Ala Thr Gln Glu Val Lys Asn Trp Met Thr Asp Thr Leu Leu Val 305 310 315 320	960
caa aat gcg aac cca gat tgt aag acc att ttg aga gca tta gga cca Gln Asn Ala Asn Pro Asp Cys Lys Thr Ile Leu Arg Ala Leu Gly Pro 325 330 335	1008
ggg gct aca tta gaa gaa atg atg aca gca tgt caa ggg gtg gga gga Gly Ala Thr Leu Glu Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly 340 345 350	1056
cct ggc cac aaa gca aga gta ttg gct gag gca atg agt caa aca aac Pro Gly His Lys Ala Arg Val Leu Ala Glu Ala Met Ser Gln Thr Asn 355 360 365	1104
agt gga aac ata atg atg cag aga agc aat ttt aaa ggc cct aga aga Ser Gly Asn Ile Met Met Gln Arg Ser Asn Phe Lys Gly Pro Arg Arg 370 375 380	1152
att gtt aaa tgt ttt aac tgt ggc aag gaa ggg cac ata gcc aga aat Ile Val Lys Cys Phe Asn Cys Gly Lys Glu Gly His Ile Ala Arg Asn 385 390 395 400	1200
tgc aga gcc cct agg aaa aaa ggc tgt tgg aaa tgt gga aaa gaa gga Cys Arg Ala Pro Arg Lys Lys Gly Cys Trp Lys Lys Gly Glu Gly 405 410 415	1248
cac caa atg aaa gac tgc act gag agg cag gct aat ttt tta ggg aaa His Gln Met Lys Asp Cys Thr Glu Arg Gln Ala Asn Phe Leu Gly Lys 420 425 430	1296

att tgg cct tcc cac aag ggg agg cca ggg aat ttc ctt cag aac aga 1344
 ile Trp Pro Ser His Lys Gly Arg Pro Gly Asn Phe Leu Gln Asn Arg
 435 440 445

cca gag cca aca gcc cca cca gca gag agc ttc agg ttc gaa gag aca 1392
 Pro Glu Pro Thr Ala Pro Pro Ala Glu Ser Phe Arg Phe Glu Glu Thr
 450 455 460

acc ccc gct cgg aaa cag gag cgg ata gaa agg gaa ccc tta act tcc 1440
 Thr Pro Ala Pro Lys Gln Glu Pro ile Glu Arg Glu Pro Leu Thr Ser
 465 470 475 480

ctc aaa tca ctc ttt ggc agc gac ccc ttg tct caa 1476
 Leu Lys Ser Leu Phe Gly Ser Asp Pro Leu Ser Gln
 485 490

<210> 5

<211> 492

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 5

Met Ala Ala Arg Ala Ser ile Leu Arg Gly Glu Lys Leu Asp Lys Trp
 1 5 10 15
 Glu Lys ile Arg Leu Arg Pro Gly Gly Lys Lys His Tyr Met Leu Lys
 20 25 30
 His ile Val Trp Ala Ser Arg Glu Leu Glu Arg Phe Ala Leu Asn Pro
 35 40 45
 Gly Leu Leu Glu Thr Ser Glu Gly Cys Lys Gln ile Met Lys Gln Leu
 50 55 60
 Gln Pro Ala Leu Gln Thr Gly Thr Glu Glu Leu Lys Ser Leu Tyr Asn
 65 70 75 80
 Thr Val Ala Thr Leu Tyr Cys Val His Glu Lys ile Glu Val Arg Asp
 85 90 95
 Thr Lys Glu Ala Leu Asp Lys ile Glu Glu Glu Gln Asn Lys Cys Gln
 100 105 110
 Gln Lys Thr Gln Gln Ala Lys Ala Ala Asp Gly Lys Val Ser Gln Asn
 115 120 125
 Tyr Pro ile Val Gln Asn Leu Gln Gly Gln Met Val His Gln Ala ile
 130 135 140
 Ser Pro Arg Thr Leu Asn Ala Trp Val Lys Val ile Glu Glu Lys Ala
 145 150 155 160
 Phe Ser Pro Glu Val ile Pro Met Phe Thr Ala Leu Ser Glu Gly Ala
 165 170 175
 Thr Pro Gln Asp Leu Asn Thr Met Leu Asn Thr Val Gly Gly His Gln
 180 185 190
 Ala Ala Met Gln Met Leu Lys Asp Thr ile Asn Glu Glu Ala Ala Glu
 195 200 205

26

```

Trp Asp Arg Leu His Pro Val His Ala Gly Pro Ile Ala Pro Gly Gln
210                215                220
Met Arg Glu Pro Arg Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu
225                230                235                240
Gln Glu Gln Ile Ala Trp Met Thr Ser Asn Pro Pro Ile Pro Val Gly
245                250                255
Asp Ile Tyr Lys Arg Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg
260                265                270
Met Tyr Ser Pro Val Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu
275                280                285
Pro Phe Arg Asp Tyr Val Asp Arg Phe Phe Lys Thr Leu Arg Ala Glu
290                295                300
Gln Ala Thr Gln Glu Val Lys Asn Trp Met Thr Asp Thr Leu Leu Val
305                310                315                320
Gln Asn Ala Asn Pro Asp Cys Lys Thr Ile Leu Arg Ala Leu Gly Pro
325                330                335
Gly Ala Thr Leu Glu Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly
340                345                350
Pro Gly His Lys Ala Arg Val Leu Ala Glu Ala Met Ser Gln Thr Asn
355                360                365
Ser Gly Asn Ile Met Met Gln Arg Ser Asn Phe Lys Gly Pro Arg Arg
370                375                380
Ile Val Lys Cys Phe Asn Cys Gly Lys Glu Gly His Ile Ala Arg Asn
385                390                395                400
Cys Arg Ala Pro Arg Lys Lys Gly Cys Trp Lys Cys Gly Lys Glu Gly
405                410                415
His Gln Met Lys Asp Cys Thr Glu Arg Gln Ala Asn Phe Leu Gly Lys
420                425                430
Ile Trp Pro Ser His Lys Gly Arg Pro Gly Asn Phe Leu Gln Asn Arg
435                440                445
Pro Glu Pro Thr Ala Pro Pro Ala Glu Ser Phe Arg Phe Glu Glu Thr
450                455                460
Thr Pro Ala Pro Lys Gln Glu Pro Ile Glu Arg Glu Pro Leu Thr Ser
465                470                475                480
Leu Lys Ser Leu Phe Gly Ser Asp Pro Leu Ser Gln
485                490

```

<210> 6

<211> 813

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<221> CDS

<222> (1)...(813)

<400> 6

```

atg agc cat att caa cgg gaa acg tct tgc tgc agg ccg cga tta aat
Met Ser His Ile Gln Arg Glu Thr Ser Cys Ser Arg Pro Arg Leu Asn
1          5          10          15

```

48

27

tcc aac atg gat gct gat tta tat ggg tat aaa tgg gct cgc gat aat Ser Asn Met Asp Ala Asp Leu Tyr Gly Tyr Lys Trp Ala Arg Asp Asn 20 25 30	96
gtc ggg caa tca ggt gcg aca atc tat cga ttg tat ggg aag ccc gat Val Gly Gln Ser Gly Ala Thr Ile Tyr Arg Leu Tyr Gly Lys Pro Asp 35 40 45	144
gcg cca gag ttg ttt ctg aaa cat ggc aaa ggt agc gtt gcc aat gat Ala Pro Glu Leu Phe Leu Lys His Gly Lys Gly Ser Val Ala Asn Asp 50 55 60	192
gtt aca gat gag atg gtc aga cta aac tgg ctg acg gaa ttt atg cct Val Thr Asp Glu Met Val Arg Leu Asn Trp Leu Thr Glu Phe Met Pro 65 70 75 80	240
ctt ccg acc atc aag cat ttt atc cgt act cct gat gat gca tgg tta Leu Pro Thr Ile Lys His Phe Ile Arg Thr Pro Asp Asp Ala Trp Leu 85 90 95	288
ctc acc act gcg atc ccc ggg aaa aca gca ttc cag gta tta gaa gaa Leu Thr Thr Ala Ile Pro Gly Lys Thr Ala Phe Gln Val Leu Glu Glu 100 105 110	336
tat cct gat tca ggt gaa aat att gtt gat gcg ctg gca gtg ttc ctg Tyr Pro Asp Ser Gly Glu Asn Ile Val Asp Ala Leu Ala Val Phe Leu 115 120 125	384
cgc cgg ttg cat tgg att cct gtt tgt aat tgt cct ttt aac agc gat Arg Arg Leu His Ser Ile Pro Val Cys Asn Cys Pro Phe Asn Ser Asp 130 135 140	432
cgc gta ttt cgt ctc gct cag gcg caa tca cga atg aat aac ggt ttg Arg Val Phe Arg Leu Ala Gln Ala Gln Ser Arg Met Asn Asn Gly Leu 145 150 155 160	480
gtt gat gcg agt gat ttt gat gac gag cgt aat ggc tgg cct gtt gaa Val Asp Ala Ser Asp Phe Asp Asp Glu Arg Asn Gly Trp Pro Val Glu 165 170 175	528
caa gtc tgg aaa gaa atg cat aag ctt ttg cca ttc tca ccg gat tca Gln Val Trp Lys Glu Met His Lys Leu Leu Pro Phe Ser Pro Asp Ser 180 185 190	576
gtc gtc act cat ggt gat ttc tca ctt gat aac ctt att ttt gac gag Val Val Thr His Gly Asp Phe Ser Leu Asp Asn Leu Ile Phe Asp Glu 195 200 205	624
ggg aaa tta ata ggt tgt att gat gtt gga cga gtc gga atc gca gac Gly Lys Leu Ile Gly Cys Ile Asp Val Gly Arg Val Gly Ile Ala Asp 210 215 220	672
oga tac cag gat ctt gcc atc cta tgg aac tgc ctc ggt gag ttt tct Arg Tyr Gln Asp Leu Ala Ile Leu Trp Asn Cys Leu Gly Glu Phe Ser 225 230 235 240	720

```

cct tca tta cag aaa cgg ctt ttt caa aaa tat ggt att gat aat cct      768
Pro Ser Leu Gln Lys Arg Leu Phe Gln Lys Tyr Gly Ile Asp Asn Pro
                245                      250                      255

```

```

gat atg aat aaa ttg cag ttt cat ttg atg ctc gat gag ttt ttc      813
Asp Met Asn Lys Leu Gln Phe His Leu Met Leu Asp Glu Phe Phe
                260                      265                      270

```

<210> 7

<211> 271

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 7

```

Met Ser His Ile Gln Arg Glu Thr Ser Cys Ser Arg Pro Arg Leu Asn
 1              5              10              15
Ser Asn Met Asp Ala Asp Leu Tyr Gly Tyr Lys Trp Ala Arg Asp Asn
      20              25              30
Val Gly Gln Ser Gly Ala Thr Ile Tyr Arg Leu Tyr Gly Lys Pro Asp
 35              40              45
Ala Pro Glu Leu Phe Leu Lys His Gly Lys Gly Ser Val Ala Asn Asp
 50              55              60
Val Thr Asp Glu Met Val Arg Leu Asn Trp Leu Thr Glu Phe Met Pro
 65              70              75              80
Leu Pro Thr Ile Lys His Phe Ile Arg Thr Pro Asp Asp Ala Trp Leu
      85              90              95
Leu Thr Thr Ala Ile Pro Gly Lys Thr Ala Phe Gln Val Leu Glu Glu
 100              105              110
Tyr Pro Asp Ser Gly Glu Asn Ile Val Asp Ala Leu Ala Val Phe Leu
 115              120              125
Arg Arg Leu His Ser Ile Pro Val Cys Asn Cys Pro Phe Asn Ser Asp
 130              135              140
Arg Val Phe Arg Leu Ala Gln Ala Gln Ser Arg Met Asn Asn Gly Leu
 145              150              155              160
Val Asp Ala Ser Asp Phe Asp Asp Glu Arg Asn Gly Trp Pro Val Glu
 165              170              175
Gln Val Trp Lys Glu Met His Lys Leu Leu Pro Phe Ser Pro Asp Ser
 180              185              190
Val Val Thr His Gly Asp Phe Ser Leu Asp Asn Leu Ile Phe Asp Glu
 195              200              205
Gly Lys Leu Ile Gly Cys Ile Asp Val Gly Arg Val Gly Ile Ala Asp
 210              215              220
Arg Tyr Gln Asp Leu Ala Ile Leu Trp Asn Cys Leu Gly Glu Phe Ser
 225              230              235              240
Pro Ser Leu Gln Lys Arg Leu Phe Gln Lys Tyr Gly Ile Asp Asn Pro
 245              250              255
Asp Met Asn Lys Leu Gln Phe His Leu Met Leu Asp Glu Phe Phe
 260              265              270

```


29

<210> 8
 <211> 5076
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 8
 ataggcgccg catgagagaa gccagaccaa attacctacc aaaaatggag aaagttoacg 60
 ttgacatoga ggaagacagc coattctcca gagcttttga cggagcttc cgcgacttg 120
 aggtagaagc aacgacggtc actgataatg accatgctaa tgcagagcg ttttcgcatc 180
 tggcttcaaa actgatcgaa acggaggtgg acccatccga cagcatcctt gacattggaa 240
 gtgcgcccgc cgcgagaatg tattctaagc acaagtatca ttgtatctgt cagatgagat 300
 gtgcggaaga tccggacaga ttgtataagt atgcaactaa gctgaagaaa aactgtaaagg 360
 aaataactga taaggaaattg gacaagaaaa tgaaggagct cgcgcgcgtc atgagcgacc 420
 ctgacctgga aactgagact atgtgctccc acgacgaaga gtctgtctgc tacgaagggc 480
 aagtgcctgt ttaccaggat gtatacgcgg ttgacggacc ctataactct ctacggctaa 540
 cctgaatgga ctacgacata gtctagtccg ccaagatggt ccggttccag ccaatgtatc 600
 cgtatgcacc aatgccttat cgcacccggt tcgcccgcgc gcgcagggcc tgggtcccca 660
 gaaccgaccc ttttctggcg atgcagggtgc aggaattaac ccgctcgatg gctaacctga 720
 cgttgaagca acgcggggac gcgccacctg aggggcccac cgtaagaaa ccgaagaagg 780
 aggcctcgca aaaaacagaaa gggggaggcc aagggaagaa gaagaagaac caagggaaga 840
 agaaggctaa gacaggggcg cctaactcga aggcacagaa tggaaacaa aagaagacca 900
 acaaaaacac aggcgaagga cagcgcatgg tcatgaaatt ggaatctgac aagacgttcc 960
 caatcatggt ggaagggaag ataaacggct acgcttctgt ggtcggaggg aagtattatc 1020
 ggcgatgca tgtggaaggc aagatcgaca acgacgttct ggcgcgcgtt aagacgaaga 1080
 aagaccccaa atacgatctt gagtatgcag atgtgccaca gaacatgcgg gccgatacat 1140
 tcaaatcac ccattgaaaa ccccaaggct attacagctg gcactcatga gcagtcacat 1200
 atgaaaatgg gogtttcaag gtgcgcgaag gacttggggc caagggaagc aggcgacac 1260
 ccattctgga taaccaggga cgggtggtcg ctattgtgct gggagggtgt aatgaaggat 1320
 ctaggacagc ctttctagtc gtcatgtgga acgagaagg agttacogtg aagtaacttc 1380
 cgggaactgt cgagcaatgg tcaactagtga ccacatgtg tctgctcgcc aatgtgactg 1440
 tccctatgtc tcaaccacca attttgctac acagaaaacc agcagagact ttggccatgc 1500
 tcagcgttaa catccctgct gggaggatca gccgtaatta ttataattgg ctgggtgctg 1560
 gctactattg tggccatgta cgtgctgacc aaccagaanc ataattgaat acagacgcaa 1620
 ttggcaagct gcttaccatag aactcgcggc gattggcatg ccgctttaa atttttattt 1680
 tatttttctt tctttttccg aatcggattt tgtttttaat atttcaaaaa aaaaaaaaat 1740
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ggggaagagc cgcgcgcgcg ctgggctacg 1800
 tcttctggcg gtctgcgacg cgaggctgga tggccttccc cattatgatt ctctctgctt 1860
 cggcgggcat cgggatggcc gcgttgcaag ccactgctgc caggcaggtg gatgaagacc 1920
 atcaggagca gcttcaagga tgcctcgcgg ctcttaccag cctaactcgt atcaactggc 1980
 cgcgtgatcgt cagcgcgatt tatgcgcgct cggcgagcac atggacggg ttggcatgga 2040
 ttgtaggcgc cgcctatata cttgtctgcc tccccgcgtt ggcgtgcggt gcattggagcc 2100
 gggccacctc gacctgaatg gaagccggcg gcacctcgct aacggattca ccactccaa 2160
 aatggagacc aatacaattct tgcggagaac tgtgaatgc caaccacacc ctggcgagaa 2220
 catatccatc gogtcgcgca tctccagcag ccgcacggcg cgcactcgg cgagcgttgg 2280
 gtctcggcca cgggtgcgca tgactgtgct cctgtctgtg aggacccggc tagcggtggc 2340
 ggggtgcttt actggttagc agaataatc acogatacgc gacgcgaagt gaagcagact 2400
 ctgctgcaaa aogtctgcga cctgagcaac aacatgaatg gtcttgggtt tccgtgtttc 2460
 gtaaatgtctg gaaacgcgga agtcagcgcc ctgcaaccatt atgttccgga tctgcactgc 2520
 aggatgtctg tggctacact gtggaacacc tacactctga ttaaocgaac gctggcactg 2580
 accctgagtg attttctctt ggtcccgccg catccatacc gccagtgttt taacctcaaa 2640
 acgttccagt aaccgggcat gttcatcacc agtaaccctg atcgtgagca tctctctctg 2700

tttcatcggt	atcattacc	ccatgaacag	aaatccccc	tacacggagg	catcagtgac	2760
caaacaggaa	aaaaccggcc	ttaacatggc	ccgctttatc	agaagccaga	cattaaagct	2820
tctggagaaa	ctcaacgagc	tggacgcgga	tgaacaggca	gacatctgtg	aatcgctcca	2880
cgaccacgct	gatgagcttt	accgcagctg	cctcgccgct	ttcggtgatg	acggtgaaaa	2940
cctctgacac	atgcagctcc	cggagacggt	cacagcttgt	ctgtaagcgg	atgcggggag	3000
cagacaagcc	cgtcagggcg	cgtcagcggg	tgctggcggg	tgctggggcg	cagccatgac	3060
ccagtcacgt	agcgatagcg	gagtgatatac	tggcttaact	atgcggcattc	agagcagatt	3120
gtactgagag	tgcacatata	atgcggtgtg	aaatcacgca	cagatgcgtg	aggagaaatt	3180
accgcatacg	gcgctctccc	gcttctctgc	tcactgaactc	gctgogctcg	gtcgttcggc	3240
tgccgcgagc	ggtatcagct	cactcaaaag	cggtaatacg	gttatccaca	gaatcagggg	3300
ataacgcagg	aaagaacatg	tgagcaaaa	gccagcaaaa	ggccagggaac	cgtaaaaagg	3360
ccgcgttgct	ggcgcttttc	cataggctcc	gccccctcga	cgagcatcac	aaaaatcgac	3420
gctcaagtca	agggtggcga	aaccgcagac	gactataaag	ataccaggcg	tttccccctg	3480
gaagctccct	cgtgcgctct	cctgttccga	cctgcgcgt	taccggatac	ctgtccgcct	3540
ttctcccttc	gggaagcggtg	gcgctttctc	atagctcagc	ctgtagggtat	ctcagttcgg	3600
tgtaggctgt	tgcctccaag	ctgggctgtg	tgcacgaacc	ccccgttcag	ccgcagcgtc	3660
gcgccttate	cgttaactat	cgtcttgagt	ccaaccgggt	aagacacgac	ttatcgccac	3720
tgccagcagc	cactggtaac	aggattagca	gagcgaggta	tgtaggcggt	gctcacagat	3780
tcttgaagtg	gtggcctaac	tacggctaca	ctagaaggac	agtatttggt	atctcgctcc	3840
tgcgtgaagc	agttaccctc	ggaaaaagag	ttggtagctc	ttgatccggc	aaaaaaacca	3900
ccgctggtag	cggctggttt	tttgtttgca	agcagcagat	tacggccaga	aaaaaaggat	3960
ctcaagaaga	tcccttgatc	ttttctacgg	ggctctgacg	tcagtggaac	gaaaactcac	4020
gttaagggat	tttggtcatg	aaacaataaa	ctgtctgctt	acataaaccg	taatacaagg	4080
gggtgtatga	gccatattca	acgggaaaacg	tcttgctcga	ggccgcgatt	aaattccaac	4140
atggatgcgtg	atttatatgg	gtataaatgg	gctcgcgata	atgtcgggca	atcaggtcgc	4200
acaactatc	gattgtatgg	gaagcccgat	gcgcagaggt	tgcttctgaa	acatggccaa	4260
ggtagcgctg	ccaatgatgt	tacagatgag	atggctcagac	taaaactggct	gacggaaatt	4320
atgcctcttc	cgaccatcaa	gcattttatc	cgtactcctg	atgatgcagt	gttactcacc	4380
actgcgatac	ccgggaaaac	agcattctag	gtattagaag	aatactctga	ttcaggtgaa	4440
aataattgtg	atgcgctggc	agtgttctcg	cgccgggtgc	attcgattcc	tggttgtaatt	4500
tgtcctttta	acagcgatcg	cgtatttctg	ctcgctcagg	cgcaatcacg	aatgaataac	4560
gggttggttg	atgcgagtga	ttttgatgac	gagcgtaagt	gctggcctgt	tgaacaagtc	4620
tggaaagaaa	tgcataagct	tttgccattc	tcacgcgatt	cagtcgtcac	tcattggtgat	4680
ttctcacttg	ataaccttat	ttttgacgag	gggaaattaa	taggttgtag	tatgattgga	4740
cgagtcggaa	tgcgagccg	ataccaggat	cttgccatcac	tatggaaactg	cctcggtgag	4800
ttttctcctt	cattacagaa	acggcttttt	caaaaatgat	gtattgataa	tctctgatatg	4860
aataaaattgc	agtttctatt	gatgctcgat	gagttttttc	aagaattctc	atgtttgaca	4920
gctttacatc	gataagcttt	aatgcggtag	tttatccagc	ttaaattgct	aacgcagtc	4980
ggcacgcgtg	atgaaatcta	acaatgcgct	catcgtctac	ctcgccacgc	tcaccctgga	5040
tgcgtcttag	aggatcccta	atacgactca	ctatag			5076

<210> 9

<211> 1026

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<221> CDS

<222> (1)...(1026)

31

<400> 9			
atg ttc ccg ttc cag cca atg tat ccg atg cag cca atg ccc tat cgc		48	
Met Phe Pro Phe Gln Pro Met Tyr Pro Met Gln Pro Met Tyr Arg			
1 5 10 15			
aac ccg ttc gcg gcc ccg cgc agg ccc tgg ttc cgc aga acc gac cct		96	
Asn Pro Phe Ala Ala Pro Arg Arg Pro Trp Phe Pro Arg Thr Asp Pro			
20 25 30			
ttt ctg gcg atg cag gtg cag gaa tta acc cgc tgc atg gct aac ctg		144	
Phe Leu Ala Met Gln Val Gln Glu Leu Thr Arg Ser Met Ala Asn Leu			
35 40 45			
acg ttc aag caa cgc cgg gac gcg cca cct gag ggg cca tcc gct aag		192	
Thr Phe Lys Gln Arg Arg Asp Ala Pro Pro Glu Gly Pro Ser Ala Lys			
50 55 60			
aaa ccg aag aag gag gcc tgc caa aaa cag aaa ggg gga ggc caa ggg		240	
Lys Pro Lys Lys Glu Ala Ser Gln Lys Lys Gln Lys Gly Gly Gln Gly			
65 70 75 80			
aag aag aag aag aac caa ggg aag aag aag gct aag aca ggg ccg cct		288	
Lys Lys Lys Lys Asn Gln Gly Lys Lys Lys Ala Lys Thr Gly Pro Pro			
85 90 95			
aat ccg aag gca cag aat gga aac aag aag aag acc aac aag aaa cca		336	
Asn Pro Lys Ala Gln Asn Gly Asn Lys Lys Lys Thr Asn Lys Lys Pro			
100 105 110			
ggc aag aga cag cgc atg gtc atg aaa ttg gaa tct gac aag acg ttc		384	
Gly Lys Arg Gln Arg Met Val Met Lys Leu Glu Ser Asp Lys Thr Phe			
115 120 125			
cca atc atg ttg gaa ggg aag ata aac ggc tac gct tgt gtg gtc gga		432	
Pro Ile Met Leu Glu Gly Lys Ile Asn Gly Tyr Ala Cys Val Val Gly			
130 135 140			
ggg aag tta ttc agg ccg atg cat gtg gaa ggc aag atc gac aac gac		480	
Gly Lys Leu Phe Arg Pro Met His Val Glu Gly Lys Ile Asp Asn Asp			
145 150 155 160			
gtt ctg gcc gcg ctt aag acg aag aaa gca tcc aaa tac gat ctt gag		528	
Val Leu Ala Ala Leu Lys Thr Lys Lys Ala Ser Lys Tyr Asp Leu Glu			
165 170 175			
tat gca gat gtg cca cag aac atg cgg gcc gat aca ttc aaa tac acc		576	
Tyr Ala Asp Val Pro Gln Asn Met Arg Ala Asp Thr Phe Lys Tyr Thr			
180 185 190			
cat gag aaa ccc caa gcc tat tac agc tgg cat cat gga gca gtc caa		624	
His Glu Lys Pro Gln Gly Tyr Tyr Ser Trp His His Gly Ala Val Gln			
195 200 205			
tat gaa aat ggg cgt ttc acg gtg ccg aaa gga gtt ggg gcc aag gga		672	
Tyr Glu Asn Gly Arg Phe Thr Val Pro Lys Gly Val Gly Ala Lys Gly			
210 215 220			

gac agc gga cga ccc att ctg gat aac cag gga cgg gtg gtc gct att 720
 Asp Ser Gly Arg Pro Ile Leu Asp Asn Gln Gly Arg Val Val Ala Ile
 225 230 235 240

gtg ctg gga ggt gtg aat gaa gga tct agg aca gcc ctt tca gtc gtc 768
 Val Leu Gly Gly Val Asn Glu Gly Ser Arg Thr Ala Leu Ser Val Val
 245 250 255

atg tgg aac gag aag gga gtt acc gtg aag tat act ccg gag aac tgc 816
 Met Trp Asn Glu Lys Gly Val Thr Val Lys Tyr Thr Pro Glu Asn Cys
 260 265 270

gag caa tgg tca cta gtg acc acc atg tgt ctg ctc gcc aat gtg acg 864
 Glu Gln Trp Ser Leu Val Thr Thr Met Cys Leu Leu Ala Asn Val Thr
 275 280 285

ttc cca tgt gct caa cca cca att tgc tac gac aga aaa cca gca gag 912
 Phe Pro Cys Ala Gln Pro Pro Ile Cys Tyr Asp Arg Lys Pro Ala Glu
 290 295 300

act ttg gcc atg ctc agc gtt aac atc cct gct ggg agg atc agc cgt 960
 Thr Leu Ala Met Leu Ser Val Asn Ile Pro Ala Gly Arg Ile Ser Arg
 305 310 315 320

aat tat tat aat tgg ctt ggt gct ggc tac tat tgt ggc cat gta cgt 1008
 Asn Tyr Tyr Asn Trp Leu Gly Ala Gly Tyr Tyr Cys Gly His Val Arg
 325 330 335

gct gac caa cca gaa aca 1026
 Ala Asp Gln Pro Glu Thr
 340

<210> 10

<211> 342

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 10

Met Phe Pro Phe Gln Pro Met Tyr Pro Met Gln Pro Met Pro Tyr Arg
 1 5 10 15
 Asn Pro Phe Ala Ala Pro Arg Arg Pro Trp Phe Pro Arg Thr Asp Pro
 20 25 30
 Phe Leu Ala Met Gln Val Gln Glu Leu Thr Arg Ser Met Ala Asn Leu
 35 40 45
 Thr Phe Lys Gln Arg Arg Asp Ala Pro Pro Glu Gly Gly Pro Ser Ala Lys
 50 55 60
 Lys Pro Lys Lys Glu Ala Ser Gln Lys Gln Lys Gly Gly Gly Gln Gly
 65 70 75 80

tgtctgcca	tgtgacgttc	ccatgtgctc	aaccaccaat	ttgctacgac	agaaaaccag	650
cagagacttt	ggccatgtctc	agcgttaacg	ttgacaaccc	ggcctacgat	gagctgctgg	720
aagcagctgt	taagtgtccc	ggaaggaaaa	ggagatccac	cgaggagctg	tttaaggatg	780
ataagctaac	ggccctttac	atggccagat	gcacatgatg	tgcagttggg	agctgccata	840
gtccaatgc	aatcgaggca	gtaaagagcg	acgggacaga	cggttatgtt	agacttcaga	900
cttctctgca	tatgggcttg	gattcctccg	gcaacttaaa	ggcgaggacc	atgcgggatg	960
acatgcacgc	gaccatataa	gagataccac	tacatcaagt	gtcaactccat	acacttgcct	1020
cggtgcacat	tgtggatggg	cacggttatt	tctcgtctgc	caggtgcccgc	caggcgagcc	1080
ccatcaccat	ggaatttaag	aaagattccg	tcacacactc	ctgctcgggt	ccgtatgaag	1140
tgaatttaa	tctctgaggg	agagaactct	atactcatcc	cccagaaacac	ggatagagc	1200
aagcgtgcc	agtctacgca	catgatgcac	agaacagagg	agcttatgtc	gagatgcacc	1260
tcccaggctc	agaagtggac	agcagtttgg	tttctcttag	cggcagttca	gtcacctgta	1320
cactctcctg	tgggactagc	gcccctgggtg	aatcgagtg	tggcggcaca	aagatctcca	1380
agaccataca	caagacaaaa	cagttcagcc	agtgacacaa	gaaggagcag	tgcagagcat	1440
atcggctgca	gaacgataag	tgggtgtata	atctcgacaa	actgcccaaa	gcagcgggag	1500
ccactctaaa	aggaataactg	catgtcccat	tcttctgtgc	agacggcaaa	tgcacccgtg	1560
ctctagcacc	agaaacctatg	ataacctctg	gtttcagatc	agtgctcactg	atgcgtgccc	1620
ctaagaatcc	cacataccta	accaccgcc	aacttctgta	tgagcctcac	tacacgcgatg	1680
agctcatatc	tgaaccagct	gttaggaatt	ttaccgtcac	cggaaaagggt	tgaggagtgt	1740
tatggggaaa	ccaccgccg	aaaaggtttt	gggacacaga	aacagcaccc	ggaaatccac	1800
atgggctacc	gcaagaggtg	ataactccat	attaccacag	ataccctatg	tccaccatcc	1860
tgggtttgtc	aatttgtgcc	gccattgcac	ccgtttccct	tgcagcgtct	acctgctgctg	1920
tttgcagatc	tagagttgog	tgcctaactc	cttaaccgct	aacacctaac	gctaggatag	1980
catcttctgt	ggctgtgctt	tgtcggccgc	gcactgccgc	ggcgagacac	acctgggagt	2040
cccttgatca	cctatggaaac	aataaaccaac	agatgtttctg	gattcaattg	ctgatccctc	2100
tggcgccttt	gtcgtagtg	actcgcctgc	tcagggtcgt	gtgctgtgtc	gtgccttttt	2160
tagtcatggc	cggcgccgca	ggcgccggcg	ccacagacac	cgcgaccacg	atgcggagcc	2220
aagcgggaat	ctcgtataac	actatagtca	acagagcagg	ctacgcacca	ctccctatca	2280
gcataaacac	aacaaagatc	aagctgatac	ctacagtga	cttggagtac	gtcaacctgc	2340
actacaaaa	aggaatggat	tcaccagcca	tcaaatgtctg	cggatctcag	gaatgcactc	2400
caacttacag	gctcgtatga	cagtgcacaa	tcttcacagg	gggtttaccg	ttcatgtggg	2460
gtggtgcata	ttgcttttgc	gacactgaga	acacccaagt	cagcaaggcc	tacgtaatga	2520
aatctgacga	ctgccttgcg	gatcatgctg	agcatatata	agcgacacac	gctccagtgc	2580
agggcttctc	caacatcaca	gtggggaac	actctattgt	gactaccgtg	tatgtgaagt	2640
ggaacacttc	tgtgaatttc	aatggggtca	aattaactgc	aggtccgctt	tccacagctt	2700
ggacaccctt	tgatcgcaaa	atcgtgcagt	atgcggggga	gatctataat	tatgattttc	2760
ctgagtatgg	ggcaggacaa	ccaggagcat	ttggagatag	accaatccag	acagcttcaa	2820
gctcagatct	gtatgccaat	accacactag	tgtcgcagat	accacaaagc	ggagcgtatc	2880
acgtgcacata	cactcagcca	ccttcgggtt	ttgagcaatg	gaagaaagat	aaagctccat	2940
cattgaaatt	taccgcccct	ttcggatgcy	aaatatatac	aaaccccatc	cgcgcgcgaa	3000
actgtactgt	agggctcaatt	ccattagcct	ttgacattcc	cgcgcgcttg	ttcaccaggg	3060
tgtcagaaac	accgcacactt	tcagcggccg	aatgcactct	tacagagtgc	gtgtatttct	3120
ccgacttttg	tgggactgcgc	acggtcaagt	actcggccag	caagtccagg	aagtgcgcag	3180
tccatgtgctc	atcaggagact	gctacctata	aagaagcagc	agtcagacta	accagacaa	3240
ggtcggcgac	tatccatttc	tgcaccgcaa	atatccaccc	ggagttcagg	ctccaaatat	3300
gcacatcata	tgttatcgtgc	aaaggtgtatt	gtcacccccc	gaagacacat	attgtgacac	3360
accctcagta	tacgcgccaa	acattttacag	cgcgggtgtc	aaaaaacccg	tggcgttggt	3420
taactcccat	gctcgggagga	tcagccgttaa	ttatttataat	tggcttggtg	ctggtcacta	3480
tttgggctct	gtacgtgctg	accacacaga	aacataatgt	aatacagcag	caatttgcga	3540
gctcgttaca	tagaactcgc	ggcgatttgc	atgcgctttt	aaaattttta	ttttattttt	3600
ctttctcttt	cgcgaatcgg	ttttgttttt	aatattttca	aaaaaaaaaa	aaaaaaaaaa	3660
aaaaaaaaaa	aaaaaaaaaa	aaaggaagaa	gcgcggcgcc	gcgcgtgggt	acgtcttctc	3720
ggcgttcggc	accgcaggct	ggatggcctt	cccattatg	attcttctcg	cttcggggg	3780
catcgggatg	ccgcggttgc	agcccatgct	gtccaggcag	gtagatgacg	acatcaggg	3840
acagctccaa	ggatcgtctg	cggctcttac	cagcctaact	tcatcactg	gaacctgat	3900

cgtaacggcg	atttatgcgc	cctcgccgag	cacatggaac	gggttgccat	ggattgtagg	3960
cgccgcccta	tacctgtct	gcctcccgc	gttggtgcgc	gggtgcacga	gcggggccac	4020
ctgcacccga	atgggaacgc	gcggcacctc	gttaacggat	tcacaccccc	aagaatttga	4080
gccaatcaat	tcttgccgag	aaactgtgaat	gcgcacaaca	acccttgcca	gaacatatcc	4140
atcgctccgc	ccatctccag	cagccgcagc	cggcgcatct	cgggcagcgt	tggttcctcg	4200
ccacgggtgc	gcattgatgt	gcctcctgcg	tggaggaccc	ggctaggctg	gcggggctgc	4260
cttactgggt	agcaaatga	atcacccgata	cgcgagcgaa	cgtgaagcga	ctgctgctgc	4320
aaacactctg	gcacctgagc	aacaacatga	atggtctctc	gtttcgtgtg	ttcgtaacct	4380
ctggaaacgc	gggaagtgcg	gcctcgacc	atgatgttcc	ggatctgcac	cgcaggatgc	4440
tgctggctac	cctgtgggac	acctacatct	gtattaacga	agcgctggca	ttgacccctga	4500
gtgatttttc	tctggtccgc	cgcgcatccat	accgccagt	gtttaacctc	acaaacctcc	4560
agtaacccga	catgttccat	atcagtaacc	cgtatogtga	gcactccttc	tcgtttcttc	4620
gggtatccgt	cccccatgaa	cagaaatccc	ccttacacgg	aggcatcagt	gaccaaacag	4680
gaaaaaacgc	cccttaacat	ggcccgcttt	atcagaagcc	agacattaac	gcttctggag	4740
aaactcaacg	agctggagcg	ggatgaacag	gcagacatct	gtgaatcgct	tcacgaccac	4800
ctgtgatgag	tttaccgcag	ctgcctcgcg	cgtttcgggtg	atgacgggtga	aaacctctga	4860
cacatgcagc	tcocggagac	ggtcacagct	tgctctgaag	cggatgcggg	gagcagacaa	4920
gcccgctcag	gcgcgtcagc	gggtggttgc	gggtgtcggg	gcgcagccat	gacccagctca	4980
cgtagcgata	cggaggtgta	tactggctta	actatgcggc	atcagagcag	gttgactctga	5040
gagtgaccca	tatatgcggt	gtgaaatacc	gcacagatgc	gtaaggagaa	aataccgcac	5100
caggcgctct	tcgccttccc	cgcctcactga	ctgcgtgcgc	tcggtgtctc	ggctgcggcg	5160
aggcgctatca	gctcaactca	aggcggtaat	accggttatcc	acagaatcac	gggataacgc	5220
aggaaagaa	atgtgagcaa	aaggccacga	aaaggccagg	aacogtaaaa	aggccgcgtt	5280
gctgcgcgtt	ttccataggc	tcgcgccccc	tgacgagcat	cacaaaaaatc	gacgtccaag	5340
tcagaggtgg	cgaaacccga	caggactata	aagataccag	gcggtttccc	ctggaaagctc	5400
cctcgtgcgc	ctctcgttcc	cgaccctgcc	gcctaccgga	tacctgtccg	cctttctccc	5460
ttcggggaag	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cgggtgaggt	5520
cgttgcgctcc	aagctgggct	gtgtgcacga	accccccggt	cagcccgacc	gctgcgcctt	5580
atccggtaac	tatcgtcttg	agtcacaacc	ggtaagacac	gacttatcgc	cactggccagc	5640
agccactgggt	aacaggatta	gcagagcgag	gtatgtaggc	gggtgctacg	agttcttgaa	5700
gtggtggcct	aactacggct	acactagaag	gacagtattt	ggatctcgcg	ctctctgtaa	5760
gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaaaaaa	ccacgcgctgg	5820
tagcgggtgg	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	5880
agatcctttg	atcttttcta	cggggtctga	cgtctcagtg	aacgaaaaatc	ccagtttaag	5940
gtattttgtc	atgaacaata	aaactgtctg	cttacataaa	cagtaataca	aggggtgttta	6000
tgagccatat	tcaacgggaa	acgtcttgct	cgaaggccgcg	attaaattcc	acaatggatg	6060
ctgatttata	tgggtataaa	tgggctcgog	ataatgtcgg	gcaatcaggt	gcgacaatct	6120
atcgatgtga	tgggaagccc	gagtgcgcag	agttgtttct	gaacatgcg	aaaggttagcg	6180
ttgccaatga	tgttcacagt	gagatggcta	gactaaactg	gctgacggaa	tttatgcctc	6240
ttccagacat	caagcatlct	atccgtactc	ctgatgatgc	atggttactc	accactcgga	6300
ttcccgggaa	acagacattc	caggtattag	aagaatatcc	tgattcaggt	gaaaaatttg	6360
tgcgtgcctg	ggcaggtgtc	ctgcgcgggt	tgcatcgaat	tcctgtttgt	aattgtccctt	6420
ttaaacagga	tcgcgtattt	cgctctcgct	aggcgcaatc	acgaatgaat	aacgggttgg	6480
ttgatgcgag	tgattttgat	gacgagcgta	atggctggcc	tggtgaacaa	gtctggaag	6540
aaatgcataa	ctttttgcca	ttctcacccg	atcagtcgt	cactcatggg	gatttctcac	6600
ttgatcaact	tatttttgac	gaggggaaat	taatagggtg	tattgatgtt	ggacgagtcg	6660
gaatcgcaga	cgcataccag	gatcttgcca	tcctatggaa	cgcctcgggt	gactttcttc	6720
cttcattaca	gaaacgggtt	tttcaaaaaa	atggattatga	taactctgat	atgaataaat	6780
tgcaagtttca	tttgatgctc	gatgagtttt	ttcaagaatt	ctcatgtttg	acagcttatc	6840
atcgataacg	tttaatgcgg	tagtttatca	cagttaaatt	gctaacgcag	tcagggaaccg	6900
tgtatgaaat	ctaacaatgc	gcctatcgtc	atcctcgcca	cgtcaacctc	ggatgctgtc	6960
tagaggatcc	ctaatacagc	tcactatag				6980

<210> 12
 <211> 2943
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; Note =
 synthetic construct

<221> CDS
 <222> (1)...(2943)

<400> 12
 atg tca cta gtg acc acc atg tgt ctg ctc gcc aat gtg acg ttc cca 48
 Met Ser Leu Val Thr Thr Met Cys Leu Leu Ala Asn Val Thr Phe Pro
 1 5 10 15
 tgt gct caa cca cca att tgc tac gac aga aaa cca gca gag act ttg 96
 Cys Ala Gln Pro Pro Ile Cys Tyr Asp Arg Lys Pro Ala Glu Thr Leu
 20 25 30
 gcc atg ctc agc gtt aac gtt gac aac cgg ggc tac gat gag ctg ctg 144
 Ala Met Leu Ser Val Asn Val Asp Asn Pro Gly Tyr Asp Glu Leu Leu
 35 40 45
 gaa gca gct gtt aag tgc ccc gga agg aaa agg aga tcc acc gag gag 192
 Glu Ala Ala Val Lys Cys Pro Gly Arg Lys Arg Arg Ser Thr Glu Glu
 50 55 60
 ctg ttt aag gag tat aag cta acg cgc cct tac atg gcc aga tgc atc 240
 Leu Phe Lys Glu Tyr Lys Leu Thr Arg Pro Tyr Met Ala Arg Cys Ile
 65 70 75 80
 aga tgt gca gtt ggg agc tgc cat agt coa ata gca atc gag gca gta 288
 Arg Cys Ala Val Gly Ser Cys His Ser Pro Ile Ala Ile Glu Ala Val
 85 90 95
 aag agc gac ggg cac gac ggt tat gtt aga ctt cag act tcc tcg cag 336
 Lys Ser Asp Gly His Asp Gly Tyr Val Arg Leu Gln Thr Ser Ser Gln
 100 105 110
 tat ggc ctg gat tcc tcc ggc aac tta aag ggc agg acc atg cgg tat 384
 Tyr Gly Leu Asp Ser Ser Gly Asn Leu Lys Gly Arg Thr Met Arg Tyr
 115 120 125
 gac atg cac ggg acc att aaa gag ata cca cta cat caa gtg tca ctc 432
 Asp Met His Gly Thr Ile Lys Glu Ile Pro Leu His Gln Val Ser Leu
 130 135 140
 cat aca tct cgc cgg tgt cac att gtg gat ggg cac ggt tat ttc ctg 480
 His Thr Ser Arg Pro Cys His Ile Val Asp Gly His Gly Tyr Phe Leu
 145 150 155 160
 ctt gcc agg tgc cgg gca ggg gac tcc atc acc atg gaa ttt aag aaa 528
 Leu Ala Arg Cys Pro Ala Gly Asp Ser Ile Thr Met Glu Phe Lys Lys
 165 170 175

gat tcc gtc aca cac tcc tgc tgc gtg cgg tat gaa gtg aaa ttt aat	576
Asp Ser Val Thr His Ser Cys Ser Val Pro Tyr Glu Val Lys Phe Asn	
180 185 190	
cct gta ggc aga gaa ctc tat act cat ccc cca gaa cac gga gta gag	624
Pro Val Gly Arg Glu Leu Tyr Thr His Pro Pro Glu His Gly Val Glu	
195 200 205	
caa gcg tgc caa gtc tac gca cat gat gca cag aac aga gga gct tat	672
Gln Ala Cys Gln Val Tyr Ala His Asp Ala Gln Asn Arg Gly Ala Tyr	
210 215 220	
gtc gag atg cac ctc cca ggc tca gaa gtg gac agc agt ttg gtt tcc	720
Val Glu Met His Leu Pro Gly Ser Glu Val Asp Ser Ser Leu Val Ser	
225 230 235 240	
ttg agc ggc agt tca gtc acc gtg aca cct cct gtt ggg act agc gcc	768
Leu Ser Gly Ser Ser Val Thr Val Thr Pro Pro Val Gly Thr Ser Ala	
245 250 255	
ctg gtg gaa tgc gag tgt ggc ggc aca aag atc tcc aag acc atc aac	816
Leu Val Glu Cys Glu Cys Gly Gly Thr Lys Ile Ser Lys Thr Ile Asn	
260 265 270	
aag aca aaa cag ttc agc cag tgc aca aag aag gag cag tgc aga gca	864
Lys Thr Lys Gln Phe Ser Gln Cys Thr Lys Lys Glu Gln Cys Arg Ala	
275 280 285	
tat cgg ctg cag aac gat aag tgg gtg tat aat tct gac aaa ctg ccc	912
Tyr Arg Leu Gln Asn Asp Lys Trp Val Tyr Asn Ser Asp Lys Leu Pro	
290 295 300	
aaa gca gcg gga gcc acc tta aaa gga aaa ctg cat gtc cca ttc ttg	960
Lys Ala Ala Gly Ala Thr Leu Lys Gly Lys Leu His Val Pro Phe Leu	
305 310 315 320	
ctg gca gac ggc aaa tgc acc gtg cct cta gca cca gaa cct atg ata	1008
Leu Ala Asp Gly Lys Cys Thr Val Pro Leu Ala Pro Glu Pro Met Ile	
325 330 335	
acc ttc ggt ttc aga tca gtg tca ctg aaa ctg cac cct aag aat ccc	1056
Thr Phe Gly Phe Arg Ser Val Ser Leu Lys Leu His Pro Lys Asn Pro	
340 345 350	
aca tat cta acc acc cgc caa ctt gct gat gag cct cac tac acg cat	1104
Thr Tyr Leu Thr Thr Arg Gln Leu Ala Asp Glu Pro His Tyr Thr His	
355 360 365	
gag ctc ata tct gaa cca gct gtt agg aat ttt acc gtc acc gga aaa	1152
Glu Leu Ile Ser Glu Pro Ala Val Arg Asn Phe Thr Val Thr Gly Lys	
370 375 380	
ggg tgg gag ttt gta tgg gga aac cac cgg cgg aaa agg ttt tgg gca	1200
Gly Trp Glu Phe Val Trp Gly Asn His Pro Pro Lys Arg Phe Trp Ala	
385 390 395 400	

cag gaa aca gca ccc gga aat cca cat ggg cta ccg cac gag gtg ata Gln Glu Thr Ala Pro Gly Asn Pro His Gly Leu Pro His Glu Val Ile 405 410 415	1248
act cat tat tac cac aga tac cct atg tcc acc atc ctg ggt ttg tca Thr His Tyr Tyr His Arg Tyr Pro Met Ser Thr Ile Leu Gly Leu Ser 420 425 430	1296
att tgt gcc gcc att gca acc gtt tcc gtt gca gcg tct acc tgg ctg Ile Cys Ala Ala Ile Ala Thr Val Ser Val Ala Ala Ser Thr Trp Leu 435 440 445	1344
ttt tgc aga tct aga gtt gog tgc cta act cct tac cgg cta aca cct Phe Cys Arg Ser Arg Val Ala Cys Leu Thr Pro Tyr Arg Leu Thr Pro 450 455 460	1392
aac gct agg ata cca ttt tgt ctg gct gtg ctt tgc tgc gcc cgc act Asn Ala Arg Ile Pro Phe Cys Leu Ala Val Leu Cys Cys Ala Arg Thr 465 470 475 480	1440
gcc cgg gcc gag acc acc tgg gag tcc ttg gat cac cta tgg aac aat Ala Arg Ala Glu Thr Thr Trp Glu Ser Leu Asp His Leu Trp Asn Asn 485 490 495	1488
aac caa cag atg ttc tgg att caa ttg ctg atc cct ctg gcc gcc ttg Asn Gln Gln Met Phe Trp Ile Gln Leu Leu Ile Pro Leu Ala Ala Leu 500 505 510	1536
atc gta gtg act cgc ctg ctc agg tgc gtg tgc tgt gtc gtg cct ttt Ile Val Val Thr Arg Leu Leu Arg Cys Val Cys Cys Val Val Pro Phe 515 520 525	1584
tta gtc atg gcc ggc gcc gca ggc gcc ggc gcc tac gag cac gcg acc Leu Val Met Ala Gly Ala Ala Gly Ala Gly Ala Tyr Glu His Ala Thr 530 535 540	1632
acg atg ccg agc caa gcg gga atc tcg tat aac act ata gtc aac aga Thr Met Pro Ser Gln Ala Gly Ile Ser Tyr Asn Thr Ile Val Asn Arg 545 550 555 560	1680
gca ggc tac gca cca ctc cct atc agc ata aca cca aca aag atc aag Ala Gly Tyr Ala Pro Leu Pro Ile Ser Ile Thr Pro Thr Lys Ile Lys 565 570 575	1728
ctg ata cct aca gtg aac ttg gag tac gtc acc tgc cac tac aaa aca Leu Ile Pro Thr Val Asn Leu Glu Tyr Val Thr Cys His Tyr Lys Thr 580 585 590	1776
gga atg gat tca cca gcc atc aaa tgc tgc gga tct cag gaa tgc act Gly Met Asp Ser Pro Ala Ile Lys Cys Cys Gly Ser Gln Glu Cys Thr 595 600 605	1824
cca act tac agg cct gat gaa cag tgc aaa gtc ttc aca ggg gtt tac Pro Thr Tyr Arg Pro Asp Glu Gln Cys Lys Val Phe Thr Gly Val Tyr 610 615 620	1872

ccg ttc atg tgg ggt ggt gca tat tgc ttt tgc gac act gag aac acc Pro Phe Met Trp Gly Gly Ala Tyr Cys Phe Cys Asp Thr Glu Asn Thr 625 630 635 640	1920
caa gtc agc aag gcc tac gta atg aaa tct gac gac tgc ctt gcg gat Gln Val Ser Lys Ala Tyr Val Met Lys Ser Asp Asp Cys Leu Ala Asp 645 650 655	1968
cat gct gaa gca tat aaa gcg cac aca gcc tca gtg cag gcg ttc ctc His Ala Glu Ala Tyr Lys Ala His Thr Ala Ser Val Gln Ala Phe Leu 660 665 670	2016
aac atc aca gtg gga gaa cac tct att gtg act acc gtg tat gtg aat Asn Ile Thr Val Gly Glu His Ser Ile Val Thr Thr Val Tyr Val Asn 675 680 685	2064
gga gaa act cct gtg aat ttc aat ggg gtc aaa tta act gca ggt ccg Gly Glu Thr Pro Val Asn Phe Asn Gly Val Lys Leu Thr Ala Gly Pro 690 695 700	2112
ctt tcc aca gct tgg aca ccc ttt gat cgc aaa atc gtg cag tat gcc Leu Ser Thr Ala Trp Thr Pro Phe Asp Arg Lys Ile Val Gln Tyr Ala 705 710 715 720	2160
ggg gag atc tat aat tat gat ttt cct gag tat ggg gca gga caa cca Gly Glu Ile Tyr Asn Tyr Asp Phe Pro Glu Tyr Gly Ala Gly Gln Pro 725 730 735	2208
gga gca ttt gga gat ata caa tcc aga aca gtc tca agc tca gat ctg Gly Ala Phe Gly Asp Ile Gln Ser Arg Thr Val Ser Ser Ser Asp Leu 740 745 750	2256
tat gcc aat acc aac cta gtg ctg cag aga ccc aaa gca gga gcg atc Tyr Ala Asn Thr Asn Leu Val Leu Gln Arg Pro Lys Ala Gly Ala Ile 755 760 765	2304
cac gtg cca tac act cag gca cct tgg ggt ttt gag caa tgg aag aaa His Val Pro Tyr Thr Gln Ala Pro Ser Gly Phe Glu Gln Trp Lys Lys 770 775 780	2352
gat aaa gct cca tca ttg aaa ttt acc gcc cct ttc gga tgc gaa ata Asp Lys Ala Pro Ser Leu Lys Phe Thr Ala Pro Phe Gly Cys Glu Ile 785 790 795 800	2400
tat aca aac ccc att cgc gcc gaa aac tgt act gta ggg tca att cca Tyr Thr Asn Pro Ile Arg Ala Glu Asn Cys Thr Val Gly Ser Ile Pro 805 810 815	2448
tta gcc ttt gac att ccc gac gcc ttg ttc acc agg gtg tca gaa aca Leu Ala Phe Asp Ile Pro Asp Ala Leu Phe Thr Arg Val Ser Glu Thr 820 825 830	2496
ccg aca ctt tca gcg gcc gaa tgc act ctt aac gag tgc gtg tat tct Pro Thr Leu Ser Ala Ala Glu Cys Thr Leu Asn Glu Cys Val Tyr Ser 835 840 845	2544

tcc gac ttt ggt ggg atc gcc acg gtc aag tac tcg gcc agc aag tca 2592
 Ser Asp Phe Gly Gly Ile Ala Thr Val Lys Tyr Ser Ala Ser Lys Ser
 850 855 860

ggc aag tgc gca gtc cat gtg cca tca ggg act gct acc cta aaa gaa 2640
 Gly Lys Cys Ala Val His Val Pro Ser Gly Thr Ala Thr Leu Lys Glu
 865 870 875 880

gca gca gtc gag cta acc gag caa ggg tcg gcg act atc cat ttc tcg 2688
 Ala Ala Val Glu Leu Thr Glu Gln Gly Ser Ala Thr Ile His Phe Ser
 885 890 895

acc gca aat atc cac ccg gag ttc agg ctc caa ata tgc aca tca tat 2736
 Thr Ala Asn Ile His Pro Glu Phe Arg Leu Gln Ile Cys Thr Ser Tyr
 900 905 910

gtt acg tgc aaa ggt gat tgt cac ccc ccg aaa gac cat att gtg aca 2784
 Val Thr Cys Lys Gly Asp Cys His Pro Pro Lys Asp His Ile Val Thr
 915 920 925

cac cct cag tat cac gcc caa aca ttt aca gcc gcg gtg tca aaa acc 2832
 His Pro Gln Tyr His Ala Gln Thr Phe Thr Ala Ala Val Ser Lys Thr
 930 935 940

gcg tgg acg tgg tta aca tcc ctg ctg gga gga tca gcc gta att att 2880
 Ala Trp Thr Trp Leu Thr Ser Leu Leu Gly Gly Ser Ala Val Ile Ile
 945 950 955 960

ata att ggc ttg gtg ctg gct act att gtg gcc atg tac gtg ctg acc 2928
 Ile Ile Gly Leu Val Leu Ala Thr Ile Val Ala Met Tyr Val Leu Thr
 965 970 975

aac cag aaa cat aat 2943
 Asn Gln Lys His Asn
 980

<210> 13

<211> 981

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<400> 13

Met Ser Leu Val Thr Thr Met Cys Leu Leu Ala Asn Val Thr Phe Pro
 1 5 10 15
 Cys Ala Gln Pro Pro Ile Cys Tyr Asp Arg Lys Pro Ala Glu Thr Leu
 20 25 30
 Ala Met Leu Ser Val Asn Val Asp Asn Pro Gly Tyr Asp Glu Leu Leu
 35 40 45

Glu Ala Ala Val Lys Cys Pro Gly Arg Lys Arg Arg Ser Thr Glu Glu
 50 55 60
 Leu Phe Lys Glu Tyr Lys Leu Thr Arg Pro Tyr Met Ala Arg Cys Ile
 65 70 75 80
 Arg Cys Ala Val Gly Ser Cys His Ser Pro Ile Ala Ile Glu Ala Val
 85 90 95
 Lys Ser Asp Gly His Asp Gly Tyr Val Arg Leu Gln Thr Ser Ser Gln
 100 105 110
 Tyr Gly Leu Asp Ser Ser Gly Asn Leu Lys Gly Arg Thr Met Arg Tyr
 115 120 125
 Asp Met His Gly Thr Ile Lys Glu Ile Pro Leu His Gln Val Ser Leu
 130 135 140
 His Thr Ser Arg Pro Cys His Ile Val Asp Gly His Gly Tyr Phe Leu
 145 150 155 160
 Leu Ala Arg Cys Pro Ala Gly Asp Ser Ile Thr Met Glu Phe Lys Lys
 165 170 175
 Asp Ser Val Thr His Ser Cys Ser Val Pro Tyr Glu Val Lys Phe Asn
 180 185 190
 Pro Val Gly Arg Glu Leu Tyr Thr His Pro Pro Glu His Gly Val Glu
 195 200 205
 Gln Ala Cys Gln Val Tyr Ala His Asp Ala Gln Asn Arg Gly Ala Tyr
 210 215 220
 Val Glu Met His Leu Pro Gly Ser Glu Val Asp Ser Ser Leu Val Ser
 225 230 235 240
 Leu Ser Gly Ser Ser Val Thr Val Thr Pro Pro Val Gly Thr Ser Ala
 245 250 255
 Leu Val Glu Cys Glu Cys Gly Gly Thr Lys Ile Ser Lys Thr Ile Asn
 260 265 270
 Lys Thr Lys Gln Phe Ser Gln Cys Thr Lys Lys Glu Gln Cys Arg Ala
 275 280 285
 Tyr Arg Leu Gln Asn Asp Lys Trp Val Tyr Asn Ser Asp Lys Leu Pro
 290 295 300
 Lys Ala Ala Gly Ala Thr Leu Lys Gly Lys Leu His Val Pro Phe Leu
 305 310 315 320
 Leu Ala Asp Gly Lys Cys Thr Val Pro Leu Ala Pro Glu Pro Met Ile
 325 330 335
 Thr Phe Gly Phe Arg Ser Val Ser Leu Lys Leu His Pro Lys Asn Pro
 340 345 350
 Thr Tyr Leu Thr Thr Arg Gln Leu Ala Asp Glu Pro His Tyr Thr His
 355 360 365
 Glu Leu Ile Ser Glu Pro Ala Val Arg Asn Phe Thr Val Thr Gly Lys
 370 375 380
 Gly Trp Glu Phe Val Trp Gly Asn His Pro Pro Lys Arg Phe Trp Ala
 385 390 395 400
 Gln Glu Thr Ala Pro Gly Asn Pro His Gly Leu Pro His Glu Val Ile
 405 410 415
 Thr His Tyr Tyr His Arg Tyr Pro Met Ser Thr Ile Leu Gly Leu Ser
 420 425 430
 Ile Cys Ala Ala Ile Ala Thr Val Ser Val Ala Ala Ser Thr Trp Leu
 435 440 445
 Phe Cys Arg Ser Arg Val Ala Cys Leu Thr Pro Tyr Arg Leu Thr Pro
 450 455 460
 Asn Ala Arg Ile Pro Phe Cys Leu Ala Val Leu Cys Cys Ala Arg Thr
 465 470 475 480

Ala Arg Ala Glu Thr Thr Trp Glu Ser Leu Asp His Leu Trp Asn Asn
 485 490 495
 Asn Gln Gln Met Phe Trp Ile Gln Leu Leu Ile Pro Leu Ala Ala Leu
 500 505 510
 Ile Val Val Thr Arg Leu Leu Arg Cys Val Cys Cys Val Pro Phe
 515 520 525
 Leu Val Met Ala Gly Ala Ala Gly Ala Tyr Glu His Ala Thr
 530 535 540
 Thr Met Pro Ser Gln Ala Gly Ile Ser Tyr Asn Thr Ile Val Asn Arg
 545 550 555 560
 Ala Gly Tyr Ala Pro Leu Pro Ile Ser Ile Thr Pro Thr Lys Ile Lys
 565 570 575
 Leu Ile Pro Thr Val Asn Leu Glu Tyr Val Thr Cys His Tyr Lys Thr
 580 585 590
 Gly Met Asp Ser Pro Ala Ile Lys Cys Cys Gly Ser Gln Glu Cys Thr
 595 600 605
 Pro Thr Tyr Arg Pro Asp Glu Gln Cys Lys Val Phe Thr Gly Val Tyr
 610 615 620
 Pro Phe Met Trp Gly Gly Ala Tyr Cys Phe Cys Asp Thr Glu Asn Thr
 625 630 635 640
 Gln Val Ser Lys Ala Tyr Val Met Lys Ser Asp Asp Cys Leu Ala Asp
 645 650 655
 His Ala Glu Ala Tyr Lys Ala His Thr Ala Ser Val Gln Ala Phe Leu
 660 665 670
 Asn Ile Thr Val Gly Glu His Ser Ile Val Thr Thr Val Tyr Val Asn
 675 680 685
 Gly Glu Thr Pro Val Asn Phe Asn Gly Val Lys Leu Thr Ala Gly Pro
 690 695 700
 Leu Ser Thr Ala Trp Thr Pro Phe Asp Arg Lys Ile Val Gln Tyr Ala
 705 710 715 720
 Gly Glu Ile Tyr Asn Tyr Asp Phe Pro Glu Tyr Gly Ala Gly Gln Pro
 725 730 735
 Gly Ala Phe Gly Asp Ile Gln Ser Arg Thr Val Ser Ser Ser Asp Leu
 740 745 750
 Tyr Ala Asn Thr Asn Leu Val Leu Gln Arg Pro Lys Ala Gly Ala Ile
 755 760 765
 His Val Pro Tyr Thr Gln Ala Pro Ser Gly Phe Glu Gln Trp Lys Lys
 770 775 780
 Asp Lys Ala Pro Ser Leu Lys Phe Thr Ala Pro Phe Gly Cys Glu Ile
 785 790 795 800
 Tyr Thr Asn Pro Ile Arg Ala Glu Asn Cys Thr Val Gly Ser Ile Pro
 805 810 815
 Leu Ala Phe Asp Ile Pro Asp Ala Leu Phe Thr Arg Val Ser Glu Thr
 820 825 830
 Pro Thr Leu Ser Ala Ala Glu Cys Thr Leu Asn Glu Cys Val Tyr Ser
 835 840 845
 Ser Asp Phe Gly Gly Ile Ala Thr Val Lys Tyr Ser Ala Ser Lys Ser
 850 855 860
 Gly Lys Cys Ala Val His Val Pro Ser Gly Thr Ala Thr Leu Lys Glu
 865 870 875 880
 Ala Ala Val Glu Leu Thr Glu Gln Gly Ser Ala Thr Ile His Phe Ser
 885 890 895
 Thr Ala Asn Ile His Pro Glu Phe Arg Leu Gln Ile Cys Thr Ser Tyr
 900 905 910

Val Thr Cys Lys Gly Asp Cys His Pro Pro Lys Asp His Ile Val Thr
 915 920 925
 His Pro Gln Tyr His Ala Gln Thr Phe Thr Ala Ala Val Ser Lys Thr
 930 935 940
 Ala Trp Thr Trp Leu Thr Ser Leu Leu Gly Gly Ser Ala Val Ile Ile
 945 950 955 960
 Ile Ile Gly Leu Val Leu Ala Thr Ile Val Ala Met Tyr Val Leu Thr
 965 970 975
 Asn Gln Lys His Asn
 980

<210> 14

<211> 12379

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 14

```

atggggcggcg catgagagaa gccagaccaa attacctacc caaaatggag aaagtccacg      60
ttgacatcga ggaagacagc ccatctctca gagctttgca gccgagcttc ccgcagtttg      120
aggtagaagc caagcaggtc actgataatg accatgctaa tgcagagcg ttttcgcatc      180
tggtctcaaa actgattcga acggaggtgg accatccga cagcatcctt gacattggaa      240
gtgcgcccg cgcagacaatg tattctaagc acaagtatca ttgatctgt ccgatgagat      300
gtcgggaaga tcggagacga ttgtataagt atgcaactaa gctgaagaaa aactgtaaag      360
aaataactga taagggaattg gacaagaaaa tgaaggagct cgcgcgcgtc atgagcgacc      420
ctgacctgga aactgagact atgtgcctcc acgacgacga gtcgtgtcgc tacgaaggcg      480
aagtgcgtgt ttaccaggat gtatacgcgg ttgacggacc gacaagtcct tatcaccaag      540
ccaataaggg agtttagagtc gcctactgga taggctttga caccacccct tttatgttta      600
agaacttgcc tggagcatat ccatcactat ctaccaactg gccgacgaa accgtgttaa      660
cggtcgttaa cataggccta tgcagctctg acgttatgga gcggtcacgt agagggatgt      720
ccattcttag aaagaagtat ttgaaacct ccaacaatgt tctattctct gttggctcga      780
ccatctacca cgagaagagg gacttactga ggagctggca cctgcgctct gtatttcaact      840
tacgtggcaa gcaaaattac acatgtcggg gtgagactat agttagttgc gacgggtacg      900
tcgttaaaag aatagctatc agtcaggccc tgtatgggaa gccctcagcg tatgctgcta      960
cgatgcaccc cyaggggattc ttgtgctgca aagtgcagaa cacattcaac ggggagaggg      1020
tctcttttcc cgtgtgcacg tatgtgccag ctacattgtg tgaccaaatg actggcatac      1080
tggcacagaa gttcagtcgc gacgacgcgc aaaaactgct gtttgggctc aaccagccta      1140
tagtcgtcaa cgttcgcacc cagagaacaa ccaataccat gaaaaattac cttttgcccg      1200
tagtggccca ggcatttgct aggtggggcaa aggaataaa ggaagatcaa gaagatgaaa      1260
ggccactagg actacgagat agacagttag tcatgggggt ttgttgggctc tttagaaggc      1320
acaagataac atctatttat aagcgcccg ataccacaa catcatcaaa gtgaacagcg      1380
atttcacctc attcgtgctg ccaggatag gcagtataac attggagatc gggctgagaa      1440
caagaatcag gaaaatgtta gaggagcaca aggagccgct acctctcatc accgcggagg      1500
acgtacaaga gctcaagtcg gcagcgcgat agcgtaaaga ggtcgtgaa gccgagaggt      1560
tgcgcgcagc tctaccacct ttggcagctg atgttgagga gccactctgt gaagccgatg      1620
tcgacttgat gttacaagag gctggggcgc gctcagtgga gacacctcgt ggcttgataa      1680
aggttaccag ctacgatggc gaggacaaga tcggtctcta cgtgtgctt tctcgcaggg      1740
ctgtactcaa gagtgaaaaa ttatcttgca tccacctctc cgtgaacaaa gtcatagtga      1800
taacacacct tggccgaaaa ggcgtttatg cgtggaacc ataccatggt aaagtatgtg      1860
tgccagaggg acatgcaata ccgtccagg accttcaagc ctgagtgaa agtgccacca      1920
ttgtgtacaa cgaactgtag ttcgtaaaca ggtacctgca ccatattgcc acacatggag      1980
gagcgctgaa cactgatgaa gaattattca aaactgtcaa gccagcgag cagcagcgcg      2040

```

aatacctgtga	cgacatcgac	aggaacacagt	gcgtcaagaa	agaactagtc	actgggctag	2100
ggctocacagg	cgagctgggt	gatcctccct	tcocatgaatt	cgccctacag	agtcctgagaa	2160
ccagcocaagc	gcgtcccttac	caagatcccaa	ccatagggggt	gtatggcggt	ccaggatcag	2220
gcaagctctgg	catcattaaa	agcgcagtc	ccaaaaaaga	tctagtgggt	agcgccaaga	2280
aagaaaaactg	tgcagaatt	ataagggagc	tcaagaaaa	gaaaggggtg	gacgtcaattg	2340
ccagaaactgt	ccagctcagtg	ctcttgaatg	gatgcaaaa	cccgctagag	accctgtata	2400
ttcagcgaagc	ttttgctgt	catgcaggt	ctctcagagc	gctcatagcc	atataaagac	2460
ctaaaaaactg	agtgctctgc	ggggatccca	aacagtgagg	tttttttaac	atgattgtgcc	2520
tgaagattgca	ttttaaccac	gagatttgca	cacaagtcct	ccacaaaagc	atctctcgcc	2580
gttgcactaa	atctgtgact	tcggtcgtct	caaccttggt	ttacgacaaa	aaaatgagaa	2640
cgaacgaatcc	gaaagagact	aagatttgta	ttgacactac	cggcagtaac	aaacctaaagc	2700
aggaagcatct	cattctcact	tggttcagag	gggtgggtgaa	gcagttgcaa	atagattaca	2760
aaggcacaoca	aataatgaag	gcagctgcct	ctcaaggggt	gaccogtaaa	ggtgtgtatg	2820
ccgttcogtta	caaggtgaat	gaaaatccct	tgtaagcaac	cacctcagaa	catgtgaaog	2880
tctactgac	ccgcacggag	gacgcagtc	tggtgaaaaa	actagcggcg	gacccatgga	2940
taaaaacact	gaactccgaag	tacccctggga	atttcaactgc	cacgatagag	gagtggaag	3000
cagcagcatga	tgcacatctg	agggacacat	tggaagagac	ggacccctac	cagctgtctcc	3060
agaaataagcg	aaaactgtgt	tggggcaagg	cttttagtgcc	ggtgctgaag	acccgtggca	3120
tagacatgac	catctgaaca	tggaaacactg	tgagattattt	tgaacggcag	aaagctccact	3180
cagcagagat	agttattgaac	caactatgag	tgaggttctt	tggactcgat	ctggactccg	3240
gtctattttt	tgcacccact	gttcogttat	ccattagga	taactcactg	gataaactcc	3300
cgctgcctcaa	catgtacggg	ctgaataaag	aagtggtccg	tcagctctct	cccgatgtacc	3360
cacaactggcc	tcggggcagtt	gccactggaa	gagttctatga	catgaacact	ggtaacactgc	3420
gcaattataga	tcocgcgata	aacctagtac	ctgtaaacag	aagactgcct	catgctttag	3480
tctcccaacca	taattgaacac	ccacagagtg	acttttcttc	attcgtcagc	aaatttgaag	3540
gcagaaactgt	ctcgtgtggtc	ggggaaaagt	gtccgtccc	aggcaaaatg	gttgaactgg	3600
tgtcagacccg	gcctgaggct	accttcagag	ctcggctgga	tttaggcaac	ccaggtgatg	3660
tgcccaataa	tgacataata	tttgttaag	tgaggacccc	atataaaac	catcactatc	3720
agcaagtgtga	agaccatgoc	attaagctta	gcattgtgac	caagaaagct	tgcttcgctc	3780
tgaatccccc	cggaacactgt	gtcagcatag	gttatggta	cgtcagacag	gcacggcaaa	3840
gcactcattgg	tgctatagcg	cggcagttca	agtttcccg	ggatgcaaa	ccgaatctct	3900
caactgaaga	gacgggaagt	ctgtttgtat	tcattgggta	cgatcgcaag	gcccgtacgc	3960
acaaatcctta	caagctttca	tcaacactga	ccaacattta	tacaggtcc	agactccaag	4020
aagcgggatg	tgcaccccta	tatcatgtgg	tgcgggggga	tattggcaag	gccacccgaag	4080
gagtgattat	aaatgctgct	aacagcaaa	gacaacctgg	cggagggggt	tgccgagcgc	4140
tgtataagaa	attcccgga	agcttogatt	tacagccgat	cgaagttaga	aaagccgcag	4200
tggctcaaaag	tgcagctaaa	catatcattc	atgcgtagg	acaaaacttc	acaaaagttt	4260
cggaaggttga	aggtgacaaa	cagttggcag	agggcttatga	tgccatgtgc	aagattgtca	4320
acgataacaa	ttcaagatca	gtagcgattc	caactgtgtc	cacgggcact	ttttccggga	4380
acaaagatgc	actaaaccca	tcattgaacc	atttgctgac	agcctttagac	accactgatg	4440
cagatgtagc	catatactgc	agggacaaga	aatgggaat	gactctcaag	gaagcagttg	4500
ctaggagaga	agcagtgga	gagatatgca	tatccgaag	actcttcaag	acagaaactg	4560
atgcagagct	ggtaggggtg	catccgaaga	gttctttggc	tgggaagga	ggctacagca	4620
caagcgatgg	caaaaacttc	tcataattgg	aagggaccaa	gtttccaccg	goggccaagg	4680
atatagcaga	aattaatgoc	tggtggcccg	tgtaacagcg	ggccaatgga	caggttatgca	4740
tgtatatctc	cgggaagaag	atgagcagta	ttaggtogaa	atgcoccgct	gaagagtggg	4800
aagccctctc	accacactagc	acgctgcctt	ctgtgtgcat	ccatgcgatg	actccagaaa	4860
gagtcacagc	ctcaaaagcc	tcacgtccag	aacaaattac	tggtgtctca	tcctttccat	4920
tgccgaagta	tagaattcact	gggtgcaaga	agatccaatg	ctccagcct	atatgtttct	4980
caccgaaagt	gccttgctat	attcatccaa	ggaagtatct	cgtgggaaca	ccacccgtag	5040
acagagactc	ggggccatcg	gcagagaaac	aatccacaga	gggaacact	gaacacagac	5100
cacttataac	cgaggatgag	accaggacta	gaacgcctga	gccgatcatc	atcgaaagag	5160
aagaagagga	tagcaataag	ttgctgtcag	atggcccgag	ccaccagggt	ctgcaagttg	5220
aggcagacat	tcaacggccg	ccctctgtat	ctagctatc	ctggtccatt	ctcatgcat	5280
ccgactttga	tgtggacagt	ttatccatag	ttgacaccc	ggaggagct	agcgtgacca	5340

gcggggcaac	gtcagccgag	actaactctt	acttcgcmaa	gagtatggag	ttcttgccgc	5400
gaccgcctgcc	tgcgcctcga	acagtatcca	ggaaccctcc	acatccccgt	ccgcgcacaa	5460
gaaccacccgt	acttcgaccc	agcagggcct	gtcgcgagaac	cagcttagtt	tcaccccgcg	5520
caggcgctgaa	taggggtgatc	actagagagg	agctcgaggc	gcttaccgcc	tcacgcactc	5580
ctagcaggtc	gggtctcgaga	accagcctgg	tctccaaccc	gccaggcgta	aataggsgtg	5640
ttacaagaga	ggaggtctbag	gcgttcgttag	cacaacaaca	atgacgggtt	gatgcgggtg	5700
catacatctt	ttctctccag	accggtcaag	ggcatttaca	acaaaaatca	gtaaaggcaa	5760
cgggtctalc	cgaagttctg	tgggagagga	cgcgaattga	gatttcgtat	gccccgcgcc	5820
tgaccacaaga	aaaagaagaa	ttactacgca	agaaattaca	gttaaatccc	acacctgcta	5880
acagaagcag	ataccagctc	aggaaggtgg	agaaatgaa	agccataaca	gctgacgta	5940
ttctgcgaag	cctagggcat	tatttgaagg	cagaaggaaa	agtggaagtgc	taccgaaccc	6000
tgcactctgt	tcctttgtat	tcacttagtg	tgaacccgtc	cttttcaagc	cccaagggtg	6060
cagtggaagc	ctgtaacgcc	atgttgaaag	agaactttcc	gactgtggct	tcttactgta	6120
ttatttcaga	gtacgatgcc	tatttggaca	tggttgacgg	agcttcacgc	tgcttagaca	6180
ctgcacgttt	ttgcctcgca	aagctgcgca	gctttccaaa	gaacactccc	tatttggaa	6240
ccacatacag	atcgccagtg	ccttcagcga	tcocagaacac	gctccagaac	gtcctggcag	6300
ctgcacaaaa	aagaatttgc	aatgtcacgc	aaatgagaga	attgcccgtg	ttggattcgg	6360
cggcctttaa	tgtggaatgc	ttcaagaaat	atgcgtgtaa	taatgaatat	tgggaaacgt	6420
ttaaagaaaa	cccatcagg	cttactgaag	aaaaactgtg	aaattacatt	accaaatcaa	6480
aaggacacaaa	agctgctgct	ctttttcgca	agacacataa	tttgaatatg	ttgcaggaca	6540
taccaattgga	caggtttgta	atggacttaa	agagagacgt	gaaagtgcac	ccagggaacaa	6600
aacatactga	agaaagggcc	aaggtacagg	tgatccaggc	tgccgatccg	ctagcaacag	6660
cgtatctgtg	cggaaatcac	cagagagctg	ttaggagatt	aaatgcggct	ctgcttcoga	6720
actatcatcat	actgtttgat	atgtcggctg	aagactttgc	cgctattata	gccagacact	6780
tcacgctgg	ggattgtggt	ctggaaactg	acatcgcgtc	gtttgataaa	agtgaggacg	6840
acgcctagcc	ttgcagccgc	ttaatgattc	tggaaagatt	aggtgtggac	gcagagctgt	6900
tgacgctgat	tggaggcgct	ttcggcgaaa	tttcatcaat	acattttgcc	actaaactta	6960
aattttaaatt	cggagccatg	atgaaatctg	gaatgttctc	cacactgttt	gtgaacacag	7020
tcattaacat	tgtaactcga	agcagagtg	tgagagaacg	gctaaacgga	tcaccatgtg	7080
cagcatctcat	tggagatgac	aatatcgtga	aaggagtcac	atccggaacaa	ttaatggcag	7140
acaggtgcgc	cacctgggtg	aatatggaag	tcaagattat	agatgctgtg	gtgggcgaga	7200
aagcgcctta	ttctctgga	gggtttattt	tgtgtgactc	cgtgaccggc	acagcgtgcc	7260
gtgtggcaga	cccccataaa	aggtctgtta	agcttggcaa	acctctggca	gcagcagatg	7320
aacatgatga	tgacaggaga	agggcatttc	atgaagagtc	aacacgcgtg	aaccgagtg	7380
gtattctttc	agagctgtgc	aaggcagtag	aatcaaggta	tgaacccgta	ggaacttcca	7440
tcatagttat	ggccatgact	actctagcta	gcagtggtta	atcattcagc	tacctgagag	7500
gggcgcctat	aaactctctac	ggctaaccctg	aatggactac	gacatagctc	agttccgcac	7560
tggtcccaatc	agtcctcattg	aaactgtacc	agtaaaactg	aagccaggaa	tggtgcggcc	7620
aaaggttata	caatggcgcgt	taacagaagt	gaaaaataaa	gcatttaaac	caattttgtg	7680
agaaattgga	aaggaaaggaa	aaattacaaa	aatggcgctc	gaaatcccat	ataactctcc	7740
aatattcgcc	ataaaaaagg	aagacagcac	taagtggaga	aaattagtag	atttcaggga	7800
acctcaataa	agaaactcaag	acttttggga	ggttcaatta	ggaataccac	accagcagc	7860
gttaaaaaag	aaaaaattcag	tgacagtagt	ggatgtggga	gagatcatat	tttcatctcc	7920
tttagatgaa	ggcttcaggga	aatatactgc	attcaccata	cctagataaa	acaaatgaa	7980
accagggatt	agatatcaat	ataatgtgct	tcocaaagga	tggaaagggt	caccagcaat	8040
attccagcgt	agcatgacaa	aaatcctaga	gccccttaga	gctaaaaatc	cagaaatagt	8100
catctatcaa	catatggcgg	cattgtatgt	aggaatcgac	ttagaaaatg	ggcaacatag	8160
agcaaaaaata	gaagagttaa	gagaacatct	attaaagggt	ggagttaaca	caccagacaa	8220
aaaaacatcag	aagaacaccc	catttctttg	gatgggtagt	gaactctatc	ctgacaaagt	8280
gacagctacag	cctatacagc	tgccagaaaa	agatagctgg	actgtcaatg	acatacagaa	8340
gttagtgga	aaattaaact	ggacaagtca	gatttaccga	gggataaag	taaggcaact	8400
ttgtaagctc	cttagggggg	ccaaagcact	aacagacata	gtaccactaa	ctgaagaagc	8460
agaattagaa	ttggcgagaga	acagggaaat	tctaaaaagaa	ccagtgcatg	gagtatatta	8520
tgaccctatca	aaagacttga	tagctgaaat	acagaaacac	ggggatgacc	aagggaacata	8580
tcaaatttac	caagaacact	tcaaaaacct	gaagacagga	aagatgcaa	aaaggaggac	8640

tacccacact	aatgatgtaa	aacagttaac	agaggcagtg	caaaaaaat	ccttggaaa	8700
catagtaaca	tggggaaaga	ctcctaata	tagactacct	atcccaaaa	aaacatggga	8760
aatatgggtg	acagactatt	ggcaagccac	atggattccc	gagtgggagt	ttgttaatac	8820
ccctccccta	gtaaaaactat	ggtaccagct	agaaaaagaa	cccatagcag	gagcagaaac	8880
tttctgaagg	cgggccttaa	ttaagtaacy	atacagcagc	aattggcag	ctgcttaaat	8940
agaactccgc	gcgatgggca	tgcgcgttta	aaatttttat	tttatttttc	ttttcttttc	9000
cgaatcggat	tttgttttta	atattttcaa	aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	9060
aaaaaataaa	aaagggaaga	gcgcggcgcc	gcgctgggct	acgtcttgct	ggcgttcggc	9120
acgcgaggtc	ggatggccct	cccattatgt	attcttctgc	cttcggcggc	catcgggatg	9180
ccgcgcttgc	agggcatgct	gtccaggcag	gtagatgacg	accatcaggg	acagcttcaa	9240
ggatcgctgc	cggtctttac	cagcctaact	togatcaact	gaccgctgat	cgtcaaggcg	9300
atttatgcgc	cctgcgcgag	cacatggaa	gggttggcag	ggattgtagg	cgccgcctta	9360
tacctgtctc	ctgcctccgc	gttgcgctgc	gggtcatgga	gcggggccac	ctcgacctga	9420
atgggaagcg	gcggcacctc	gctaaccgat	tcaccactcc	aagaattgga	gcacatcaat	9480
tcttgcgagg	aactgtgaat	gcgcaaacca	acocctggca	gaacatatcc	atcgcgctcg	9540
ccatctccag	cagccgcacg	cggcgcatct	cgggcaagct	tgggtcctgt	ccaaggggtg	9600
cgatgatcgt	gcctctctgc	ttgaggaccc	ggctaggctg	gcgggggtgc	cttactgggtt	9660
agcagaatga	atcaccgata	cgcgagcgaa	cgtgaagcga	ctgctgcctg	aaacagctct	9720
gcgcctgggc	aaacaacatga	atggtcttcg	gtttccgtgt	ttcgtaaaag	ctggaaaagc	9780
ggaagtccgc	gcctgcaccc	attatgttcc	ggatctgcag	cgcaggatgc	tgctggctac	9840
ctctgtggaa	acctacaact	gtattaaaga	agocgtggca	ttgacctgga	gtgatttttc	9900
tctgttccgc	cgcgctccat	accgcagbtt	gtttaccctc	acaaagcttc	cgtaaccggg	9960
catgttctac	atcagtaacc	cgtatcgtga	gcactcctct	tcgtttctac	ggatcattta	10020
cccccatgaa	cagaaatacc	ccctacacgc	agggatcagc	gacaaaacag	gaaaaaacgc	10080
cccttaacat	ggccgcgttt	atcagaagcc	agacattaac	gcttctggag	aaactcaagc	10140
agctggacgc	ggatgaaacg	gcagacatct	gtgaattcgt	tcacgaccac	gctgatgagc	10200
tttaccgcag	ctgcctcgcc	cgtttccgtg	atgacgggtg	aaacctctga	cacatgcagc	10260
tcocgcgagc	ggtaacagct	tgtctgtaag	cggatgcggc	gagcagacaa	gcccgtcagg	10320
gcgcgctcag	gggtgttgcc	gggtgtcgcc	gcgcagccat	gacccagtca	cgtagcgata	10380
gcggagtgtg	taactggctta	actatcgggc	atcagagcag	attgtactga	gagtgaccca	10440
tatatgcggg	gtgaataacc	gcacagatgc	gtaaggagaa	aataccgcac	cagcgctcct	10500
tcgcgtctct	cgcctcactga	ctcgctgcgc	tcggctcgtc	ggctgcggcg	agcggtatca	10560
gctcactcaa	agggcggtat	acggttatcc	acagaaatcag	gggataacgc	aggaagaagc	10620
atgtgcagaa	aaggccagca	aaaggccagg	aacctgaata	agggcgcgct	gctggcgctt	10680
ttccatcgct	tcgcgcccc	tgacgagcat	acaaaaatc	gacgctcaag	tcagaggtgg	10740
cgaaaccoga	caggatcata	aagataccag	gcgtttcccc	ctggaaagctc	cctcgctgcg	10800
ttctctgttc	cgacctgcgc	gcttaccgga	taactgtcgc	cccttctccc	ttcggaagc	10860
gtggcgcttg	ctcatagctc	acgctgtagg	tatctcagbt	cgggttaggt	cgttcgctcc	10920
aaactgggct	gtgtgcagca	acccccggtt	cagcccgacc	gctgcgcctt	atccggtaac	10980
tatcgtcttg	agctccaaac	ggtaagacac	gacttatcgc	cactggcagc	agccactgggt	11040
acaaggatta	gcagagcgag	gtatgtaggc	gggtctacag	agttcttgaa	gtggtggcct	11100
aactacggct	acactagaag	gacagtattt	ggatatctgc	ccctgctgaa	gcaggttaac	11160
ttcggaaaaa	gagtggtgat	ctcttgatcc	ggcaaaaaaa	ccaccgcgtg	tagcgggtgtg	11220
ttttttgttt	gcaagcagca	gattaaogcc	agaaaaaaag	gatctcaaga	agatcccttg	11280
atctttttta	gggggtctga	cgctcagtg	acgaaaaact	cagcttaagg	gattttggct	11340
atcaacaaata	aaactgtctg	cttacataaa	cagtaataca	aggggtgtta	tgagccatat	11400
tcacaggggaa	acgtcttgct	cgaggccgcg	atataattcc	aacatggatg	ctgatttata	11460
tggtgtataaa	tggtgcgcgc	ataatgtcgg	gcaatcaggt	gcgacaatct	atcgatttga	11520
tggggaagccc	gaggtgccag	agttgtttct	gaacataggg	aaaggttagc	ttgccaatga	11580
tgttacagat	gagatggtca	gactaaactg	gctgacggaa	tttatgcctc	ttccgacocat	11640
caagcatctt	atccgtactc	ctgatgatgc	agtttatact	accactgoga	tcoccgggaa	11700
aaacagcttc	caggtattag	aagaatatcc	tgattcaggt	gaaaataatt	ttgatgcgct	11760
ggcaggtgtc	ctcgcgcggg	tgcatctgat	tcctgtttgt	aattgtcctt	ttaacagcga	11820
tcogtatatt	cgctcgcctc	agcgccaatc	acgaatgaat	acaggtttgt	ttgatgcgag	11880
tgattttgat	gacgagcgta	atgctgtggc	tggtgaacaa	gtctggaaa	aaatgcataa	11940

```

gtttttgcca ttctcaccgg attcagtcgt cactcatggt gattttctcac ttgataaacct 12000
tatttttgac gaggggaaat taataggttg tattgatgtt ggacgagtcg gaatgcgaga 12060
ccgataccag gatcttgcca tctatggaa ctgcctcggg gagttttctc ctctattaca 12120
gaaacggctt tttcaaaaat atggtattga taatcctgat atgaataaat tgcagtttca 12180
tttgatgctc gatgagtttt tctaagaatt ctcatgttg acagcttacc atcgataagc 12240
tttaatgcgg tatgttatca cagttaaatt gctaacgcag tcaggcaccc tgatgaaat 12300
ctaacaatgc gctcatgctc atcctcgga cegtcaccct ggatgctgct tagaggatcc 12360
ctaatacgac tcactatag

```

<210> 15

<211> 1323

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<221> CDS

<222> (1)...(1323)

<400> 15

```

atg cca atc agt ccc att gaa act gta cca gta aaa ctg aag cca gga 48
Met Pro Ile Ser Pro Ile Glu Thr Val Pro Val Lys Leu Lys Pro Gly

```

```

1 5 10 15

```

```

atg gat ggc cca aag gtt aaa caa tgg cgg tta aca gaa gtg aaa ata 96
Met Asp Gly Pro Lys Val Lys Gln Trp Pro Leu Thr Glu Val Lys Ile

```

```

20 25 30

```

```

aaa gca tta aca gca att tgt gaa gaa atg gaa aag gaa gga aaa att 144
Lys Ala Leu Thr Ala Ile Cys Glu Glu Met Glu Lys Glu Gly Lys Ile

```

```

35 40 45

```

```

aca aaa att ggg cct gaa aat cca tat aac act cca ata ttc gcc ata 192
Thr Lys Ile Gly Pro Glu Asn Pro Tyr Asn Thr Pro Ile Phe Ala Ile

```

```

50 55 60

```

```

aaa aag gaa gac agc act aag tgg aga aaa tta gta gat ttc agg gaa 240
Lys Lys Glu Asp Ser Thr Lys Trp Arg Lys Leu Val Asp Phe Arg Glu

```

```

65 70 75 80

```

```

ctc aat aaa aga act caa gac ttt tgg gag gtt caa tta gga ata cca 288
Leu Asn Lys Arg Thr Gln Asp Phe Trp Glu Val Gln Leu Gly Ile Pro

```

```

85 90 95

```

```

cac cca gca ggg tta aaa aag aaa aaa tca gtg aca gta ctg gat gtg 336
His Pro Ala Gly Leu Lys Lys Lys Lys Ser Val Thr Val Leu Asp Val

```

```

100 105 110

```

```

gga gat gca tat ttt tca gtt cct tta gat gaa ggc ttc agg aaa tat 384
Gly Asp Ala Tyr Phe Ser Val Pro Leu Asp Glu Gly Phe Arg Lys Tyr

```

```

115 120 125

```

act gca ttc acc ata cct agt ata aac aat gaa aca cca ggg att aga Thr Ala Phe Thr Ile Pro Ser Ile Asn Asn Glu Thr Pro Gly Ile Arg 130 135 140	432
tat caa tat aat gtg ctt cca caa gga tgg aaa ggg tca cca gca ata Tyr Gln Thr Asn Val Leu Pro Gln Gly Trp Lys Gly Ser Pro Ala Ile 145 150 155 160	480
ttc cag gct agc atg aca aaa atc cta gag ccc ttt aga gct aaa aat Phe Gln Ala Ser Met Thr Lys Ile Leu Glu Pro Phe Arg Ala Lys Asn 165 170 175	528
cca gaa ata gtc atc tat caa cat atg gcg gca ttg tat gta gga tct Pro Glu Ile Val Ile Tyr Gln His Met Ala Ala Leu Tyr Val Gly Ser 180 185 190	576
gac tta gaa ata ggg caa cat aga gca aaa ata gaa gag tta aga gaa Asp Leu Glu Ile Gly Gln His Arg Ala Lys Ile Glu Glu Leu Arg Glu 195 200 205	624
cat cta tta aag tgg gga ttt acc aca cca gac aaa aaa cat cag aaa His Leu Leu Lys Trp Gly Phe Thr Thr Pro Asp Lys Lys His Gln Lys 210 215 220	672
gaa ccc cca ttt ctt tgg atg ggg tat gaa ctc cat cct gac aaa tgg Glu Pro Pro Phe Leu Trp Met Gly Tyr Glu Leu His Pro Asp Lys Trp 225 230 235 240	720
aca gta cag cct ata cag ctg cca gaa aaa gat agc tgg act gtc aat Thr Val Gln Pro Ile Gln Leu Pro Glu Lys Asp Ser Trp Thr Val Asn 245 250 255	768
gac ata cag aag tta gtg gga aaa tta aac tgg aca agt cag att tac Asp Ile Gln Lys Leu Val Gly Lys Leu Asn Trp Thr Ser Gln Ile Tyr 260 265 270	816
cca ggg att aaa gta agg caa ctt tgt aag ctc ctt agg ggg acc aaa Pro Gly Ile Lys Val Arg Gln Leu Cys Lys Leu Leu Arg Gly Thr Lys 275 280 285	864
gca cta aca gac ata gta cca cta act gaa gaa gca gaa tta gaa ttg Ala Leu Thr Asp Ile Val Pro Leu Thr Glu Glu Ala Glu Leu Glu Leu 290 295 300	912
gca gag aac agg gaa att cta aaa gaa cca gtg cat gga gta tat tat Ala Glu Asn Arg Glu Ile Leu Lys Glu Pro Val His Gly Val Tyr Tyr 305 310 315 320	960
gac cca tca aaa gac ttg ata gct gaa ata cag aaa cag ggg gat gac Asp Pro Ser Lys Asp Leu Ile Ala Glu Ile Gln Lys Gln Gly Asp Asp 325 330 335	1008
caa tgg aca tat caa att tac caa gaa cca ttc aaa aac ctg aag aca Gln Trp Thr Tyr Gln Ile Tyr Gln Glu Pro Phe Lys Asn Leu Lys Thr 340 345 350	1056

gga aag tat gca aaa agg agg act acc cac act aat gat gta aaa cag 1104
 Gly Lys Tyr Ala Lys Arg Arg Thr Thr His Thr Asn Asp Val Lys Gln
 355 360 365

tta aca gag gca gtg caa aaa ata tcc ttg gaa agc ata gta aca tgg 1152
 Leu Thr Glu Ala Val Gln Lys Ile Ser Leu Glu Ser Ile Val Thr Trp
 370 375 380

gga aag act cct aaa ttt aga cta ccc atc caa aaa gaa aca tgg gaa 1200
 Gly Lys Thr Pro Lys Phe Arg Leu Pro Ile Gln Lys Glu Thr Trp Glu
 385 390 395 400

ata tgg tgg aca gac tat tgg caa gcc aca tgg att cct gag tgg gag 1248
 Ile Trp Trp Thr Asp Tyr Trp Gln Ala Thr Trp Ile Pro Glu Trp Glu
 405 410 415

ttt gtt aat acc cct ccc cta gta aaa cta tgg tac cag cta gaa aaa 1296
 Phe Val Asn Thr Pro Pro Leu Val Lys Leu Trp Tyr Gln Leu Glu Lys
 420 425 430

gaa ccc ata gca gga gca gaa act ttc 1323
 Glu Pro Ile Ala Gly Ala Glu Thr Phe
 435 440

<210> 16

<211> 441

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 16

Met Pro Ile Ser Pro Ile Glu Thr Val Pro Val Lys Leu Lys Pro Gly
 1 5 10 15
 Met Asp Gly Pro Lys Val Lys Gln Trp Pro Leu Thr Glu Val Lys Ile
 20 25 30
 Lys Ala Leu Thr Ala Ile Cys Glu Glu Met Glu Lys Glu Gly Lys Ile
 35 40 45
 Thr Lys Ile Gly Pro Glu Asn Pro Tyr Asn Thr Pro Ile Phe Ala Ile
 50 55 60
 Lys Lys Glu Asp Ser Thr Lys Trp Arg Lys Leu Val Asp Phe Arg Glu
 65 70 75 80
 Leu Asn Lys Arg Thr Gln Asp Phe Trp Glu Val Gln Leu Gly Ile Pro
 85 90 95
 His Pro Ala Gly Leu Lys Lys Lys Lys Ser Val Thr Val Leu Asp Val
 100 105 110
 Gly Asp Ala Tyr Phe Ser Val Pro Leu Asp Glu Gly Phe Arg Lys Tyr
 115 120 125
 Thr Ala Phe Thr Ile Pro Ser Ile Asn Asn Glu Thr Pro Gly Ile Arg
 130 135 140

50

Tyr Gln Tyr Asn Val Leu Pro Gln Gly Trp Lys Gly Ser Pro Ala Ile
 145 150 155 160
 Phe Gln Ala Ser Met Thr Lys Ile Leu Glu Pro Phe Arg Ala Lys Asn
 165 170 175
 Pro Glu Ile Val Ile Tyr Gln His Met Ala Ala Leu Tyr Val Gly Ser
 180 185 190
 Asp Leu Glu Ile Gly Gln His Arg Ala Lys Ile Glu Glu Leu Arg Glu
 195 200 205
 His Leu Leu Lys Trp Gly Phe Thr Thr Pro Asp Lys Lys His Gln Lys
 210 215 220
 Glu Pro Pro Phe Leu Trp Met Gly Tyr Glu Leu His Pro Asp Lys Trp
 225 230 235 240
 Thr Val Gln Pro Ile Gln Leu Pro Glu Lys Asp Ser Trp Thr Val Asn
 245 250 255
 Asp Ile Gln Lys Leu Val Gly Lys Leu Asn Trp Thr Ser Gln Ile Tyr
 260 265 270
 Pro Gly Ile Lys Val Arg Gln Leu Cys Lys Leu Leu Arg Gly Thr Lys
 275 280 285
 Ala Leu Thr Asp Ile Val Pro Leu Thr Glu Glu Ala Glu Leu Glu Leu
 290 295 300
 Ala Glu Asn Arg Glu Ile Leu Lys Glu Pro Val His Gly Val Tyr Tyr
 305 310 315 320
 Asp Pro Ser Lys Asp Leu Ile Ala Glu Ile Gln Lys Gln Gly Asp Asp
 325 330 335
 Gln Trp Thr Tyr Gln Ile Tyr Gln Glu Pro Phe Lys Asn Leu Lys Thr
 340 345 350
 Gly Lys Tyr Ala Lys Arg Arg Thr Thr His Thr Asn Asp Val Lys Gln
 355 360 365
 Leu Thr Glu Ala Val Gln Lys Ile Ser Leu Glu Ser Ile Val Thr Trp
 370 375 380
 Gly Lys Thr Pro Lys Phe Arg Leu Pro Ile Gln Lys Glu Thr Trp Glu
 385 390 395 400
 Ile Trp Trp Thr Asp Tyr Trp Gln Ala Thr Trp Ile Pro Glu Trp Glu
 405 410 415
 Phe Val Asn Thr Pro Pro Leu Val Lys Leu Trp Tyr Gln Leu Glu Lys
 420 425 430
 Glu Pro Ile Ala Gly Ala Glu Thr Phe
 435 440

<210> 17

<211> 13584

<212> DNA

<213> Artificial Sequence

<220>

 <223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 17

atgggcgccg	catgagagaa	gccagacca	attacctacc	caaaatggag	aaagtccagc	60
ttgacatcga	ggaagacagc	ccattcctca	gagctttgca	gcggagcttc	ccgcagtttg	120
aggtagaagc	caagcaggctc	actgataatg	accatgctaa	tgccagagcg	ttttcgctac	180
tggtttcaaa	actgatcgaa	acggaggctg	acccatccga	cacgatccct	gacattggaa	240
gtgcgcgcgc	ccgcagaatg	tattctaagc	acaagtatca	ttgtatctgt	ccgatgagat	300
gtgcggaaga	tccggacaga	ttgtataagt	atgcaactaa	gctgaagaaa	aactgtaagg	360

aaataactga	taaggaattg	gacaagaaaa	tgaaggagct	cgccgcgcgtc	atgagcgcacc	420
ctgacctggg	aactgagact	atgtgcctcc	acgacgacga	gtcgtgtcgc	tacgaaggcgc	480
aagtgcctgt	ttaccaggat	gtatacgcgg	ttgacggacc	gacaagtctc	tatcaccaagc	540
ccaataaggc	agtttagatc	gcctactgga	taggctttga	caccaccctc	ctttagtctta	600
agaaactggc	tggagcatat	ccatcatact	ctaccaactg	ggccgcacga	accggtttaa	660
cggtctgtaa	cataggcccta	tgcagctctg	acgttatgga	ggcggtcacgt	agagggtagt	720
ccatctcttag	aaagaagtat	ttgaaacctt	ccaacaatgt	tctattctct	gttggtcgca	780
ccatctacga	cgagaagagg	gacttactga	ggagctggca	ctcgcgctct	gtatttccat	840
tacgtggcaa	gcaaaattac	acatgtcggg	gtgagactat	agtttagtgc	gacgggtacg	900
tcgtaaaag	aatagctatc	agtcacggcc	tgtatgggaa	gccttcaggc	tatgtcgtcta	960
cgatgcaccg	cgaggggattc	ttgtgtctga	aagtgcacga	cacattcaac	ggggagcagg	1020
tctcttttcc	cgtgtgcacg	tatgtgccag	ctacattgtg	tgaccaaatg	actggcattc	1080
tggccaacaga	tgtcagtgcg	gacgaogcgc	aaaaaactgt	ggttgggctc	aaccagcgta	1140
tgtctgtcca	cggtcgcacc	cagagaacaa	ccaataccat	gaaaaattac	cttttgcgcc	1200
tagttggccca	ggcatttgct	aggtggggcaa	aggaataata	ggaagatcaa	gaagatgaaa	1260
ggccactagg	actacgagat	agacagttag	tcatgggggtg	ttgttgggct	tttagaaggc	1320
aaaagataac	actcatcttat	aagcgcccg	ataccacaac	catcatcaaa	gtggaacagc	1380
atctccactc	attcgtctgt	cccaggatag	gcagtaaacat	attgggagctc	gggctgagaa	1440
caagaatcag	gaaaaagtta	gaggagcaca	aggagccgtc	acctctcatt	accgcgcagg	1500
acgtacaaga	agctaatgtg	gcagcccgat	agcgtaaggaa	ggtgcgtgaa	gccgagaggt	1560
tgcgcgcagc	tctaccacct	ttggcagctg	atggttgagg	gcccaactctg	gaagccgatg	1620
tcgcactgat	gttacaagag	gctggggcgc	gctcagtgga	gacacctgtg	ggcttgataa	1680
aggttaccag	ctacgtaggc	gaggacaaga	tcggctctta	cgtgtgtgct	tctccgcagg	1740
ctgtactcaa	gactgaaaaa	ttatcttgca	tccacctctc	cgtgaaacaa	gtcatagtga	1800
taacacacctc	tggccgaaaa	ggggcgttat	ccgtggaaac	ataccatggt	aaagttagtg	1860
tgccagcagg	ccagcaataa	cccgctcaag	actttcaagc	ctcgtagtga	agtgccacca	1920
ttgtgtacaa	cgaaactgtg	ttcgttaaaa	ggtacotgca	ccatatggcc	acacatggag	1980
gcgcgtgtaa	catgtatgaa	gaattattaca	aaactgtcaa	gccacagcag	cacgacggcg	2040
aataacctgta	cgacatcgac	aggaacaact	gcgtcaagaa	agaactagtc	actgggctag	2100
ggctccacag	cgagctgggt	gatctccct	tccatgaatt	cgctcacag	agtcgtgaga	2160
cacgaccagc	cgctcccttac	caagtaccaa	ccataggggg	gtatggcggt	ccaggatcag	2220
gcaagctctg	catcatataa	agcgacgtca	ccaaaaaaga	tctagtgggt	agcgccaaga	2280
aagaaaactg	tgcagaaatt	ataagggacg	tcaagaaaat	gaaagggtcg	gacgtcaatg	2340
ccagaactgt	ggactcagtg	ctcttgaatg	gatgcaaaac	cccgttagag	accctgtata	2400
ttgacgaagg	ttttgcttgt	catgcaggta	ctctcagagc	gctcatagcc	attataagac	2460
ctaaaaaggc	agtgctctgc	ggggatccca	aacagtgcgg	tttttttaac	atgatgtggc	2520
tgaagtgca	ttttaaccac	gagatttgca	cacaagctct	ccacaaaagc	attctctcgc	2580
gttgactaaa	atctgtgact	toggctcgtc	caacctgtgt	ttacgacaaa	aaaattgaga	2640
cgaagaaatc	gaaagagact	aagatttgta	ttgacctac	cggcagatcc	aaacctaaag	2700
aggaagctct	catcttcact	tgtttcagag	gggtgggtgaa	gcagttgcaa	atagattaca	2760
aaggcaacga	aataatgacg	gcagctgcct	ctcaagggtc	gaccctgaaa	gttgtgtatg	2820
cogtctggta	caagggtgaa	gaaaatctcc	tgtagcaccc	caactcgaaa	catgtgaagc	2880
tctactgac	cgcgacggag	gaccgcatcg	tgtgggaaac	actagccggc	gacctgtgga	2940
taaaaactac	gactgccaag	tacctgggga	atttcaactgc	ccagatagag	gagtggtcaa	3000
cagagcatga	tggcatcatg	agggacatct	tggagagacc	ggacctaac	gacgtcttcc	3060
agaataaggc	aaaactgtgt	tggggccaag	ctttagtgcc	ggtgctgaag	accgctggca	3120
tagacatgac	catctgaaca	tggaaacact	tggattattt	tgaaacggac	aaagctcaat	3180
gacagagat	agtatgaaac	caactatgcg	tgaggtctct	ttgactcgat	ctggactccg	3240
gtctattttc	tggaccacct	gttccgttat	ccattaggaa	taatactcgg	gataactccc	3300
cgtcgcttaa	cattgacagg	ctgaataaag	aagtggtccg	tcagctctct	cgcaggtacc	3360
cacaactccc	tcgggcagtg	gccactggaa	gagttctatg	catgaacact	gttgactcgc	3420
gcaattatga	tcccgcgata	aacctagtag	ctgtaaaacg	aagactgcct	catgctttag	3480
tctctcaacca	taatgaacac	ccacagagtg	acttttctct	attcgctgac	aaattgaagg	3540
gcgaagactg	ctcgttggtc	ggggaaaagt	tgtccgtccc	aggcgaatg	gttgactggg	3600
tgtcagaccg	gctcgaggct	accttcagag	ctcggtgga	tttaggcatt	ccagggtgat	3660

tgcccaata	tgacataata	tttgtaagt	tgaggacccc	atataaatat	catcactatc	3720
agcagtggtga	agaccatgcc	atataagctta	gcattgttgac	caagaaaagct	tgctctgcac	3780
tgatcccggt	cggaacctgt	gtcagcatag	gttatgtgta	cgtgcacagg	gccagcgaaa	3840
gcacatctgg	tgctatagcg	cggcagttca	agttttcccg	ggtatgcmaa	ccgaaaacct	3900
cacttggaaga	gcaggaagtt	ctgtttgtat	tcatttggtta	cgtatgcgaag	gcccgtaacg	3960
accaactcta	caagcttttca	tcaaccttga	ccaacattta	tacaggttcc	agactccacg	4020
agccgggatg	tgacacctca	tatcatgttg	tgccggggga	tattgccacg	gccaccgaag	4080
gagtgattat	aaatgtctgt	aacagcaaa	gacaaactgg	cggaggggtg	tgccgagcgc	4140
tgatataaga	attcccgtaa	agcttctgatt	tacagccgat	cgaaagttaga	aaagcgcgac	4200
tggtcaaaag	tgacagctaaa	catatcattc	atgcgtagg	accaaacttc	aacaaagtgt	4260
cggagggtga	aggtgacaaa	cagttggcag	aggcttatga	gtccatcgct	aagattgtca	4320
cgataacaaa	ttacaagtca	gtagcgattc	cactgttgct	cacggcgatc	ttttccggga	4380
acaaagatcg	actaacccaa	tcattgaaac	atttgctgac	agcttttagac	accactgatg	4440
cagatgtgac	catatactgc	agggacaaga	aatgggaaat	gactctcaag	gaagcagttg	4500
ctaggagaga	agcagtgagg	gagatatgca	tatccgacga	ctcttcagtg	acagaacctgg	4560
atgcagagct	ggtaggggtg	catccgaaga	gttcttttgg	tggaagggaag	ggctacacga	4620
caagcgatgg	caaaacttttc	tcattatttg	aagggaacca	gtttccaccg	gcggccagaag	4680
atatagcaga	aatttaagtc	atgtggcccg	ttgcaacgga	ggccaatcag	caggtatgca	4740
tgataactct	cggaaaagc	atgagcagta	ttaggtcgaa	atgcccctgc	gaagagtcgg	4800
aagcctcttc	accactcagc	acgctgcctt	gcttgtgcat	ccatgccatg	actccgaaaa	4860
gagtaacgag	ctcaaaagcc	tcacgtccag	aacaaattac	tggtgtgctca	tcttttccat	4920
tgccgaagta	tagaatcact	ggtgtgcaga	agatccaatg	ctcccagctc	atattgtttct	4980
cacccgaagt	gcctcggtat	attcatccaa	ggaagtattc	cgtggaaaaca	ccaccggtag	5040
acagagactcc	ggagccatcg	gcagagaaac	aatccacaga	ggggacacct	gaaccaaccac	5100
catctataac	cgaggatgag	accaggaacta	gaacgcctga	gccgatcctc	atcgaaagag	5160
aagaagagga	tgcataaagt	ttgctgtcag	atggcccgcg	caccacagtg	ctgcaagctg	5220
agcgagacat	tccggggcgc	ccctctgtat	ctagctcctc	ctggctccatt	cctcatgatc	5280
cggactttga	tgtaggcagt	ttatccatc	ttgacacctc	ggagggagct	agcgtgacca	5340
gcggggcaac	gtcagccgag	actaacctct	acttcgcaaa	gagtatggag	tttctggcgc	5400
gaacoggtg	tgccgctcga	acagtatcca	ggaacctccc	acatcccgct	cgcgcacaaa	5460
gaacacccgtc	acttgcaacc	agcagggcct	gctcgagaac	cagcctagtt	tcacccccgc	5520
cagggctgaa	taggggtgat	actagagagg	agctcgaggc	gcttaccacc	tcacgcactc	5580
ctagcaggtc	gggtctcgaga	accagcctgg	tctccaaacc	gccaggcgta	ataaggggtga	5640
ttacaagaga	ggagttttgag	gcgttctgtg	cacaacaaca	atgacgggtt	gatcgggggtg	5700
catacactct	ttctctcgac	accgggtcaag	ggcattttaca	acaaaatcca	gtaaggccaaa	5760
cggtgtctac	cgaaagtggg	ttggagagga	ccgaatttga	gatttctgat	ccccccgcgc	5820
tcgacacnaga	aaaangangaa	ttactacgca	agaaattaca	gttaaatccc	acactcgtcta	5880
acagaagcag	ataccagctcc	aggaaggttg	agaacatgaa	agccataaca	gctagacgta	5940
ttctcgaaag	cctaggagct	tatttgaagg	cagaaggaaa	agtggagtcg	taccgaacct	6000
tgcatcctgt	tcctttgtat	tcactcagtg	tgaacctgtg	cttttcaagc	cccaaggtcg	6060
cagtgggaag	ctgtaacgcc	atgttgaagg	agaactttcc	gactgtggct	tcttatctga	6120
ttttccaga	gtatcatgcc	tatttggaca	tggttgaagc	agcttcatgc	tgcttagaca	6180
ctgcgcagtt	ttgcctgcga	aagctgcgca	gctttccaaa	gaacacactc	tatttggaaac	6240
ccacaatacg	atccggcagtg	ccttcaagcg	tccagaacac	gctccagaaac	gtcctggcag	6300
ctgccacaaa	aagaatttgc	aatgtcacgc	aaatgagaga	attgcccgtga	ttggattcgg	6360
cggcctttaa	tgtggaaatgc	ttcaagaaat	atgctgttaa	taatatgaat	tggaagagct	6420
ttaagaaaaa	ccccctcagg	cttactgaag	aaaaagtggt	aaattacatt	accaaattaa	6480
aaggacacaaa	agctgtctgt	ctttttgcga	agacacataa	tttgaatatg	ttgcagagaca	6540
taccaatgga	caggtttgtga	atggacttaa	agagagacgt	gaaagtgcct	ccaggaacaa	6600
aacatactga	agaaacggccc	aaggtacagg	tgatccagcg	tgccgactcg	ctagcaaacag	6660
cgatctctgt	cggaaatccac	cgagagctgg	ttaggagatt	aaatgcggtc	ctgcttccga	6720
acattctatac	actgtttgat	atgtcggctg	aagactttga	cgctattata	gccagagact	6780
tcagctcgg	ggattgtggt	ctggaaaactg	acatcgctgc	gttttagaaa	agtgagagcg	6840
acgcatcggc	ctcgaccggc	ttaatgatct	tggaaagact	aggtgtggac	gcagagctgt	6900
tgacgctgat	tgaggcggtc	ttcgccgaaa	tttcatcaat	acatttgcgc	actaaaacta	6960

aattttaaatt	ogggagccatg	atgaaatctg	gaatgttctc	cacactgttt	gtgaacacag	7020
tcatataacat	tgtaatcgca	agcagagtg	tgagagaacg	gctaaccogga	tcacactgtg	7080
cagcattcat	tgagagtgac	aatatcgtga	aaggagtcga	atcggaacaa	ttaatggcag	7140
acaggtgcgc	cacctgggtg	aatatggaag	tcaagattat	agatgctgtg	gtggggcgaga	7200
aagcgcccta	tttctgtgga	gggtttattt	tgtgtgaact	cgtgacogcg	acagcgctgc	7260
gtgtggcgaga	ccccctaaaa	agggctgttta	agcttggcaa	acctctggca	gcagacgatg	7320
aacatgatga	tgacaggaga	agggcattgc	atgaagagtc	aacacgctgg	aacagcgatg	7380
tgtattcttc	agagctgtgc	aaggcagtag	aatacaagga	tgaacccctg	ggacattcca	7440
tcatagtatt	ggccatgact	actctageta	gcagtggtta	atcattcagc	tacctgagag	7500
gggcccctat	aactctctac	ggctaactcg	aatggactac	gacatagtct	agtcocggaa	7560
gatgagagtg	atggggatgc	agaggaattg	gccacaatgg	tggatatggg	gcaacctagg	7620
ctttgtggatg	ataataattt	gtagggtggg	ggggaaactg	aacttgtggg	tcacagtcta	7680
ttatggggta	cctgtgtgga	aagaagcaaa	aactactcta	ttctgtgcat	cagatgctaa	7740
agcatatgat	aaagaagtag	ataatgtctg	ggctacacat	gctctgtgac	ccacagaccg	7800
caacccacga	gaaatagttt	tggaaaatgt	aacagaaaa	tttaacatgt	ggaaaaatga	7860
catgtgtggat	cagatgcatg	aggatataat	cagtttatgg	gatcaaaagg	taaaaccatg	7920
tgtaaaagtg	ccccactct	gtgtcacttt	aaatgtgaca	aatgcacctg	cctacaataa	7980
tagactgcat	ggaagaatga	aaaattgtct	tttcaataca	accacagaga	taagagatga	8040
gaacacagaaa	cggtagtcac	ttttttataa	acctgatgta	gtgccactta	ataggagaga	8100
agagaataat	gggacaggag	agtatataat	aataaattgc	aattcctcaa	ccataacaca	8160
agcgtgtcca	aaggtcaact	ttgacccaat	toctatacat	tattgtgctc	cagctgggta	8220
tggattctcta	aagtgttaata	ataagacatt	caatggggca	ggacacatga	ataatgtcag	8280
cacagatcaaa	tgtacacatg	gaattatgoc	agtggtatca	actcaattac	tgttaaatgg	8340
tagcttagca	gaggaagaga	taataattag	atctgaaaa	ctgcaaaaac	atatcaaatg	8400
ataatattgct	caacttaata	aatctgtaga	aattgtgtgt	acaagaccca	acaataatca	8460
agaaaaaagt	ataagagtag	gaccaggaca	aacattctat	gcaacagggt	aaataattagg	8520
aaacataaaga	gaagcacatt	gtaacattag	taaaagttaac	tggaccagta	ctttagaaca	8580
ggtaaagaaa	aaattaaaag	aacactacaa	taagacaata	gaattttaacc	ccacctcagg	8640
aggggatcta	gaagttacaa	cacatagctt	taattgtaga	ggagaatttt	tctattgcaa	8700
tacacacaaa	ctgttttcaa	acaacagtag	tccaacacac	gaaaccaatca	cactcccatg	8760
caagataaaa	acaattataa	acatgtggca	gaaggtagga	cgagcaatgt	atgccctcc	8820
cattgaagga	aacataacat	gtaaatcaaa	tatcacagga	ctactattga	cagctgattg	8880
aggaagaagt	acaacaaatg	agatattoag	acogggagga	ggaaatatga	aggaacaatg	8940
gagaaagtga	ttatataaat	ataaagtgtg	agaaattgag	ccatttgggg	tagcacccac	9000
taaatcaaaa	aggagagtg	tggagagaga	aaaaagagca	gtgggactag	gagctgtact	9060
cctgtgggtc	ttgggagcag	cagggaagcac	tatgggcgcg	gcgtcaataa	cgcgtacoggt	9120
acaggccaga	caactgttgt	ctggatatgt	gcaacagcaa	agcaatttgc	tgagagctat	9180
agagggcgaa	cagcatatgt	tgcacactac	gtctctgggg	attaagcagc	tcagacaag	9240
agtcctggct	atagagagat	acctaaaaga	tcaacagctc	ctagggctct	ggggctgctc	9300
tggaaaaatc	atctgcacaa	ctgctgtgcc	tgggaactcc	agttggagta	ataaattcca	9360
agaagatatt	tgggtataca	tgacctggat	gcagtgggat	agagaaatta	gtaattacac	9420
aggccaacata	tattaggttac	ttgaaagactc	gcacaaacag	caggagaaaa	gtgaaaaaga	9480
tttatatgca	ttggacagtt	ggaaaaactt	gtggaaattg	tttaacataa	caaatgtggt	9540
gtggtatata	aaaattatca	tcatgatagt	agaggctgt	atagggttga	gaataatttt	9600
tgggtgaact	gctatagtag	aaagagttag	gcagggatac	tcacctttgt	cgtttcagac	9660
ccctaccccca	agccggaggg	gtcccgacag	gctcggaaga	atcgagaaga	aaggtggaga	9720
gcaagacaaa	gacagatcca	tctgattagt	gagcggattc	ttagcaacttg	ctgggacga	9780
ctctgcccag	ctgtgcctct	tccagctacca	ccacttgaga	gaactccat	tgattgcagc	9840
gagacgagcg	gaactctctg	gacgcagcag	tctcagggga	ctgcagagag	ggtgggaagc	9900
ccttaagtag	ctgggaaatc	tgtgtcagta	tgggggtctg	gagctaaaaa	gaagtgtgat	9960
taaaactgtt	gataccatag	caatagcagt	agctgaagag	acagatagga	ttcttgaagt	10020
aatacagaga	attgttagag	ctatccggca	catacctata	agaataagac	agggcttga	10080
agocagcttg	caataaataa	ttaagtaacc	gatacagcag	caattggcaa	gctgcttaca	10140
tagaactcgc	ggcgatggc	atgcgcgctt	aaaaatttta	ttttattttt	tttttttttt	10200
tccgaatogg	aaattgtttt	taattattca	aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	10260

aaaaaaaa	aaaaaaaa	gaagagcgcg	gcgcgcgcgc	gggctacgtc	ttgctggcgt	10320
tccgcatcgc	aggctggatg	gccttcccac	ttatgattct	tctcgcttcc	ggcgccatcg	10380
ggatgcgcgc	gttgcagccg	atgctgtcca	ggcaggtaga	tgacgaccat	caggcgacgc	10440
ttcaaggatc	gctcgcggct	cttaccagcc	taacttcgat	cacttgaccg	ctgatcgta	10500
cggcgattta	tgccgcctcg	gcgagcacat	ggaacgggtt	ggcatggatt	gtaggcgccg	10560
ccctaaccct	tgtctgcctc	cccgcggtgc	gtcgcgggtg	atggagccgg	gccacctcga	10620
ccatgaatgga	agccggcggc	acctcgctaa	cggattcacc	actccaagaa	ttgagcccaa	10680
tccattctct	cgggaactgt	tgaatgcgca	aaccaacctt	tggcagaaca	ttccatcgcc	10740
gtccgcacat	tccagcagcc	gcacgcggcg	catctcggcg	agcggtgggt	ccctggccacg	10800
ggctgcgcgc	atcgctgtcc	tgtcgttgag	gacccggcta	ggctggcggg	gtgccttac	10860
tggttagcag	aatgaatcac	cgatacgcga	gcgaacgtga	agcgactgct	gctgcaaaac	10920
gtctgcgcgc	tgagcaacaa	catgaatggt	cttcggttcc	cgtgtttcgt	aaagtctgag	10980
aaocggcgaa	tcagcgcctt	gcaccattat	gttcgggac	tgcatcgacg	gatgctcgtg	11040
gctaccctct	ggaaacaccta	catctgtatt	aacgaagcgc	tgccattgac	ccctgagtgt	11100
ttttctctctg	tcccgcgcga	tcataaccgc	cagttgttla	ccctcaaac	gttccagtaa	11160
cgggcatgct	tcatacatg	taaccgctat	cgtgagcatc	ctctctcgtt	tcactcggtat	11220
catctaccct	atgaacagaa	atccccctta	acaggaggca	tcagtgacca	aacagcaaaa	11280
aaocgcacct	aacatggccc	gctttatcag	aagccagaca	ttacgcttcc	tggaagaaat	11340
caacgagctg	gcgcgggatg	aacaggcaga	catctgtgaa	tcgcttccag	accacgctga	11400
tgagctttac	cgcagctgcc	tccgcgcggt	cggtgatgac	gggtgaaaa	ctgcacacat	11460
gcagctcccg	gagacggcca	cagcttgtct	gtaagcggat	gcocggagca	gacaagcccg	11520
tcaggggcgcg	tcacgggggtg	ttggcggggtg	tcggggcgca	gcgatgacc	agtcacgtag	11580
cgtatagcga	gtgtatactg	gcttaactat	gcggcatcag	agcagattgt	actgagagtg	11640
caccatatact	cggggtgtgaa	ataccgcaca	gatcgctaag	gagaaataac	gcactcaggc	11700
gctcttccgc	ttctcgcctc	actgactcgc	tgccctcggt	cgttcggctg	cggcgagcgg	11760
tatcagctca	ctcaaggcgc	gtaatacggc	tatccacaga	atcaggggat	aacgcaggaa	11820
agaaactatg	agcaaaaggc	cagcaaaagg	ccaggaaacg	taaaaaggcc	cggttgtctg	11880
cgtttttcca	taggtctcgc	cccctgacg	agcatcacaa	aaatcgacg	tcagtcaga	11940
gggtggcgaaa	cccgcacgga	ctataaagat	accaggcggt	tcccctcggg	agctccctcg	12000
tgcgctctcc	tgttccgacc	ctgcgcctta	ccggatagct	gtccgccttt	ctcccttcgg	12060
gaagcgtggc	gctttctcat	agctcaegct	gtaggatatc	cagttcggtg	ttagctgttc	12120
gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgacgcgtgc	gccttatccg	12180
gtaactatcg	tcttgagtc	aaccgggtaa	gacacgactt	atcgccactg	gcagcagcca	12240
ctggtaacag	gatttagcga	gcgaggtatg	taggcgggtg	tacagagttc	ttgaagtgtg	12300
ggcctaacta	cgcctacact	agaaggacag	tatttggtat	ctgcgctctg	ctggaagccg	12360
ttaccctcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	12420
gtgggtttttt	tggttgcaag	cagcagatta	cgcgacagaa	aaaaggatct	caagaagacc	12480
cttttgatctt	ttctacgggg	ttctgacgtc	atgggaacga	aaactcaggt	taagggtatt	12540
tggtcatgaa	caataaaact	gtctgcttac	ataaacagta	atacaagggg	tggttatgagc	12600
catattcaac	gggaacaagtc	ttgctcgagg	cgcgcattaa	atcccaact	ggatgctgat	12660
ttatatgggt	ataaatgggc	tcgcgataat	gtcgggcaat	caggtgcgac	aatctatcga	12720
ttgatgggga	agcccgatgc	gccagagtgt	ttctgaaac	atggcaaaag	tagcgttgcc	12780
aatgatgta	cagatgagat	ggtcagacta	aactggctga	cggaaatttat	gcctctccgc	12840
accatcaang	atttttccg	tactcctgat	gatgcaggtt	tactcaaac	tgcatcccc	12900
gggaaacag	cattccaggt	attagaagaa	tattcctggt	caggtgaaaa	tattgttgat	12960
gcgctggcag	tggttctcgc	ccggttgcct	tcgatctcgt	ttgttaattg	tccttttaac	13020
agcgatcgcg	tattttgctt	cgcgcaggcg	caatcacgaa	tgaataacgg	tttgtgtgat	13080
gcgagtgatt	ttgatgacga	gcgttaatggc	tgccctgttg	aacaagctcg	gaagaagatg	13140
catcaagctt	tgccatcttc	accggattca	gtcgtcacct	atgggtgatt	ctcacttgat	13200
aaacttattt	ttgacgaggg	gaatataata	ggttgatatt	atggttgacg	agtcggaatc	13260
gcagaccgat	accaggatct	tgccatccta	tggaactgct	ctcggtgagt	ttctcttcca	13320
ttacagaaac	ggctttttca	aaaatatggt	attgataatc	ctgatatgaa	taaattgcag	13380
ttactattga	tgctcagatga	gtttttctaa	gaattctcat	gtttgacagc	ttatcatoga	13440
taacgctttaa	tcgggtagtt	tatcacagtt	aaatgtctaa	cgcagctcag	caacggttat	13500
gaaattctaac	aatgcgctca	tctgcatcct	cggcaacgtc	accctggatg	ctgtctagag	13560

gacccctaata acgactcact atag

13584

<210> 18

<211> 2532

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence; Note =
synthetic construct

<221> CDS

<222> (1) ... (2532)

<400> 18

atg aga gtg atg ggg ata cag agg aat tgg cca caa tgg tgg ata tgg 48

Met Arg Val Met Gly Ile Gln Arg Asn Trp Pro Gln Trp Ile Trp

1

5

10

15

ggc acc tta ggc ttt tgg atg ata ata att tgt agg gtg gtg ggg aac 96

Gly Thr Leu Gly Phe Trp Met Ile Ile Cys Arg Val Val Gly Asn

20

25

30

ttg aac ttg tgg gtc aca gtc tat tat ggg gta cct gtg tgg aaa gaa 144

Leu Asn Leu Trp Val Thr Val Tyr Tyr Gly Val Pro Val Trp Lys Glu

35

40

45

gca aaa act act cta ttc tgt gca tca gat gct aaa gca tat gat aaa 192

Ala Lys Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys Ala Tyr Asp Lys

50

55

60

gaa gta cat aat gtc tgg gct aca cat gcc tgt gta ccc aca gac ccc 240

Glu Val His Asn Val Trp Ala Thr His Ala Cys Val Pro Thr Asp Pro

65

70

75

80

aac cca cga gaa ata gtt ttg gaa aat gta aca gaa aat ttt aac atg 288

Asn Pro Arg Glu Ile Val Leu Glu Asn Val Thr Glu Asn Phe Asn Met

85

90

95

tgg aaa aat gac atg gtg gat cag atg cat gag gat ata atc agt tta 336

Trp Lys Asn Asp Met Val Asp Gln Met His Glu Asp Ile Ile Ser Leu

100

105

110

tgg gat caa agc cta aaa cca tgt gta aag ttg acc cca ctc tgt gtc 384

Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr Pro Leu Cys Val

115

120

125

act tta aat tgt aca aat gca cct gcc tac aat aat agc atg cat gga 432

Thr Leu Asn Cys Thr Asn Ala Pro Ala Tyr Asn Asn Ser Met His Gly

130

135

140

gaa atg aaa aat tgc tct ttc aat aca acc aca gag ata aga gat agg 480

Glu Met Lys Asn Cys Ser Phe Asn Thr Thr Thr Glu Ile Arg Asp Arg

145

150

155

160

aaa cag aaa gcg tat gca ctt ttt tat aaa cct gat gta gtg cca ctt Lys Gln Lys Ala Tyr Ala Leu Phe Tyr Lys Pro Asp Val Val Pro Leu 165 170 175	528
aat agg aga gaa gag aat aat ggg aca gga gag tat ata tta ata aat Asn Arg Arg Glu Glu Asn Asn Gly Thr Gly Glu Tyr Ile Leu Ile Asn 180 185 190	576
tgc aat tcc tca acc ata aca caa gcc tgt cca aag gtc act ttt gac Cys Asn Ser Ser Thr Ile Thr Gln Ala Cys Pro Lys Val Thr Phe Asp 195 200 205	624
cca att cct ata cat tat tgt gct cca gct ggt tat gcg att cta aag Pro Ile Pro Ile His Tyr Cys Ala Pro Ala Gly Tyr Ala Ile Leu Lys 210 215 220	672
tgt aat aat aag aca ttc aat ggg aca gga cca tgc aat aat gtc agc Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys Asn Asn Val Ser 225 230 235 240	720
aca gta caa tgt aca cat gga att atg cca gtg gta tca act caa tta Thr Val Gln Cys Thr His Gly Ile Met Pro Val Val Ser Thr Gln Leu 245 250 255	768
ctg tta aat ggt agc cta gca gaa gaa gag ata ata att aga tct gaa Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Ile Ile Ile Arg Ser Glu 260 265 270	816
aat ctg aca aac aat atc aaa aca ata ata gtc cac ctt aat aaa tct Asn Leu Thr Asn Asn Ile Lys Thr Ile Ile Val His Leu Asn Lys Ser 275 280 285	864
gta gaa att gtg tgt aca aga ccc aac aat aat aca aga aaa agt ata Val Glu Ile Val Cys Thr Arg Pro Asn Asn Asn Thr Arg Lys Ser Ile 290 295 300	912
agg ata gga cca gga caa aca ttc tat gca aca ggt gaa ata ata gga Arg Ile Gly Pro Gly Gln Thr Phe Tyr Ala Thr Gly Glu Ile Ile Gly 305 310 315 320	960
aac ata aga gaa gca cat tgt aac att agt aaa agt aac tgg acc agt Asn Ile Arg Glu Ala His Cys Asn Ile Ser Lys Ser Asn Trp Thr Ser 325 330 335	1008
act tta gaa cag gta aag aaa aaa tta aaa gaa cac tac aat aag aca Thr Leu Glu Gln Val Lys Lys Lys Leu Lys Glu His Tyr Asn Lys Thr 340 345 350	1056
ata gaa ttt aac cca ccc tca gga ggg gat cta gaa gtt aca aca cat Ile Glu Phe Asn Pro Pro Ser Gly Gly Asp Leu Glu Val Thr Thr His 355 360 365	1104
agc ttt aat tgt aga gga gaa ttt ttc tat tgc aat aca aca aaa ctg Ser Phe Asn Cys Arg Gly Glu Phe Phe Tyr Cys Asn Thr Thr Lys Leu 370 375 380	1152

ttt tca aac aac agt gat tca aac aac gaa acc atc aca ctc cca tgc Phe Ser Asn Asn Ser Asp Ser Asn Asn Glu Thr Ile Thr Leu Pro Cys 385 390 395 400	1200
aag ata aaa caa att ata aac atg tgg cag aag gta gga cga gca atg Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Arg Ala Met 405 410 415	1248
tat gcc cct ccc att gaa gga aac ata aca tgt aaa tca aat atc aca Tyr Ala Pro Pro Ile Glu Gly Asn Ile Thr Cys Lys Ser Asn Ile Thr 420 425 430	1296
gga cta cta ttg aca cgt gat gga gga aag aat aca aca aat gag ata Gly Leu Leu Leu Thr Arg Asp Gly Gly Lys Asn Thr Thr Asn Glu Ile 435 440 445	1344
ttc aga cgg gga gga gga aat atg aag gac aat tgg aga agt gaa tta Phe Arg Pro Gly Gly Gly Asn Met Lys Asp Asn Trp Arg Ser Glu Leu 450 455 460	1392
tat aaa tat aaa gtg gta gaa att gag cca ttg gga gta gca ccc act Tyr Lys Tyr Lys Val Val Glu Ile Glu Pro Leu Gly Val Ala Pro Thr 465 470 475 480	1440
aaa tca aaa agg aga gtg gtg gag aga gaa aaa aga gca gtg gga cta Lys Ser Lys Arg Arg Val Val Glu Arg Glu Lys Arg Ala Val Gly Leu 485 490 495	1488
gga gct gta ctc ctt ggg ttc ttg gga gca gca gga agc act atg ggc Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly 500 505 510	1536
gcg gcg tca ata acg ctg acg gta cag gcc aga caa ctg ttg tct ggt Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly 515 520 525	1584
ata gtg caa cag caa agc aat ttg ctg aga gct ata gag gcg caa cag Ile Val Gln Gln Gln Ser Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln 530 535 540	1632
cat atg ttg caa ctc acg gtc tgg ggc att aag cag ctc cag aca aga His Met Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Thr Arg 545 550 555 560	1680
gtc ttg gct ata gag aga tac cta aag gat caa cag ctc cta ggg ctt Val Leu Ala Ile Glu Arg Tyr Leu Lys Asp Gln Gln Leu Leu Gly Leu 565 570 575	1728
tgg ggc tgc tct gga aaa atc atc tgc acc act gct gtg cct tgg aac Trp Gly Cys Ser Gly Lys Ile Ile Cys Thr Thr Ala Val Pro Trp Asn 580 585 590	1776
tcc agt tgg agt aat aaa tct caa gaa gat att tgg gat aac atg acc Ser Ser Trp Ser Ser Asn Lys Ser Gln Glu Asp Ile Trp Asp Asn Met Thr 595 600 605	1824

tgg atg cag tgg gat aga gaa att agt aat tac aca ggc aca ata tat Trp Met Gln Trp Asp Arg Glu Ile Ser Asn Tyr Thr Gly Thr Ile Tyr 610 615 620	1872
agg tta ctt gaa gac tgg caa aac cag cag gag aaa aat gaa aaa gat Arg Leu Leu Glu Asp Ser Gln Asn Gln Gln Glu Lys Asn Glu Lys Asp 625 630 635 640	1920
tta tta gca ttg gac agt tgg aaa aac ttg tgg aat tgg ttt aac ata Leu Leu Ala Leu Asp Ser Trp Lys Asn Leu Trp Asn Trp Phe Asn Ile 645 650 655	1968
aca aat tgg ctg tgg tat ata aaa ata ttc atc atg ata gta gga ggc Thr Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly 660 665 670	2016
ttg ata ggt ttg aga ata att ttt ggt gta ctc gct ata gtg aaa aga Leu Ile Gly Leu Arg Ile Ile Phe Gly Val Leu Ala Ile Val Lys Arg 675 680 685	2064
gtt agg cag gga tac tca cct ttg tgg ttt cag acc ctt acc cca agc Val Arg Gln Gly Tyr Ser Pro Leu Ser Phe Gln Thr Leu Thr Pro Ser 690 695 700	2112
cgg agg ggt ccc gac agg ctc gga aga atc gaa gaa gaa ggt gga gag Pro Arg Gly Pro Asp Arg Leu Gly Arg Ile Glu Glu Glu Gly Gly Glu 705 710 715 720	2160
caa gac aaa gac aga tcc att cga tta gtg agc gga ttc tta gca ctt Gln Asp Lys Asp Arg Ser Ile Arg Leu Val Ser Gly Phe Leu Ala Leu 725 730 735	2208
gcc tgg gac gat ctg cgg agc ctg tgc ctc ttc agc tac cac cac ttg Ala Trp Asp Asp Leu Arg Ser Leu Cys Leu Phe Ser Tyr His His Leu 740 745 750	2256
aga gac ttc ata ttg att gca gcg aga gca gcg gaa ctt ctg gga cgc Arg Asp Phe Ile Leu Ile Ala Ala Arg Ala Ala Glu Leu Leu Gly Arg 755 760 765	2304
agc agt ctc agg gga ctg cag aga ggg tgg gaa gcc ctt aag tat ctg Ser Ser Leu Arg Gly Leu Gln Arg Gly Trp Glu Ala Leu Lys Tyr Leu 770 775 780	2352
gga aat ctt gtg cag tat ggg ggt ctg gag cta aaa aga agt gct att Gly Asn Leu Val Gln Tyr Gly Gly Leu Glu Leu Lys Arg Ser Ala Ile 785 790 795 800	2400
aaa ctg ttt gat acc ata gca ata gca gta gct gaa gga aca gat agg Lys Leu Phe Asp Thr Ile Ala Ile Ala Val Ala Glu Gly Thr Asp Arg 805 810 815	2448
att ctt gaa gta ata cag aga att tgt aga gct atc cgc cac ata cct Ile Leu Glu Val Ile Gln Arg Ile Cys Arg Ala Ile Arg His Ile Pro 820 825 830	2496

ata aga ata aga cag ggc ttt gaa gca gct ttg caa
 ile arg ile arg gln gly phe glu ala ala leu gln
 835 840

2532

<210> 19
 <211> 844
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence; Note =
 synthetic construct

<400> 19
 Met Arg Val Met Gly Ile Gln Arg Asn Trp Pro Gln Trp Trp Ile Trp
 1 5 10 15
 Gly Thr Leu Gly Phe Trp Met Ile Ile Ile Cys Arg Val Val Gly Asn
 20 25 30
 Leu Asn Leu Trp Val Thr Val Tyr Tyr Gly Val Pro Val Trp Lys Glu
 35 40 45
 Ala Lys Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys Ala Tyr Asp Lys
 50 55 60
 Glu Val His Asn Val Trp Ala Thr His Ala Cys Val Pro Thr Asp Pro
 65 70 75 80
 Asn Pro Arg Glu Ile Val Leu Glu Asn Val Thr Glu Asn Phe Asn Met
 85 90 95
 Trp Lys Asn Asp Met Val Asp Gln Met His Glu Asp Ile Ile Ser Leu
 100 105 110
 Trp Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr Pro Leu Cys Val
 115 120 125
 Thr Leu Asn Cys Thr Asn Ala Pro Ala Tyr Asn Asn Ser Met His Gly
 130 135 140
 Glu Met Lys Asn Cys Ser Phe Asn Thr Thr Thr Glu Ile Arg Asp Arg
 145 150 155 160
 Lys Gln Lys Ala Tyr Ala Leu Phe Tyr Lys Pro Asp Val Val Pro Leu
 165 170 175
 Asn Arg Arg Glu Glu Asn Asn Gly Thr Gly Glu Tyr Ile Leu Ile Asn
 180 185 190
 Cys Asn Ser Ser Thr Ile Thr Gln Ala Cys Pro Lys Val Thr Phe Asp
 195 200 205
 Pro Ile Pro Ile His Tyr Cys Ala Pro Ala Gly Tyr Ala Ile Leu Lys
 210 215 220
 Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys Asn Asn Val Ser
 225 230 235 240
 Thr Val Gln Cys Thr His Gly Ile Met Pro Val Val Ser Thr Gln Leu
 245 250 255
 Leu Leu Asn Gly Ser Leu Ala Glu Glu Glu Ile Ile Arg Ser Glu
 260 265 270
 Asn Leu Thr Asn Asn Ile Lys Thr Ile Ile Val His Leu Asn Lys Ser
 275 280 285
 Val Glu Ile Val Cys Thr Arg Pro Asn Asn Asn Thr Arg Lys Ser Ile
 290 295 300

Arg Ile Gly Pro Gly Gln Thr Phe Tyr Ala Thr Gly Glu Ile Ile Gly
 305 310 315 320
 Asn Ile Arg Glu Ala His Cys Asn Ile Ser Lys Ser Asn Trp Thr Ser
 325 330 335
 Thr Leu Glu Gln Val Lys Lys Lys Leu Lys Glu His Tyr Asn Lys Thr
 340 345 350
 Ile Glu Phe Asn Pro Pro Ser Gly Gly Asp Leu Glu Val Thr Thr His
 355 360 365
 Ser Phe Asn Cys Arg Gly Glu Phe Phe Tyr Cys Asn Thr Thr Lys Leu
 370 375 380
 Phe Ser Asn Asn Ser Asp Ser Asn Asn Glu Thr Ile Thr Leu Pro Cys
 385 390 395 400
 Lys Ile Lys Gln Ile Ile Asn Met Trp Gln Lys Val Gly Arg Ala Met
 405 410 415
 Tyr Ala Pro Pro Ile Glu Gly Asn Ile Thr Cys Lys Ser Asn Ile Thr
 420 425 430
 Gly Leu Leu Leu Thr Arg Asp Gly Gly Lys Asn Thr Thr Asn Glu Ile
 435 440 445
 Phe Arg Pro Gly Gly Gly Asn Met Lys Asp Asn Trp Arg Ser Glu Leu
 450 455 460
 Tyr Lys Tyr Lys Val Val Glu Ile Glu Pro Leu Gly Val Ala Pro Thr
 465 470 475 480
 Lys Ser Lys Arg Arg Val Val Glu Arg Glu Lys Arg Ala Val Gly Leu
 485 490 495
 Gly Ala Val Leu Leu Gly Phe Leu Gly Ala Ala Gly Ser Thr Met Gly
 500 505 510
 Ala Ala Ser Ile Thr Leu Thr Val Gln Ala Arg Gln Leu Leu Ser Gly
 515 520 525
 Ile Val Gln Gln Gln Ser Asn Leu Leu Arg Ala Ile Glu Ala Gln Gln
 530 535 540
 His Met Leu Gln Leu Thr Val Trp Gly Ile Lys Gln Leu Gln Thr Arg
 545 550 555 560
 Val Leu Ala Ile Glu Arg Tyr Leu Lys Asp Gln Gln Leu Leu Gly Leu
 565 570 575
 Trp Gly Cys Ser Gly Lys Ile Ile Cys Thr Thr Ala Val Pro Trp Asn
 580 585 590
 Ser Ser Trp Ser Asn Lys Ser Gln Glu Asp Ile Trp Asp Asn Met Thr
 595 600 605
 Trp Met Gln Trp Asp Arg Glu Ile Ser Asn Tyr Thr Gly Thr Ile Tyr
 610 615 620
 Arg Leu Leu Glu Asp Ser Gln Asn Gln Gln Glu Lys Asn Glu Lys Asp
 625 630 635 640
 Leu Leu Ala Leu Asp Ser Trp Lys Asn Leu Trp Asn Trp Phe Asn Ile
 645 650 655
 Thr Asn Trp Leu Trp Tyr Ile Lys Ile Phe Ile Met Ile Val Gly Gly
 660 665 670
 Leu Ile Gly Leu Arg Ile Ile Phe Gly Val Leu Ala Ile Val Lys Arg
 675 680 685
 Val Arg Gln Gly Tyr Ser Pro Leu Ser Phe Gln Thr Leu Thr Pro Ser
 690 695 700
 Pro Arg Gly Pro Asp Arg Leu Gly Arg Ile Glu Glu Glu Gly Gly Glu
 705 710 715 720
 Gln Asp Lys Asp Arg Ser Ile Arg Leu Val Ser Gly Phe Leu Ala Leu
 725 730 735

Ala	Trp	Asp	Asp	Leu	Arg	Ser	Leu	Cys	Leu	Phe	Ser	Tyr	His	His	Leu
			740					745					750		
Arg	Asp	Phe	Ile	Leu	Ile	Ala	Ala	Arg	Ala	Ala	Glu	Leu	Leu	Gly	Arg
		755					760					765			
Ser	Ser	Leu	Arg	Gly	Leu	Gln	Arg	Gly	Trp	Glu	Ala	Leu	Lys	Tyr	Leu
		770				775					780				
Gly	Asn	Leu	Val	Gln	Tyr	Gly	Gly	Leu	Glu	Leu	Lys	Arg	Ser	Ala	Ile
785					790					795					800
Lys	Leu	Phe	Asp	Thr	Ile	Ala	Ile	Ala	Val	Ala	Glu	Gly	Thr	Asp	Arg
			805						810						815
Ile	Leu	Glu	Val	Ile	Gln	Arg	Ile	Cys	Arg	Ala	Ile	Arg	His	Ile	Pro
		820						825					830		
Ile	Arg	Ile	Arg	Gln	Gly	Phe	Glu	Ala	Ala	Leu	Gln				
		835					840								